

***-homomorphisms between groupoid C^* -algebras**

Fuyuta Komura

Abstract. In this paper, we investigate *-homomorphisms between C^* -algebras associated to étale groupoids. First, we prove that such a *-homomorphism can be described by closed invariant subsets, groupoid homomorphisms and cocycles under some assumptions. Then we prove C^* -rigidity results for étale groupoids which are not necessarily effective. As another application, we investigate certain subgroups of the automorphism groups of groupoid C^* -algebras. More precisely, we show that the groups of automorphisms that globally preserve the function algebras on the unit spaces are isomorphic to certain semidirect product groups. As a corollary, we show that, if group actions on groupoid C^* -algebras fix the function algebras on the unit spaces, then the actions factors through the abelianizations of the acting groups.

1. Introduction

The main subject in the present paper is C^* -algebras associated with étale groupoids, that is, groupoid C^* -algebras. The theory of groupoid C^* -algebras is initiated by Renault in [16]. It is known that many C^* -algebras are realized as groupoid C^* -algebras (see [18], for example). It is a natural task to characterize properties of groupoid C^* -algebras in terms of étale groupoids. For instance, see [1] for the relation between nuclearity of groupoid C^* -algebras and amenability of topological groupoids. In addition, simplicity of full groupoid C^* -algebras is investigated in [2]. Recently, the authors in [3] established the Galois correspondence result between étale groupoids and twisted groupoid C^* -algebras. In [11], the author studied certain submodules in groupoid C^* -algebras and analyzed discrete group coactions on groupoid C^* -algebras. In the present paper, we investigate *-homomorphisms between groupoid C^* -algebras. First of all, we explain our motivation to study *-homomorphisms between groupoid C^* -algebras.

For an étale groupoid G , one can construct a (reduced) groupoid C^* -algebra $C_r^*(G)$ and a commutative C^* -subalgebra $C_0(G^{(0)}) \subset C_r^*(G)$. Renault proved a C^* -rigidity result in [17]. Namely, for effective étale groupoids G_1 and G_2 , Renault proved that G_1 and G_2 are isomorphic as étale groupoids if inclusions of C^* -algebras $C_0(G_1^{(0)}) \subset C_r^*(G_1)$ and $C_0(G_2^{(0)}) \subset C_r^*(G_2)$ are isomorphic. This result connects the bridge between étale groupoids and C^* -algebras and has various applications. Indeed, one can deduce a C^* -rigidity result for C^* -algebras associated with dynamical systems from Renault's

result (see [13], for example). Now, assume that inclusions $C_0(G_1^{(0)}) \subset C_r^*(G_1)$ and $C_0(G_2^{(0)}) \subset C_r^*(G_2)$ are isomorphic. One may wonder how many isomorphisms between $C_0(G_1^{(0)}) \subset C_r^*(G_1)$ and $C_0(G_2^{(0)}) \subset C_r^*(G_2)$ exist. To solve this problem, it is sufficient to determine the following group

$$\text{Aut}_{C_0(G_1^{(0)})}(C_r^*(G_1)) := \{\varphi \in \text{Aut}(C_r^*(G_1)) \mid \varphi(C_0(G_1^{(0)})) = C_0(G_1^{(0)})\}.$$

Therefore, we are motivated to investigate $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$ for an effective étale groupoid G . In Corollary 3.2.2, we prove that $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$ is isomorphic to the semidirect group of $\text{Aut}(G)$ and $Z(G, \mathbb{T})$, where $Z(G, \mathbb{T})$ is the abelian group of \mathbb{T} -valued 1-cocycles on G . We remark that the almost same result is obtained in [14, Proposition 5.7 (1)] and we obtain a slightly more direct proof by using our main theorem, that is, Theorem 3.1.1. In addition, we shall remark that the similar result has been already obtained for von Neumann algebras arising from equivalence relations in [8, Theorem 3]. In the present paper, we deal with an analogue of [8, Theorem 3] for groupoid C^* -algebras.

As stated above, our purpose in this paper is to study $*$ -homomorphisms between groupoid C^* -algebras such as elements in $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$. For example, a similar attempt succeeded in [5, Theorem 6.3] for isomorphisms between regular C^* -inclusions in terms of coordinate systems. In this paper, we begin with the study of general $*$ -homomorphisms between groupoid C^* -algebras which need not to be isomorphisms. Our main theorem is Theorem 3.1.1, which asserts that a $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ can be described in terms of underlying étale groupoids G and H under assumptions that H is effective and φ has some compatibility with $C_0(G^{(0)})$ and $C_0(H^{(0)})$. Taking into account that previous works like [17] relies on the effectiveness of underlying étale groupoids, it seems noteworthy that we do not assume the effectiveness of G . As a direct application of Theorem 3.1.1, we prove that surjective $*$ -homomorphisms between groupoid C^* -algebras induce quotients of étale groupoids (Corollary 3.1.11). Corollary 3.1.11 generalizes Renault's result in [17], which asserts that $*$ -isomorphisms between groupoid C^* -algebras which preserve the Cartan subalgebras induce isomorphisms between étale groupoids. From Corollary 3.1.11, we obtain the following variant of the rigidity results for not necessarily effective étale groupoids: for étale (not necessarily effective) groupoids G which have closed $\text{Iso}(G)^\circ$ and some amenability condition, the quotient groupoids $G/\text{Iso}(G)^\circ$ are invariants for the inclusion of C^* -algebras $(C_r^*(G), C_0(G^{(0)}))$ (Corollary 3.1.12). In other words, $(C_r^*(G), C_0(G^{(0)}))$ remembers $G/\text{Iso}(G)^\circ$ even if an étale groupoid G is not effective.

As another important application of Theorem 3.1.1, we prove the structure theorem of $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$ (Corollary 3.2.2). More precisely, we show that $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$ is isomorphic to the semidirect product of $\text{Aut}(G)$ and $Z(G, \mathbb{T})$. In particular, it turns out that $Z(G, \mathbb{T})$ corresponds to a certain abelian subgroup of $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$. In Corollary 3.2.6, we prove that $Z(G, \mathbb{T})$ corresponds to

$$\text{FAut}_{C_0(G^{(0)})}(C_r^*(G)) := \{\varphi \in \text{Aut}(C_r^*(G)) \mid \varphi(a) = a \text{ for all } a \in C_0(G^{(0)})\}.$$

As a corollary, we show that a group action on a groupoid C^* -algebra factors through its abelianization if the fixed point subalgebra contains $C_0(G^{(0)})$ (Corollary 3.2.8).

This paper is organized as follows. In Section 1, we recall fundamental facts about étale groupoids, groupoid C^* -algebras and inverse semigroups. In Section 2, we prove our main theorems about *-homomorphisms between groupoid C^* -algebras. Our goal in the first subsection, Section 3.1, is Theorem 3.1.1. Toward Theorem 3.1.1, we first prove that *-homomorphisms between groupoid C^* -algebras induce groupoid homomorphisms and \mathbb{T} -valued 1 cocycles (Lemma 3.1.6 and Lemma 3.1.8). Then we prove that every *-homomorphism can be described by closed invariant subsets, groupoid homomorphisms and \mathbb{T} -valued 1-cocycles (Proposition 3.1.9 and Proposition 3.1.10). As a special case of Theorem 3.1.1, we observe that surjective *-homomorphisms induce quotients of étale groupoids in Corollary 3.1.11. This immediately implies a variant of rigidity result for not necessarily effective groupoids (Corollary 3.1.12). In Section 3.2, we investigate $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$ for an effective étale groupoid G . First, we prove that $\text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$ is isomorphic to the natural semidirect product of $\text{Aut}(G)$ and $Z(G, \mathbb{T})$ (Corollary 3.2.2). Then we observe that $Z(G, \mathbb{T})$ corresponds to $\text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$ (Corollary 3.2.6). In addition, we show that an endomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(G)$ which fixes $C_0(G^{(0)})$ pointwisely becomes an automorphism automatically (Corollary 3.2.7). As a by-product of the analysis of $\text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$, we show that a group action on a groupoid C^* -algebra factors through its abelianization if the fixed point algebra contains $C_0(G^{(0)})$ (Corollary 3.2.8).

2. Preliminaries

In this section, we recall fundamental notions about étale groupoids, groupoid C^* -algebras and inverse semigroups.

2.1. Étale groupoids

We recall the basic notions on étale groupoids. See [15, 18] for more details.

A groupoid is a set G together with a distinguished subset $G^{(0)} \subset G$, domain and range maps $d, r: G \rightarrow G^{(0)}$ and a multiplication

$$G^{(2)} := \{(\alpha, \beta) \in G \times G \mid d(\alpha) = r(\beta)\} \ni (\alpha, \beta) \mapsto \alpha\beta \in G$$

such that

- (1) for all $x \in G^{(0)}$, $d(x) = x$ and $r(x) = x$ hold,
- (2) for all $\alpha \in G$, $\alpha d(\alpha) = r(\alpha)\alpha = \alpha$ holds,
- (3) for all $(\alpha, \beta) \in G^{(2)}$, $d(\alpha\beta) = d(\beta)$ and $r(\alpha\beta) = r(\alpha)$ hold,
- (4) if $(\alpha, \beta), (\beta, \gamma) \in G^{(2)}$, we have $(\alpha\beta)\gamma = \alpha(\beta\gamma)$,
- (5) every $\gamma \in G$, there exists $\gamma' \in G$ which satisfies $(\gamma', \gamma), (\gamma, \gamma') \in G^{(2)}$ and $d(\gamma) = \gamma'\gamma$ and $r(\gamma) = \gamma\gamma'$.

Since the element γ' in (5) is uniquely determined by γ , γ' is called the inverse of γ and denoted by γ^{-1} . We call $G^{(0)}$ the unit space of G . A subgroupoid of G is a subset of G which is closed under the inversion and multiplication. For $U \subset G^{(0)}$, we define $G_U := d^{-1}(U)$ and $G^U := r^{-1}(U)$. We define also $G_x := G_{\{x\}}$ and $G^x := G^{\{x\}}$ for $x \in G^{(0)}$. A subset $F \subset G^{(0)}$ is said to be invariant if $d(\alpha) \in F$ implies $r(\alpha) \in F$ for all $\alpha \in G$. If $F \subset G^{(0)}$ is invariant, $G_F \subset G$ is a subgroupoid and the unit space of G_F is F .

A topological groupoid is a groupoid equipped with a topology where the multiplication and the inverse are continuous. A topological groupoid is said to be étale if the domain map is a local homeomorphism. Note that the range map of an étale groupoid is also a local homeomorphism. In this paper, although there exist important étale groupoids that are not Hausdorff, we assume that étale groupoids are always locally compact Hausdorff unless otherwise stated. Hence, we mean locally compact Hausdorff étale groupoids by étale groupoids.

A subset U of an étale groupoid G is called a bisection if the restrictions $d|_U$ and $r|_U$ are injective. It follows that $d|_U$ and $r|_U$ are homeomorphism onto their images if U is a bisection since d and r are open maps.

An étale groupoid G is said to be effective if $G^{(0)}$ coincides with the interior of $\text{Iso}(G)$, where

$$\text{Iso}(G) := \{\alpha \in G \mid d(\alpha) = r(\alpha)\}$$

is the isotropy of G . An étale groupoid G is said to be topologically principal if

$$\{x \in G^{(0)} \mid G_x \cap G^x = \{x\}\}$$

is dense in $G^{(0)}$. If G is topologically principal, then G is effective. If G is second countable and effective, then G is topologically principal (see [17, Proposition 3.6]).

A groupoid homomorphism $c: G \rightarrow \mathbb{T}$ is called a \mathbb{T} -valued 1-cocycle, where \mathbb{T} denotes the circle group. Because we only consider a \mathbb{T} -valued 1-cocycle in this paper, we often simply call it a cocycle. We let $Z(G, \mathbb{T})$ denote the set of all continuous cocycles $c: G \rightarrow \mathbb{T}$. Then $Z(G, \mathbb{T})$ is an abelian group with respect to the pointwise product.

2.2. Groupoid C^* -algebras

We recall the definition of groupoid C^* -algebras.

Let G be an étale groupoid. Then $C_c(G)$, the vector space of compactly supported continuous \mathbb{C} -valued functions on G , is a $*$ -algebra with respect to the multiplication and the involution defined by

$$f * g(\gamma) := \sum_{\alpha\beta=\gamma} f(\alpha)g(\beta), \quad f^*(\gamma) := \overline{f(\gamma^{-1})},$$

where $f, g \in C_c(G)$ and $\gamma \in G$. The left regular representation $\lambda_x: C_c(G) \rightarrow B(\ell^2(G_x))$ at $x \in G^{(0)}$ is defined by

$$\lambda_x(f)\delta_\alpha := \sum_{\beta \in G_{r(\alpha)}} f(\beta)\delta_{\beta\alpha},$$

where $f \in C_c(G)$ and $\alpha \in G_x$. The reduced norm $\|\cdot\|_r$ on $C_c(G)$ is defined by

$$\|f\|_r := \sup_{x \in G^{(0)}} \|\lambda_x(f)\|$$

for $f \in C_c(G)$. We often omit the subscript ‘ r ’ of $\|\cdot\|_r$ if there is no chance to confuse. The reduced groupoid C^* -algebra $C_r^*(G)$ is defined to be the completion of $C_c(G)$ with respect to the reduced norm. Note that $C_c(G^{(0)}) \subset C_c(G)$ is a $*$ -subalgebra and this inclusion extends to the inclusion $C_0(G^{(0)}) \subset C_r^*(G)$.

For a closed invariant subset $F \subset G^{(0)}$, the closed subgroupoid $G_F \subset G$ is étale with respect to the relative topology. It is well known that the restriction

$$C_c(G) \ni f \mapsto f|_{G_F} \in C_c(G_F)$$

extends to the surjective $*$ -homomorphism $C_r^*(G) \rightarrow C_r^*(G_F)$. In addition, the reduced groupoid C^* -algebra $C_r^*(G)$ can be embedded into $C_0(G)$ as in the following, which is originally proved by Renault in [16, Proposition II 4.2]. See also [18, Proposition 9.3.3] for the proof.

Proposition 2.2.1 (Evaluation). *Let G be an étale groupoid. For $a \in C_r^*(G)$, $j(a) \in C_0(G)$ is defined by*

$$j(a)(\alpha) := \langle \delta_\alpha | \lambda_{d(\alpha)}(a) \delta_{d(\alpha)} \rangle$$

for $\alpha \in G$.¹ Then $j: C_r^(G) \rightarrow C_0(G)$ is a norm decreasing injective linear map. Moreover, j is an identity map on $C_c(G)$.*

Remark 2.2.2. Since $j: C_r^*(G) \rightarrow C_0(G^{(0)})$ is injective, we may identify $j(a)$ with a . Hence, we often regard a as a function on G and simply denote $j(a)$ by a .

Finally, we recall facts about normalizers.

Definition 2.2.3. Let A be a C^* -algebra and $D \subset A$ be a C^* -subalgebra. An element $n \in A$ is called a normalizer for D if $nDn^* \cup n^*Dn \subset D$ holds. We denote the set of normalizers for D by $N(A, D)$.

For $a \in C_r^*(G)$, we denote the open support of a by

$$\text{supp}^\circ(a) := \{\alpha \in G \mid a(\alpha) \neq 0\}.$$

Note that $\text{supp}^\circ(a)$ is open in G . Normalizers for $C_0(G^{(0)})$ and bisections in G are intimately related as follows.

Proposition 2.2.4 ([17, Proposition 4.8]). *Let G be an étale groupoid and $U \subset G$ be an open set. If U is a bisection, then every elements in $C_c(U)$ is a normalizer. Moreover, if $n \in C_r^*(G)$ is a normalizer and G is effective, then $\text{supp}^\circ(n) \subset G$ is an open bisection.*

¹In this paper, inner products of Hilbert spaces are linear with respect to the right variables.

2.3. Inverse semigroups

We recall the basic notions about inverse semigroups. See [12] or [15] for more details. An inverse semigroup S is a semigroup where for every $s \in S$ there exists a unique $s^* \in S$ such that $s = ss^*s$ and $s^* = s^*ss^*$. We denote the set of all idempotents in S by $E(S) := \{e \in S \mid e^2 = e\}$. It is known that $E(S)$ is a commutative subsemigroup of S . An inverse semigroup which consists of idempotents is called a (meet) semilattice of idempotents. A zero element is a unique element $0 \in S$ such that $0s = s0 = 0$ holds for all $s \in S$. An inverse semigroup with a unit is called an inverse monoid. By a subsemigroup of S , we mean a subset of S that is closed under the product and inverse of S . A map $\varphi: S \rightarrow T$ between inverse semigroups S and T is called a semigroup homomorphism if $\varphi(st) = \varphi(s)\varphi(t)$ holds for all $s, t \in S$. Note that a semigroup homomorphism automatically preserves generalized inverses (i.e., $\varphi(s^*) = \varphi(s)^*$ holds for all $s \in S$).

For a topological space X , we denote by I_X the set of all homeomorphisms between open sets in X . Then I_X is an inverse semigroup with respect to the product defined by the composition of maps. For an inverse semigroup S , an inverse semigroup action $\alpha: S \curvearrowright X$ is a semigroup homomorphism $S \ni s \mapsto \alpha_s \in I_X$. In this paper, we always assume that every action α satisfies $\bigcup_{e \in E(S)} \text{dom}(\alpha_e) = X$.

2.4. Inverse semigroups associated to inclusions of C^* -algebras

Following [7, Proposition 13.3], we associate inverse semigroups of slices to inclusions of C^* -algebras.

Definition 2.4.1. Let $D \subset A$ be an inclusion of C^* -algebras. A slice is a norm closed subspace $M \subset A$ such that $DM \cup MD \subset M$ and $M \subset N(A, D)$. The set of all slices is denoted by $\mathcal{S}(A, D)$.

Proposition 2.4.2 ([7, Proposition 13.3]). *Let $D \subset A$ be an inclusion of C^* -algebras. Assume that D has an approximate unit for A . For $M, N \in \mathcal{S}(A, D)$, define MN to be the closure of the linear span of*

$$\{xy \in A \mid x \in M, y \in N\}.$$

Then $\mathcal{S}(A, D)$ is an inverse semigroup under this operation. The generalized inverse of $M \in \mathcal{S}(A, D)$ is $M^ := \{x^* \in A \mid x \in M\}$.*

Let G be an étale groupoid and $\text{Bis}(G)$ denotes the set of all open bisections in G . For $U, V \in \text{Bis}(G)$, their product is defined by

$$UV := \{\alpha\beta \in G \mid \alpha \in U, \beta \in V, d(\alpha) = r(\beta)\}.$$

Then $UV \in \text{Bis}(G)$ and $\text{Bis}(G)$ is an inverse semigroup with respect to this product. Note that $U^* \in \text{Bis}(G)$ is given by

$$U^{-1} := \{\alpha^{-1} \in G \mid \alpha \in U\}.$$

For $U \in \text{Bis}(G)$ and $f \in C_c(U)$, we have $f^*f \in C_0(G^{(0)})$ and

$$\|f\|_r^2 = \|f^*f\|_r = \sup_{x \in G^{(0)}} |f^*f(x)| = \sup_{\alpha \in U} |f(\alpha)|^2.$$

Hence the reduced norm coincides with the supremum norm on $C_c(U)$ and we may identify the closure of $C_c(U)$ in $C_r^*(G)$ with $C_0(U)$. Note that $C_0(U) \subset C_r^*(G)$ is a $C_0(G^{(0)})$ -subbimodule.

Now we associate a $C_0(G^{(0)})$ -subbimodule $C_0(U) \subset C_r^*(G)$ to a bisection $U \in \text{Bis}(G)$. This gives a semigroup homomorphism as in the following.

Theorem 2.4.3 ([11, Proposition 1.4.3., Corollary 2.1.6.]). *Let G be an étale groupoid. Then the map*

$$\Psi: \text{Bis}(G) \ni U \mapsto C_0(U) \in \mathcal{S}(C_r^*(G), C_0(G^{(0)}))$$

is an injective semigroup homomorphism. If G is effective, then Ψ is an isomorphism.

2.5. Étale groupoids associated to inverse semigroup actions

Many étale groupoids arise from actions of inverse semigroups to topological spaces. We recall how to construct an étale groupoid from an inverse semigroup action.

Let X be a locally compact Hausdorff space. Recall that I_X is the inverse semigroup of homeomorphisms between open sets in X . For $e \in E(S)$, we denote the domain of α_e by D_e^α . Then α_s is a homeomorphism from $D_{s^*s}^\alpha$ to $D_{ss^*}^\alpha$. We often omit α of D_e^α if there is no chance to confuse.

For an action $\alpha: S \curvearrowright X$, we associate an étale groupoid $S \ltimes_\alpha X$ as the following. First we put the set $S * X := \{(s, x) \in S \times X \mid x \in D_{s^*s}^\alpha\}$. Then we define an equivalence relation \sim on $S * X$ by declaring that $(s, x) \sim (t, y)$ holds if

$$x = y \text{ and there exists } e \in E(S) \text{ such that } x \in D_e^\alpha \text{ and } se = te.$$

Set $S \ltimes_\alpha X := S * X / \sim$ and denote the equivalence class of $(s, x) \in S * X$ by $[s, x]$. The unit space of $S \ltimes_\alpha X$ is X , where X is identified with the subset of $S \ltimes_\alpha X$ via the injective map

$$X \ni x \mapsto [e, x] \in S \ltimes_\alpha X, \quad x \in D_e^\alpha.$$

The domain map and range maps are defined by

$$d([s, x]) = x, \quad r([s, x]) = \alpha_s(x)$$

for $[s, x] \in S \ltimes_\alpha X$. The product of $[s, \alpha_t(x)], [t, x] \in S \ltimes_\alpha X$ is $[st, x]$. The inverse is $[s, x]^{-1} = [s^*, \alpha_s(x)]$. Then $S \ltimes_\alpha X$ is a groupoid in these operations. For $s \in S$ and an open set $U \subset D_{s^*s}^\alpha$, define

$$[s, U] := \{[s, x] \in S \ltimes_\alpha X \mid x \in U\}.$$

These sets form an open basis of $S \ltimes_\alpha X$. In these structures, $S \ltimes_\alpha X$ is a locally compact étale groupoid, although $S \ltimes_\alpha X$ is not necessarily Hausdorff. In this paper, we only treat inverse semigroup actions $\alpha: S \curvearrowright X$ such that $S \ltimes_\alpha X$ become Hausdorff.

Example 2.5.1. Let G be an étale groupoid. For $U \in \text{Bis}(G)$, put $\theta_U = r|_U \circ d|_U^{-1}$. Then $\theta_U: d(U) \rightarrow r(U)$ is a homeomorphism and we obtain an action $\theta: \text{Bis}(G) \curvearrowright G^{(0)}$. We call this action the canonical action of $\text{Bis}(G)$. Assume that an inverse subsemigroup $S \subset \text{Bis}(G)$ satisfies

- (1) $G = \bigcup_{U \in S} U$, and
- (2) for every $U, V \in S$ and $\alpha \in U \cap V$, there exists $W \in S$ such that $\alpha \in W \subset U \cap V$.

By [6, Proposition 5.4], G is isomorphic to $S \ltimes_\theta G^{(0)}$. Indeed, the map

$$\Phi: S \ltimes_\alpha G^{(0)} \ni [U, x] \mapsto \alpha \in G$$

is an isomorphism, where α is the unique element in U such that $d(\alpha) = x$.

We will use the following proposition to construct a groupoid homomorphism. The proof is left to the readers.

Proposition 2.5.2. *Let $\alpha: S \curvearrowright X$ and $\beta: T \curvearrowright Y$ be actions of inverse semigroups S and T on topological spaces X and Y . Assume that a continuous map $\sigma: X \rightarrow Y$ and a semigroup homomorphism $\psi: S \rightarrow T$ satisfies the following condition:*

$$\text{If } x \in X \text{ and } s \in S \text{ satisfies } x \in D_{s^*s}^\alpha, \text{ then } \sigma(x) \in D_{\psi(s)^*\psi(s)}^\beta \text{ and } \beta_{\psi(s)}(\sigma(x)) = \sigma(\alpha_s(x)) \text{ hold.}$$

Then the map

$$\Phi: S \ltimes_\alpha X \ni [s, x] \rightarrow [\psi(s), \sigma(x)] \in T \ltimes_\beta Y$$

is a continuous groupoid homomorphism. If $\sigma: X \rightarrow Y$ is locally homeomorphic, then Φ is also locally homeomorphic.

3. Main theorems

3.1. *-homomorphisms between groupoid C^* -algebras

In this subsection, we investigate *-homomorphisms between groupoid C^* -algebras. Our goal in this subsection is to show the next theorem.

Theorem 3.1.1. *Let G be an étale groupoid and H be an étale effective groupoid. Assume that we are given a *-homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ such that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ holds and $\varphi(C_0(G^{(0)}))$ is an ideal of $C_0(H^{(0)})$. Then there exist*

- (1) a closed invariant subset $F \subset G^{(0)}$,
- (2) a locally homeomorphic groupoid homomorphism $\Phi: G_F \rightarrow H$ such that $\Phi|_{U \cap G_F}$ is a homeomorphism onto its image for each $U \in \text{Bis}(G)$,² and
- (3) a continuous cocycle $c: G_F \rightarrow \mathbb{T}$,

which satisfy the following property.

²In particular, $\Phi|_F$ is a homeomorphism onto the open subset of $H^{(0)}$.

For each $U \in \text{Bis}(G)$, the following diagram is commutative:

$$\begin{array}{ccc} C_0(U) & \xrightarrow{\varphi_U} & C_0(\Phi(U \cap G_F)) \\ q_U \downarrow & \nearrow \varphi_{\Phi, c, U} & \\ C_0(U \cap G_F) & & \end{array},$$

where $\varphi_U: C_0(U) \rightarrow C_0(\Phi(U \cap G_F))$ is the restriction $\varphi|_{C_0(U)}$, $q_U: C_0(U) \rightarrow C_0(U \cap G_F)$ is a surjective bounded linear map defined by $q_U(f) = f|_{U \cap G_F}$ for $f \in C_0(U)$ and $\varphi_{\Phi, c, U}: C_0(U \cap G_F) \rightarrow C_0(\Phi(U \cap G_F))$ is a linear isometric isomorphism defined by

$$\varphi_{\Phi, c, U}(f)(\delta) := c(\Phi^{-1}(\delta)) f(\Phi^{-1}(\delta))$$

for $f \in C_0(U \cap G_F)$ and $\delta \in \Phi(U \cap G_F)$.

Moreover, assume that there exists a *-homomorphism $\tilde{\varphi}: C_r^*(G_F) \rightarrow C_r^*(H)$ with the following commutative diagram:³

$$\begin{array}{ccc} C_r^*(G) & \xrightarrow{\varphi} & C_r^*(H), \\ q \downarrow & \nearrow \tilde{\varphi} & \\ C_r^*(G_F) & & \end{array}$$

where $q: C_r^*(G) \rightarrow C_r^*(G_F)$ denote the *-homomorphism induced by the restriction

$$C_c(G) \ni f \mapsto f|_{G_F} \in C_c(G_F).$$

Then the formula

$$\tilde{\varphi}(f)(\delta) = \sum_{\alpha \in \Phi^{-1}(\{\delta\})} c(\alpha) f(\alpha)$$

holds for all $f \in C_c(G_F)$ and $\delta \in H$.

In short, the first half of Theorem 3.1.1 states that the local property of a given *-homomorphism φ can be described in terms of underlying étale groupoids. The latter half states that φ itself can be described in terms of étale groupoids if there exists $\tilde{\varphi}$. In Example 3.1.14 and the text above it, we will investigate the condition that $\tilde{\varphi}$ exists.

First, we summarize standing assumptions in this subsection. In the entirety of this subsection, we assume that G and H are étale groupoids. Moreover, we assume that

³Note that this $\tilde{\varphi}$ is unique if it exists since q is surjective. In Example 3.1.14, we give an example such that $\tilde{\varphi}$ does not exist.

H is effective except for Lemma 3.1.2 and Lemma 3.1.3. In addition, we assume that $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ is a $*$ -homomorphism such that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$. Except for Lemma 3.1.2, we assume that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ is an ideal.

Since $\ker \varphi \cap C_0(G^{(0)})$ is an ideal of $C_0(G^{(0)})$, there exists a closed subset $F \subset G^{(0)}$ such that $C_0(G^{(0)} \setminus F) = \ker \varphi \cap C_0(G^{(0)})$ holds. By [18, Lemma 10.3.1], F is an invariant set of G and therefore $G_F := d^{-1}(F)$ is a closed subgroupoid of G .

For $U \in \text{Bis}(G)$, recall that $C_0(U)$ is a $C_0(G^{(0)})$ -subbimodule of $C_r^*(G)$ in the natural way. We put $\varphi_U := \varphi|_{C_0(U)}$.

Lemma 3.1.2. *Let G and H be étale groupoids and $U \in \text{Bis}(G)$. Assume that a $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ satisfies $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$. Then there exists an isometric linear map $\widetilde{\varphi_U}: C_0(U \cap G_F) \rightarrow C_0(H^{(0)})$ that makes the following diagram commutative:*

$$\begin{array}{ccc} C_0(U) & \xrightarrow{\varphi_U} & \varphi(C_0(U)), \\ q_U \downarrow & \nearrow \widetilde{\varphi_U} & \\ C_0(U \cap G_F) & & \end{array}$$

where $q_U: C_0(U) \rightarrow C_0(U \cap G_F)$ denotes the surjective bounded linear map defined by the restriction. In particular, $\varphi(C_0(U)) \subset C_r^*(H)$ is a closed linear subspace.

Proof. Recall that $F \subset G^{(0)}$ is a closed invariant subset of G such that $C_0(G^{(0)} \setminus F) = \ker \varphi \cap C_0(G^{(0)})$. We claim that $\|\varphi(m)\| = \sup_{\alpha \in U \cap G_F} |m(\alpha)|$ holds for all $m \in C_0(U)$. Since U is a bisection, m is a normalizer for $C_0(G^{(0)})$ by Proposition 2.2.4. Hence $\varphi(m^*m) \in C_0(H^{(0)})$ follows from $m^*m \in C_0(G^{(0)})$ and $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$. By the definition of $F \subset G^{(0)}$, there exists an injective $*$ -homomorphism $\widetilde{\varphi_{G^{(0)}}}: C_0(F) \rightarrow C_0(H^{(0)})$ that makes the following diagram commutative:

$$\begin{array}{ccc} C_0(G^{(0)}) & \xrightarrow{\varphi_{G^{(0)}}} & C_0(H^{(0)}), \\ q_{G^{(0)}} \downarrow & \nearrow \widetilde{\varphi_{G^{(0)}}} & \\ C_0(F) & & \end{array}$$

where $\varphi_{G^{(0)}}$ is the restriction of φ to $C_0(G^{(0)})$ and $q_{G^{(0)}}: C_0(G^{(0)}) \rightarrow C_0(F)$ denotes the $*$ -homomorphism defined by the restriction. Hence, we obtain

$$\|\varphi(m)\|^2 = \|\varphi(m^*m)\| = \|q_{G^{(0)}}(m^*m)\| = \sup_{x \in F} |m^*m(x)| = \sup_{\alpha \in U \cap G_F} |m(\alpha)|^2$$

and therefore $\|\varphi(m)\| = \sup_{\alpha \in U \cap G_F} |m(\alpha)|$.

Now, we obtain an isometry $\widetilde{\varphi_U}: C_0(U \cap G_F) \rightarrow \varphi(C_0(U))$ that makes the following diagram commutative:

$$\begin{array}{ccc} C_0(U) & \xrightarrow{\varphi_U} & \varphi(C_0(U)) \\ \downarrow q_U & \nearrow \widetilde{\varphi_U} & \\ C_0(U \cap G_F) & & \end{array}$$

where $q_U: C_0(U) \rightarrow C_0(U \cap G_F)$ denotes the surjective bounded linear map defined by the restriction. Since $C_0(U \cap G_F)$ is complete and $\widetilde{\varphi_U}$ is an isometry, $\varphi(C_0(U)) = \widetilde{\varphi_U}(C_0(U \cap G_F))$ is a closed linear subspace of $C_r^*(H)$. ■

In the rest of this subsection, we assume that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ is an ideal of $C_0(H^{(0)})$.

Lemma 3.1.3. *Assume that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ is an ideal of $C_0(H^{(0)})$. Then*

$$\varphi(C_0(U)) \subset C_r^*(H)$$

is a $C_0(H^{(0)})$ -subbimodule and

$$\varphi(C_0(U)) \subset N(C_r^*(H), C_0(H^{(0)}))$$

holds for all $U \in \text{Bis}(G)$.

Proof. Take $m \in C_0(U)$ and $a, b \in C_0(H^{(0)})$. Let $\{e_i\}_{i \in I} \subset C_0(G^{(0)})$ be an approximate identity for $C_r^*(G)$. Since we assume that $\varphi(C_0(G^{(0)}))$ is an ideal of $C_0(H^{(0)})$, $a\varphi(e_i)$ and $\varphi(e_i)b$ are contained in $\varphi(C_0(G^{(0)}))$ for all $i \in I$. There exists $f, g \in C_0(G^{(0)})$ such that $\varphi(f) = a\varphi(e_i)$ and $\varphi(g) = \varphi(e_i)b$ hold. Now we have

$$a\varphi(e_i)\varphi(m)\varphi(e_i)b = \varphi(fmg) \in \varphi(C_0(U))$$

since $C_0(U) \subset C_r^*(G)$ is a $C_0(G^{(0)})$ -subbimodule. Since $\{e_i m e_i\}_{i \in I}$ converges to m , we obtain $a\varphi(m)b \in \varphi(C_0(U))$ by Lemma 3.1.2. Hence, $\varphi(C_0(U))$ is a $C_0(H^{(0)})$ -subbimodule.

Next, we show $\varphi(C_0(U)) \subset N(C_r^*(H), C_0(H^{(0)}))$. Take $m \in C_0(U)$ and $h \in C_0(H^{(0)})$. Let $\{e_i\}_{i \in I} \subset C_0(G^{(0)})$ be an approximate identity for $C_r^*(G)$. Since $\varphi(e_i)h \in \varphi(C_0(G^{(0)}))$ and $m \in N(C_r^*(G), C_0(G^{(0)}))$, we have $\varphi(m)\varphi(e_i)h\varphi(m^*) \in \varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$. Hence we obtain $\varphi(m)h\varphi(m)^* \in C_0(H^{(0)})$. One can show $\varphi(m)^*h\varphi(m) \in C_0(H^{(0)})$ in the same way. Therefore, $\varphi(m)$ is a normalizer for $C_0(H^{(0)})$ and we obtain

$$\varphi(C_0(U)) \subset N(C_r^*(H), C_0(H^{(0)})).$$

■

In the rest of this subsection, we assume that an étale groupoid H is effective.

Lemma 3.1.4. *Assume that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ is an ideal and H is effective. Then there exists a semigroup homomorphism $\psi: \text{Bis}(G) \rightarrow \text{Bis}(H)$ such that $\varphi(C_0(U)) = C_0(\psi(U))$ holds for all $U \in \text{Bis}(G)$.*

Proof. Take $U \in \text{Bis}(G)$. Then $\varphi(C_0(U)) \subset C_r^*(H)$ is a closed $C_0(H^{(0)})$ -subbimodule that consists of normalizers for $C_0(H^{(0)})$ by Lemma 3.1.2 and Lemma 3.1.3. Hence, there exists a unique $\psi(U) \in \text{Bis}(H)$ such that $\varphi(C_0(U)) = C_0(\psi(U))$ holds by Theorem 2.4.3.

We show the map $\psi: \text{Bis}(G) \rightarrow \text{Bis}(H)$ is a semigroup homomorphism. Take $U_1, U_2 \in \text{Bis}(G)$. Then we have

$$\begin{aligned} C_0(\psi(U_1)\psi(U_2)) &= C_0(\psi(U_1))C_0(\psi(U_2)) \\ &= \varphi(C_0(U_1))\varphi(C_0(U_2)) \\ &= \varphi(C_0(U_1U_2)) \\ &= C_0(\psi(U_1U_2)). \end{aligned}$$

Thus, we obtain $\psi(U_1)\psi(U_2) = \psi(U_1U_2)$ and ψ is a semigroup homomorphism. ■

We let

$$\psi: \text{Bis}(G) \rightarrow \text{Bis}(H)$$

denote the semigroup homomorphism defined in Lemma 3.1.4.

Lemma 3.1.5. *Put*

$$T := \{U \cap G_F \in \text{Bis}(G_F) \mid U \in \text{Bis}(G)\}.$$

Then T is an inverse subsemigroup of $\text{Bis}(G_F)$ and there exists a subsemigroup homomorphism $\tilde{\psi}: T \rightarrow \text{Bis}(H)$ that makes the following diagram commutative:

$$\begin{array}{ccc} \text{Bis}(G) & \xrightarrow{\psi} & \text{Bis}(H), \\ \downarrow Q & \nearrow \tilde{\psi} & \\ T & & \end{array}$$

where $Q: \text{Bis}(G) \rightarrow T$ is the semigroup homomorphism defined by $Q(U) = U \cap G_F$ for each $U \in \text{Bis}(G)$.

Proof. One can show that T is an inverse subsemigroup of $\text{Bis}(G_F)$ by straightforward calculations. It is sufficient to show that $\psi(U_1) = \psi(U_2)$ holds for $U_1, U_2 \in \text{Bis}(G)$ which satisfy $U_1 \cap G_F = U_2 \cap G_F$. It follows that $U_1 \cap G_F = U_1 \cap U_2 \cap G_F$ from $U_1 \cap G_F = U_2 \cap G_F$. By Lemma 3.1.2, there exists an isometry $\widetilde{\varphi_{U_1}}: C_0(U_1 \cap G_F) \rightarrow \varphi(C_0(U_1))$ that makes the following diagram commutative:

$$\begin{array}{ccc} C_0(U_1) & \xrightarrow{\varphi_{U_1}} & \varphi(C_0(U_1)), \\ \downarrow q_{U_1} & \nearrow \widetilde{\varphi_{U_1}} & \\ C_0(U_1 \cap G_F) & & \end{array}$$

where $q_{U_1}: C_0(U_1) \rightarrow C_0(U_1 \cap G_F)$ denotes the surjective bounded linear map defined by the restriction. Now we have

$$\begin{aligned}\varphi(C_0(U_1)) &= \widehat{\varphi_{U_1}}(C_0(U_1 \cap G_F)) = \widehat{\varphi_{U_1}}(C_0(U_1 \cap U_2 \cap G_F)) \\ &= \widehat{\varphi_{U_1}}(q_{U_1}(C_0(U_1 \cap U_2))) = \varphi(C_0(U_1 \cap U_2)).\end{aligned}$$

Thus we obtain $\psi(U_1) = \psi(U_1 \cap U_2)$. Replacing U_1 with U_2 , we obtain

$$\psi(U_2) = \psi(U_1 \cap U_2)$$

and therefore $\psi(U_1) = \psi(U_2)$. ■

Since we assume that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ is an ideal, there exists an open set $V \subset H^{(0)}$ such that $\varphi(C_0(G^{(0)})) = C_0(V)$. In addition, since F satisfies $C_0(G^{(0)} \setminus F) = \ker \varphi \cap C_0(G^{(0)})$, there exists a *-homomorphism $\widehat{\varphi_{G^{(0)}}}: C_0(F) \rightarrow C_0(V)$ that makes the following diagram commutative:

$$\begin{array}{ccc} C_0(G^{(0)}) & \xrightarrow{\varphi_{G^{(0)}}} & C_0(V), \\ q_{G^{(0)}} \downarrow & \nearrow \widehat{\varphi_{G^{(0)}}} & \\ C_0(F) & & \end{array}$$

where $q_{G^{(0)}}$ denotes the *-homomorphism defined by the restriction. One can see that $\widehat{\varphi_{G^{(0)}}}$ is indeed an *-isomorphism. By Gelfand–Naimark duality, there exists a homeomorphism $\widehat{\varphi_{G^{(0)}}}: V \rightarrow F$ such that $\widehat{\varphi_{G^{(0)}}}(f)(y) = f(\widehat{\varphi_{G^{(0)}}}(y))$ holds for all $f \in C_0(F)$ and $y \in V$. Put $\sigma := \widehat{\varphi_{G^{(0)}}}^{-1}: F \rightarrow V \subset H^{(0)}$. We regard the range of σ as $H^{(0)}$ rather than V .

Now, we obtain a semigroup homomorphism $\tilde{\psi}: T \rightarrow \text{Bis}(H)$ and a homeomorphism $\sigma: F \rightarrow H^{(0)}$.

Lemma 3.1.6. *The above maps $\sigma: F \rightarrow H^{(0)}$ and $\tilde{\psi}: T \rightarrow \text{Bis}(H)$ satisfy the condition in Proposition 2.5.2 for the canonical actions of bisections $T \curvearrowright F$ and $\text{Bis}(H) \curvearrowright H^{(0)}$.*

Proof. Take $x \in F$ and $U \in \text{Bis}(G)$ such that $x \in d(U \cap G_F)$. First, we show that $\sigma(x) \in d(\tilde{\psi}(U \cap G_F))$. We have

$$\begin{aligned}C_0(\sigma(d(U \cap G_F))) &= C_0(\widehat{\varphi_{G^{(0)}}}^{-1}(d(U \cap G_F))) \\ &= \widehat{\varphi_{G^{(0)}}}(C_0(d(U \cap G_F))) = C_0(\tilde{\psi}(d(U \cap G_F))).\end{aligned}$$

Note that we use the condition that $\widehat{\varphi_{G^{(0)}}}$ is injective to deduce the second equality. Thus, we obtain $\sigma(d(U \cap G_F)) = \tilde{\psi}(d(U \cap G_F))$. Since $\tilde{\psi}$ is a semigroup homomorphism, we have $\tilde{\psi}(d(U \cap G_F)) = d(\tilde{\psi}(U \cap G_F))$. Therefore we obtain

$$\sigma(x) \in \sigma(d(U \cap G_F)) = d(\tilde{\psi}(U \cap G_F)).$$

Since we assume that $x \in d(U \cap G_F)$, there exists $\alpha \in U \cap G_F$ such that $d(\alpha) = x$. In addition, since $\sigma(x) \in d(\tilde{\psi}(U \cap G_F))$, there exists $\delta \in \tilde{\psi}(U \cap G_F)$ such that $d(\delta) = \sigma(x)$. In order to complete the proof, it is sufficient to show $\sigma(r(\alpha)) = r(\delta)$. Instead, we show that $r(\alpha) = \widehat{\varphi_{G^{(0)}}}(r(\delta))$. Take $n \in C_0(U)$ such that $n(\alpha) \neq 0$. For all $f \in C_0(G^{(0)})$, we have

$$\begin{aligned} |n(\alpha)|^2 f(r(\alpha)) &= n^* f n(d(\alpha)) = q_{G^{(0)}}(n^* f n)(x) = q_{G^{(0)}}(n^* f n)(\widehat{\varphi_{G^{(0)}}}(d(\delta))) \\ &= \widehat{\varphi_{G^{(0)}}}(q_{G^{(0)}}(n^* f n))(d(\delta)) = \varphi_{G^{(0)}}(n^* f n)(d(\delta)) \\ &= \varphi(n)^* \varphi(f) \varphi(n)(d(\delta)) = \varphi(n^* n)(d(\delta)) \varphi(f)(r(\delta)) \\ &= \varphi(n^* n)(\sigma(x)) \widehat{\varphi_{G^{(0)}}}(q_{G^{(0)}}(f))(r(\delta)) \\ &= n^* n(x) q_{G^{(0)}}(f)(\widehat{\varphi_{G^{(0)}}}(r(\delta))) = |n(\alpha)|^2 f(\widehat{\varphi_{G^{(0)}}}(r(\delta))). \end{aligned}$$

Hence, $f(r(\alpha)) = f(\widehat{\varphi_{G^{(0)}}}(r(\delta)))$ holds for all $f \in C_0(G^{(0)})$. Therefore we obtain $r(\alpha) = \widehat{\varphi_{G^{(0)}}}(r(\delta))$ by Urysohn's lemma. ■

By Lemma 3.1.6 and Proposition 2.5.2, we obtain the groupoid homomorphism $\Phi: T \ltimes F \rightarrow \text{Bis}(H) \ltimes H^{(0)}$. Since $\Phi|_F: F \rightarrow H^{(0)}$ is a homeomorphism onto its image, Φ is locally homeomorphic. Since we may apply Example 2.5.1 to the canonical actions $T \curvearrowright F$ and $\text{Bis}(H) \curvearrowright H^{(0)}$, $T \ltimes F$ and $\text{Bis}(H) \ltimes H^{(0)}$ are isomorphic to G_F and H respectively. Thus we obtain the groupoid homomorphism from G_F to H and denote it by Φ again. This Φ is given explicitly as follows. Take $\alpha \in G_F$ and $U \in \text{Bis}(G)$ with $\alpha \in U$. Then there exists $\alpha' \in \psi(U)$ such that $\sigma(d(\alpha)) = d(\alpha')$. This α' is nothing but $\Phi(\alpha)$. In the proof of Lemma 3.1.6, we obtained $\sigma(d(U \cap G_F)) = \tilde{\psi}(d(U \cap G_F))$. In addition, we have $\tilde{\psi}(d(U \cap G_F)) = d(\tilde{\psi}(U \cap G_F)) = d(\psi(U))$. Therefore, we have the following commutative diagram:

$$\begin{array}{ccc} U \cap G_F & \xrightarrow{\Phi} & \psi(U) \\ d \downarrow & & \downarrow d \\ d(U \cap G_F) & \xrightarrow{\sigma} & d(\psi(U)). \end{array}$$

In particular, Φ gives a homeomorphism from $U \cap G_F$ to $\psi(U)$ since the vertical domain maps and σ are homeomorphisms. Note that we have $\Phi|_F = \sigma = \widehat{\varphi_{G^{(0)}}}^{-1}$.

Lemma 3.1.7. *Fix $\alpha \in G_F$. Take $U \in \text{Bis}(G)$ and $n \in C_0(U)$ such that $\alpha \in U$ and $n(\alpha) \neq 0$ hold. Then $|\varphi(n)(\Phi(\alpha))| = |n(\alpha)|$ holds. Moreover, the value*

$$\frac{\varphi(n)(\Phi(\alpha))}{n(\alpha)} \in \mathbb{T}$$

is independent of the choice of U and n .

Proof. First, we show $|\varphi(n)(\Phi(\alpha))| = |n(\alpha)|$. This follows from the next calculation:

$$\begin{aligned} |\varphi(n)(\Phi(\alpha))|^2 &= \varphi(n^*n)(d(\Phi(\alpha))) = \varphi(n^*n)(\Phi(d(\alpha))) \\ &= \widehat{\varphi_{G^{(0)}}}(q_{G^{(0)}}(n^*n)(\Phi(d(\alpha)))) \\ &= q_{G^{(0)}}(n^*n)(d(\alpha)) = |n(\alpha)|^2. \end{aligned}$$

Next, we show that the value

$$\frac{\varphi(n)(\Phi(\alpha))}{n(\alpha)} \in \mathbb{T}$$

is independent of the choice of U and n . Consider $W \in \text{Bis}(G)$ and $m \in C_0(W)$ such that $\alpha \in W$ and $m(\alpha) \neq 0$ hold. Then we have

$$n^*m(d(\alpha)) = \overline{n(\alpha)}m(\alpha)$$

and

$$\varphi(n^*m)(\Phi(d(\alpha))) = \varphi(n^*m)(d(\Phi(\alpha))) = \overline{\varphi(n)(\Phi(\alpha))}\varphi(m)(\Phi(\alpha)).$$

Take $f \in C_c(d(U \cap W))$ such that $f(d(\alpha)) = 1$. Then we have $nf, mf \in C_c(U \cap W)$ and $(nf)^*(mf) \in C_0(G^{(0)})$. Combining with

$$\begin{aligned} n^*m(d(\alpha)) &= (nf)^*(mf)(d(\alpha)) = \varphi((nf)^*(mf))(\Phi(d(\alpha))) \\ &= \overline{\varphi(n)(\Phi(\alpha))}\varphi(m)(\Phi(\alpha)) \end{aligned}$$

and $\varphi(n)(\Phi(\alpha))/n(\alpha) \in \mathbb{T}$, we obtain

$$\frac{\varphi(m)(\Phi(\alpha))}{m(\alpha)} = \frac{\overline{n(\alpha)}}{\varphi(n)(\Phi(\alpha))} = \frac{\varphi(n)(\Phi(\alpha))}{n(\alpha)}.$$

Thus, the value

$$\frac{\varphi(n)(\Phi(\alpha))}{n(\alpha)} \in \mathbb{T}$$

is independent of the choice of U and n . ■

Lemma 3.1.8. Define $c: G_F \rightarrow \mathbb{T}$ by

$$c(\alpha) := \frac{\varphi(n)(\Phi(\alpha))}{n(\alpha)},$$

where $n \in C_0(U)$ and $U \in \text{Bis}(G)$ satisfies $\alpha \in U$ and $n(\alpha) \neq 0$.⁴ Then $c: G_F \rightarrow \mathbb{T}$ is a continuous groupoid homomorphism.

⁴Note that c is well defined by Lemma 3.1.7.

Proof. First, we show that c is continuous. Fix $\alpha \in G_F$. Take $U \in \text{Bis}(G)$ and $n \in C_0(U)$ so that $\alpha \in U$ and $n(\alpha) \neq 0$ holds. Then we have

$$c(\gamma) = \frac{\varphi(n)(\Phi(\gamma))}{n(\gamma)}$$

for all $\gamma \in \text{supp}^\circ(n)$. Since $\varphi(n) \circ \Phi$ and n are continuous, c is also continuous at α . Since $\alpha \in G_F$ is arbitrary, c is continuous on G_F .

Next, we show that c is a groupoid homomorphism. Fix $\alpha, \beta \in G_F$ with $d(\alpha) = r(\beta)$. Take $U_1, U_2 \in \text{Bis}(G)$, $n \in C_0(U_1)$ and $m \in C_0(U_2)$ so that $\alpha \in U_1$, $\beta \in U_2$, $n(\alpha) \neq 0$ and $m(\beta) \neq 0$ hold. Then we have $nm \in C_0(UV)$ and $nm(\alpha\beta) = n(\alpha)m(\beta) \neq 0$. Hence, we obtain

$$c(\alpha\beta) = \frac{\varphi(nm)(\Phi(\alpha\beta))}{nm(\alpha\beta)} = \frac{\varphi(n)(\Phi(\alpha))\varphi(m)(\Phi(\beta))}{n(\alpha)m(\beta)} = c(\alpha)c(\beta).$$

Here, the second equality follows from the fact that $\varphi(n)$ and $\varphi(m)$ are supported on bisections that contain $\Phi(\alpha)$ and $\Phi(\beta)$ respectively. Therefore, $c: G_F \rightarrow \mathbb{T}$ is a continuous groupoid homomorphism. \blacksquare

Thus, we have obtained the locally homeomorphic groupoid homomorphism $\Phi: G_F \rightarrow H$ and the continuous cocycle $c: G_F \rightarrow \mathbb{T}$. From the remark beneath Lemma 3.1.6, $\Phi|_{U \cap G_F}$ is a homeomorphism onto its image for each $U \in \text{Bis}(G)$. Now, we are ready to show the first half of Theorem 3.1.1.

Proposition 3.1.9. *Fix $U \in \text{Bis}(G)$. Define a homeomorphism*

$$\Phi_U := \Phi|_{U \cap G_F}: U \cap G_F \rightarrow \Phi(U \cap G_F).$$

In addition, define $\varphi_{\Phi, c, U}: C_0(U \cap G_F) \rightarrow C_0(\Phi(U \cap G_F))$ to be

$$\varphi_{\Phi, c, U}(f)(\delta) := c(\Phi_U^{-1}(\delta))f(\Phi_U^{-1}(\delta))$$

for $f \in C_0(U \cap G_F)$ and $\delta \in \Phi(U \cap G_F)$. Then $\varphi_{\Phi, c, U}$ is a linear isometric isomorphism. Moreover, $\varphi_U = \varphi_{\Phi, c, U} \circ q_U$ and $\widetilde{\varphi_U} = \varphi_{\Phi, c, U}$ hold.

Proof. It follows that $\varphi_{\Phi, c, U}$ is a linear isometric isomorphism from the direct calculation. Note that the inverse map of $\varphi_{\Phi, c, U}$ is given by

$$\varphi_{\Phi, c, U}^{-1}(g)(\alpha) = \overline{c(\alpha)}g(\Phi_U(\alpha))$$

for $g \in C_0(\Phi(U \cap G_F))$ and $\alpha \in U \cap G_F$.

We show $\varphi_U = \varphi_{\Phi, c, U} \circ q_U$. Take $g \in C_0(U)$ and $\delta \in \Phi(U \cap G_F)$. Assume that $g(\Phi_U^{-1}(\delta)) \neq 0$. Then we have

$$c(\Phi_U^{-1}(\delta)) = \frac{\varphi(g)(\Phi(\Phi_U^{-1}(\delta)))}{g(\Phi_U^{-1}(\delta))} = \frac{\varphi(g)(\delta)}{g(\Phi_U^{-1}(\delta))} = \frac{\varphi_U(g)(\delta)}{q_U(g)(\Phi_U^{-1}(\delta))}.$$

Thus, we obtain

$$\varphi_U(g)(\delta) = c(\Phi_U^{-1}(\delta))q_U(g)(\Phi_U^{-1}(\delta)) = \varphi_{\Phi,c,U} \circ q_U(g)(\delta).$$

If $g(\Phi_U^{-1}(\delta)) = 0$, we have

$$\begin{aligned} |\varphi_U(g)(\delta)|^2 &= \varphi(g^*g)(d(\delta)) = g^*g(\Phi_U^{-1}(d(\delta))) \\ &= g^*g(d(\Phi_U^{-1}(\delta))) = |g(\Phi_U^{-1}(\delta))|^2 = 0 \end{aligned}$$

and therefore

$$\varphi_{\Phi,c,U} \circ q_U(g)(\delta) = c(\Phi_U^{-1}(\delta))g(\Phi_U^{-1}(\delta)) = 0.$$

Hence, we obtain

$$\varphi_U(g)(\delta) = \varphi_{\Phi,c,U} \circ q_U(g)(\delta).$$

for all $g \in C_0(U)$ and $\delta \in \Phi(U \cap G_F)$. The last assertion follows from the fact that we have $\varphi_U = \widetilde{\varphi}_U \circ q_U$ and q_U is surjective. ■

Define a *-homomorphism $\varphi_{\Phi,c}: C_c(G_F) \rightarrow C_c(H)$ to be

$$\varphi_{\Phi,c}(f)(\delta) := \sum_{\alpha \in \Phi^{-1}(\delta)} c(\alpha)f(\alpha)$$

for $f \in C_c(G_F)$ and $\delta \in H$. Note that the right hand side of the above formula is a finite sum since Φ is locally homeomorphic and the support of f is compact. In addition, we use the condition that Φ is injective on F to check that $\varphi_{\Phi,c}$ preserves the multiplications. Thus one can check that $\varphi_{\Phi,c}: C_c(G_F) \rightarrow C_c(H)$ is actually a *-homomorphism. We shall remark that $\varphi_{\Phi,c}$ does not always extend to $C_r^*(G_F)$ if G_F is not amenable. We will give a relevant example in Example 3.1.14.

Finally, we complete the proof of Theorem 3.1.1.

Proposition 3.1.10. *Assume that there exists a *-homomorphism*

$$\widetilde{\varphi}: C_r^*(G_F) \rightarrow C_r^*(H)$$

with the following commutative diagram:

$$\begin{array}{ccc} C_r^*(G) & \xrightarrow{\varphi} & C_r^*(H), \\ \downarrow q & \nearrow \widetilde{\varphi} & \\ C_r^*(G_F) & & \end{array}$$

where $q: C_r^*(G) \rightarrow C_r^*(G_F)$ denote the *-homomorphism induced by the restriction. Then $\widetilde{\varphi}(f) = \varphi_{\Phi,c}(f)$ holds for all $f \in C_c(G_F)$. In particular, $\varphi_{\Phi,c}$ extends to $C_r^*(G_F)$ and $\widetilde{\varphi} = \varphi_{\Phi,c}$ holds.

Proof. Fix $U \in \text{Bis}(G)$. By Proposition 3.1.9, we have $\tilde{\varphi}(f) = \varphi_{\Phi, c}(f)$ for all $f \in C_c(U \cap G_F)$. Since $C_c(G_F)$ is the linear span of $\bigcup_{U \in \text{Bis}(G)} C_c(U \cap G_F)$, we obtain $\tilde{\varphi}(f) = \varphi_{\Phi, c}(f)$ for all $f \in C_c(G_F)$. ■

Now, we have completed the proof of Theorem 3.1.1. In the last of this subsection, we give some remarks about our results in this subsection. First, we present a special case of Theorem 3.1.1. Applying Theorem 3.1.1 for surjective $*$ -homomorphisms, we obtain quotients of underlying étale groupoids (Corollary 3.1.11). We refer the readers to [9, Section 3] for the quotient étale groupoids. We remark that the quotient étale groupoid $G/\text{Iso}(G)^\circ$ appearing in Corollary 3.1.11 is nothing but the groupoid of germs associated with G in the sense of [17, Proposition 3.2].

Corollary 3.1.11. *Let G be an étale groupoid and H be an étale effective groupoid. Assume that there exists a surjective $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ such that $\varphi(C_0(G^{(0)})) = C_0(H^{(0)})$. Then there exists a closed invariant subset $F \subset G^{(0)}$ such that H is isomorphic to $G_F/\text{Iso}(G_F)^\circ$ as étale groupoids. Moreover, if φ is injective on $C_0(G^{(0)})$ (and therefore $C_0(G^{(0)})$ and $C_0(H^{(0)})$ are isomorphic), then H is isomorphic to $G/\text{Iso}(G)^\circ$.*

Proof. Applying Theorem 3.1.1, we obtain a closed invariant $F \subset G^{(0)}$ and a locally homeomorphic groupoid homomorphism $\Phi: G_F \rightarrow H$. Since we assume that φ is surjective, it turns out that $\Phi: G_F \rightarrow H$ is also surjective. Indeed, assume that $\Phi: G_F \rightarrow H$ is not surjective. Then there exists $\delta \in H \setminus \Phi(G_F)$. Using Proposition 2.2.1, define $\text{ev}_\delta: C_r^*(H) \rightarrow \mathbb{C}$ by

$$\text{ev}_\delta(a) := j(a)(\delta)$$

for $a \in C_r^*(H)$. Then, for each $U \in \text{Bis}(G)$, we have

$$\varphi(C_c(U)) \subset C_0(\Phi(U \cap G_F)) \subset \ker \text{ev}_\delta$$

by the formula $\varphi_U = \varphi_{\Phi, c, U} \circ q_U$ in Theorem 3.1.1. In addition, by the linearity of φ and the partition of unity argument, we obtain $\varphi(C_c(G)) \subset \ker \text{ev}_\delta$. Hence we obtain $\varphi(C_r^*(G)) \subset \ker \text{ev}_\delta$ by the continuity of φ . This contradicts to the assumption that φ is surjective and fact that $\ker \text{ev}_\delta \subsetneq C_r^*(H)$. Therefore, $\Phi: G_F \rightarrow H$ is surjective.

Now, one can see $\Phi^{-1}(H^{(0)}) = \text{Iso}(G_F)^\circ$ since H is effective. Therefore, Φ induces an isomorphism $G_F/\text{Iso}(G_F)^\circ \simeq H$ by the fundamental theorem on homomorphisms [10, Proposition 2.2]. Now, the last assertion is obvious since F coincides with $G^{(0)}$ if φ is injective on $C_0(G^{(0)})$. ■

Combining Corollary 3.1.11 and groupoid quotients, we obtain the following rigidity result (Corollary 3.1.12) for not necessarily effective groupoids.

Corollary 3.1.12. *Let G and H be étale (not necessarily effective) groupoids and $Q: H \rightarrow H/\text{Iso}(H)^\circ$ be the quotient map. Assume that there exists a surjective $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ such that φ give an isomorphism between $C_0(G^{(0)})$ and $C_0(H^{(0)})$.*

In addition, assume that $\text{Iso}(H)^\circ \subset H$ is closed and the *-homomorphism

$$\varphi_Q: C_c(H) \ni f \mapsto \left(\delta \mapsto \sum_{\alpha \in Q^{-1}(\delta)} f(\alpha) \right) \in C_c(H/\text{Iso}(H)^\circ)$$

extends to the *-homomorphism from $C_r^*(H)$ to $C_r^*(H/\text{Iso}(H)^\circ)$.⁵ Then $G/\text{Iso}(G)^\circ$ is isomorphic to $H/\text{Iso}(H)^\circ$.

Proof. First, $H/\text{Iso}(H)^\circ$ is Hausdorff since we assume that $\text{Iso}(H)^\circ \subset H$ is closed [9, Proposition 3.11]. We denote the extension of φ_Q by $\varphi_Q: C_r^*(H) \rightarrow C_r^*(H/\text{Iso}(H)^\circ)$ again. Then φ_Q is surjective and the restriction $\varphi_Q|_{C_0(H^{(0)})}$ is an isomorphism onto $C_0((H/\text{Iso}(H)^\circ)^{(0)})$ by [9, Proposition 3.13 and Lemma 3.14]. In addition, note that $H/\text{Iso}(H)^\circ$ is effective. Therefore we may apply Corollary 3.1.11 for $\varphi_Q \circ \varphi: C_r^*(G) \rightarrow C_r^*(H/\text{Iso}(H)^\circ)$ and this yields an isomorphism $G/\text{Iso}(G)^\circ \simeq H/\text{Iso}(H)^\circ$. ■

Remark 3.1.13. We remark that the converse of Corollary 3.1.12 does not hold. For example, let $G := \{e\}$ be the trivial group and $H := \mathbb{Z}$. Then both of $G/\text{Iso}(G)^\circ$ and $H/\text{Iso}(H)^\circ$ are trivial groups although there does not exist a surjective *-homomorphism from $C_r^*(G) = \mathbb{C}$ to $C_r^*(H) = C(\mathbb{T})$.

From Corollary 3.1.12, we deduce that the quotient groupoids $G/\text{Iso}(G)^\circ$ define an invariant for inclusions of C^* -algebras $(C_r^*(G), C_0(G^{(0)}))$ associated with étale amenable groupoids G such that $\text{Iso}(G)^\circ \subset G$ are closed. Namely, for étale amenable groupoids G and H such that $\text{Iso}(G)^\circ \subset G$ and $\text{Iso}(H)^\circ \subset H$ are closed, $G/\text{Iso}(G)^\circ$ and $H/\text{Iso}(H)^\circ$ are isomorphic if there exists a *-isomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ such that $\varphi(C_0(G^{(0)})) = C_0(H^{(0)})$. In particular, since the orbit spaces of G and $G/\text{Iso}(G)^\circ$ are homeomorphic, the orbit spaces $G^{(0)}/G$ also define an invariant for inclusions of C^* -algebras $(C_r^*(G), C_0(G^{(0)}))$ associated with étale amenable groupoids G with closed $\text{Iso}(G)^\circ$. Note that $G^{(0)}/G$ is the quotient space of $G^{(0)}$ with respect to the equivalence relation defined by declaring $x \sim y$ if there exists $\alpha \in G$ such that $d(\alpha) = x$ and $r(\alpha) = y$.

Next, we investigate the condition that $\tilde{\varphi}$ in Proposition 3.1.10 exists. Recall that an étale groupoid G is said to be inner exact if the following sequence

$$0 \rightarrow C_r^*(G_{G^{(0)} \setminus F}) \xrightarrow{\iota} C_r^*(G) \xrightarrow{q} C_r^*(G_F) \rightarrow 0$$

is exact for each closed invariant subset $F \subset G^{(0)}$, where ι is the inclusion map and q is the restriction. If G is inner exact, then $\tilde{\varphi}$ in Proposition 3.1.10 exists. In particular, $\tilde{\varphi}$ always exists for all amenable groupoids, since amenable groupoids are inner exact (see [18, Definition 10.1.2, Theorem 10.1.4 and Proposition 10.3.2] for this fact and the amenability of étale groupoids). In addition, if φ is injective on $C_0(G^{(0)})$, then $\tilde{\varphi}$ exists and $\tilde{\varphi}$ is nothing but φ since $F = G^{(0)}$ holds. In particular, no matter whether G is amenable or not, every *-automorphisms on $C_r^*(G)$ that preserves $C_0(G^{(0)})$ comes from groupoid automorphisms and continuous cocycles on G (Corollary 3.2.2).

⁵This assumption holds if H is amenable. See [18, Definition 10.1.2] for the definition of amenable groupoids.

On the other hand, Example 3.1.14 is an example such that $\tilde{\varphi}$ does not exist, although the étale groupoid G in Example 3.1.14 is not effective. To our knowledge, it is an open problem whether $\tilde{\varphi}$ always exists or not under the assumption that G is effective.

Example 3.1.14 (HLS groupoid). We give an example where $\tilde{\varphi}$ does not exist.

Let G be the HLS groupoid associated with the free group $\Gamma := F_2$ (see [19] for details) and $H := \{e\}$ be the trivial group. Then we have $C_r^*(H) = \mathbb{C}$. Recall that

$$G = \coprod_{n \in \mathbb{N} \cup \{\infty\}} \Gamma_n \times \{n\},$$

where Γ_n is a finite quotient group of Γ for $n \in \mathbb{N}$ and $\Gamma_\infty = \Gamma$ (see [19] for the precise definition). Define a $*$ -homomorphism $\varphi: C_c(G) \rightarrow \mathbb{C}$ by

$$\varphi(f) := \sum_{s \in \Gamma} f((s, \infty))$$

for $f \in C_c(G)$. Then φ extends to the $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow \mathbb{C}$ since $C_r^*(G)$ coincides with the universal groupoid C^* -algebra $C^*(G)$ by [19, Theorem 1.2]. Note that the corresponding closed invariant set is $F = \{\infty\}$. Since F_2 is not amenable, $\tilde{\varphi}: C_r^*(G_F) \rightarrow \mathbb{C}$ does not exist by [4, Theorem 2.6.8].

Example 3.1.15. From a $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ such that $\varphi(C_0(G^{(0)})) \subset C_0(H^{(0)})$ is an ideal, we have constructed a closed invariant subset $F \subset G^{(0)}$, a locally homeomorphic groupoid homomorphism $\Phi: G_F \rightarrow H$ such that $\Phi|_F$ is injective and a continuous cocycle $c: G_F \rightarrow \mathbb{T}$. If G is amenable, this construction is bijective. Indeed, for such $F \subset G^{(0)}$, $\Phi: G_F \rightarrow H$ and $c: G_F \rightarrow \mathbb{T}$, the $*$ -homomorphism $\varphi_{\Phi, c}: C_c(G_F) \rightarrow C_r^*(H)$ in Proposition 3.1.10 extends to $C_r^*(G_F)$ by [18, Theorems 9.2.3 and 10.1.4]. Thus, we obtain a $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ which induces F , Φ and c . If G is not amenable, these constructions need not to be bijective. We give such an example.

Let Γ be a countably infinite discrete group and $X := \Gamma \cup \{\infty\}$ be the one point compactification of Γ . The left multiplication $\Gamma \curvearrowright \Gamma$ extends to the action on X and we denote it by $\sigma: \Gamma \curvearrowright X$. Putting $G := \Gamma \ltimes_\sigma X$, then G is an effective étale groupoid. In addition, $F := \{\infty\}$ is a closed invariant subset of G . Let $H := \{e\}$ be the trivial group, $\Phi: G_F \rightarrow H$ be the unique group homomorphism and $c = 1$. If Γ is not amenable, then there does not exist a $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ that induces F , Φ and c . Actually, there does not exist a nonzero $*$ -homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ if Γ is not amenable. If such $\varphi: C_r^*(G) \rightarrow C_r^*(H)$ exists, then one obtains a nonzero $*$ -homomorphism $C_r^*(\Gamma) \rightarrow \mathbb{C}$ by composing φ with the canonical inclusion $C_r^*(\Gamma) \rightarrow C(X) \rtimes \Gamma = C_r^*(G)$. This contradicts to the non-amenability of Γ (see [4, Theorem 2.6.8], for example).

3.2. Automorphism groups of $C_r^*(G)$ that globally preserve $C_0(G^{(0)})$

In this subsection, we investigate automorphism groups of $C_r^*(G)$ that globally preserve $C_0(G^{(0)})$.

Definition 3.2.1. Let $D \subset A$ be an inclusion of C^* -algebras. We define $\text{Aut}_D(A)$ and $\text{FAut}_D(A)$ to be

$$\begin{aligned}\text{Aut}_D(A) &:= \{\varphi \in \text{Aut}(A) \mid \varphi(D) = D\}, \\ \text{FAut}_D(A) &:= \{\varphi \in \text{Aut}(A) \mid \varphi(a) = a \text{ for all } a \in D\}.\end{aligned}$$

Note that $\text{Aut}_D(A)$ and $\text{FAut}_D(A)$ are subgroups of $\text{Aut}(A)$ and we have

$$\text{FAut}_D(A) \subset \text{Aut}_D(A).$$

For an étale groupoid G , recall that we define

$$Z(G, \mathbb{T}) := \{c: G \rightarrow \mathbb{T} \mid c \text{ is a continuous groupoid homomorphism}\}.$$

Then $Z(G, \mathbb{T})$ is an abelian group with respect to the pointwise product. In addition, $\text{Aut}(G)$ naturally acts on $Z(G, \mathbb{T})$. Indeed, for $\Phi \in \text{Aut}(G)$ and $c \in Z(G, \mathbb{T})$, $\Phi.c \in Z(G, \mathbb{T})$ is defined by

$$\Phi.c(\alpha) := c(\Phi^{-1}(\alpha))$$

for $\alpha \in G$. In the next corollary, we consider the semidirect product $\text{Aut}(G) \ltimes Z(G, \mathbb{T})$ with respect to this action. Recall that the product of $\text{Aut}(G) \ltimes Z(G, \mathbb{T})$ is

$$(\Phi_1, c_1) \cdot (\Phi_2, c_2) = (\Phi_1 \circ \Phi_2, (\Phi_2^{-1}.c_1) \cdot c_2)$$

for $(\Phi_1, c_1), (\Phi_2, c_2) \in \text{Aut}(G) \ltimes Z(G, \mathbb{T})$.

Corollary 3.2.2 ([14, Proposition 5.7]). *Let G be an effective étale groupoid. For $c \in Z(G, \mathbb{T})$ and $\Phi \in \text{Aut}(G)$, define $\varphi_{\Phi, c}: C_c(G) \rightarrow C_c(G)$ by*

$$\varphi_{\Phi, c}(f)(\alpha) := c(\Phi^{-1}(\alpha))f(\Phi^{-1}(\alpha))$$

for $f \in C_c(G)$ and $\alpha \in G$. Then the map $\varphi_{\Phi, c}$ extends to an element of $\text{Aut}_{C_0(G^{(0)})}(C_r^(G))$. Moreover, the map*

$$\Psi: \text{Aut}(G) \ltimes Z(G, \mathbb{T}) \ni (\Phi, c) \mapsto \varphi_{\Phi, c} \in \text{Aut}_{C_0(G^{(0)})}(C_r^*(G))$$

is a group isomorphism.

Remark 3.2.3. The above result is almost same as [14, Proposition 5.7 (1)]. In the proof of [14, Proposition 5.7], Renault's theory about Cartan C^* -subalgebras [17] is used. In this paper, we obtained the proof of Corollary 3.2.2 in a slightly more direct way without Renault's theory.

Proof of Corollary 3.2.2. We show that Ψ is a group homomorphism. Take

$$(\Phi_1, c_1), (\Phi_2, c_2) \in \text{Aut}(G) \ltimes Z(G, \mathbb{T}).$$

For $f \in C_c(G)$ and $\gamma \in G$, we have

$$\begin{aligned}
 & \varphi_{\Phi_1, c_1} \circ \varphi_{\Phi_2, c_2}(f)(\gamma) \\
 &= \varphi_{\Phi_1, c_1}(\varphi_{\Phi_2, c_2}(f))(\gamma) \\
 &= c_1(\Phi_1^{-1}(\gamma))\varphi_{\Phi_2, c_2}(f)(\Phi_1^{-1}(\gamma)) \\
 &= c_1(\Phi_1^{-1}(\gamma))c_2(\Phi_2^{-1}(\Phi_1^{-1}(\gamma)))f(\Phi_2^{-1}(\Phi_1^{-1}(\gamma))) \\
 &= (\Phi_2^{-1} \cdot c_1)((\Phi_1 \circ \Phi_2)^{-1}(\gamma))c_2((\Phi_1 \circ \Phi_2)^{-1}(\gamma))f((\Phi_1 \circ \Phi_2)^{-1}(\gamma)) \\
 &= \varphi_{\Phi_1 \circ \Phi_2, (\Phi_2^{-1} \cdot c_1) \cdot c_2}(f)(\gamma) \\
 &= \varphi_{(\Phi_1, c_1)(\Phi_2, c_2)}(f)(\gamma).
 \end{aligned}$$

Thus Ψ is a group homomorphism. By Theorem 3.1.1, Ψ is surjective. Remark that, if φ in Theorem 3.1.1 is an automorphism, then the corresponding invariant closed subset $F \subset G^{(0)}$ is the whole set $G^{(0)}$ since φ is injective on $C_0(G^{(0)})$ and F is given by

$$C_0(G^{(0)} \setminus F) = \ker \varphi \cap C_0(G^{(0)}).$$

To show that Ψ is injective, take $(\Phi_1, c_1), (\Phi_2, c_2) \in \text{Aut}(G) \ltimes Z(G, \mathbb{T})$ and assume $\varphi_{\Phi_1, c_1} = \varphi_{\Phi_2, c_2}$. If $\Phi_1 \neq \Phi_2$, then $\Phi_1^{-1}(\gamma) \neq \Phi_2^{-1}(\gamma)$ holds for some $\gamma \in G$. By Urysohn's lemma, there exists $f \in C_c(G)$ such that $f(\Phi_1^{-1}(\gamma)) \neq 0$ and $f(\Phi_2^{-1}(\gamma)) = 0$ hold. It follows that $\varphi_{\Phi_1, c_1}(f)(\gamma) \neq 0$ and $\varphi_{\Phi_2, c_2}(f)(\gamma) = 0$, which contradicts to $\varphi_{\Phi_1, c_1} = \varphi_{\Phi_2, c_2}$. Hence we obtain $\Phi_1 = \Phi_2$. To show $c_1 = c_2$, take $\gamma \in G$ and $f \in C_c(G)$ so that $f(\Phi_1^{-1}(\gamma)) = 1$. Then we have

$$c_1(\gamma) = \varphi_{\Phi_1, c_1}(f)(\gamma) = \varphi_{\Phi_1, c_2}(f)(\gamma) = c_2(\gamma).$$

Thus we obtain $c_1 = c_2$. Therefore we have shown that Ψ is injective. In conclusion, Ψ is a group isomorphism. \blacksquare

Finally, we investigate $\text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$. In the following propositions, we assume that an étale groupoid G is topologically principal. Recall that an étale Hausdorff groupoid G is effective if G is topologically principal. The converse is true if G is second countable.

Proposition 3.2.4. *Let G be a topologically principal étale groupoid and $\Phi: G \rightarrow G$ be a continuous groupoid automorphism. Assume that $\Phi(x) = x$ holds for all $x \in G^{(0)}$. Then $\Phi = \text{id}$.*

Proof. Put $A = \{x \in G^{(0)} \mid d^{-1}(x) \cap r^{-1}(x) = \{x\}\}$. Then $A \subset G^{(0)}$ is dense. Since $d: G \rightarrow G^{(0)}$ is an open map, $d^{-1}(A) \subset G$ is dense. We show that $\Phi(\alpha) = \alpha$ holds for all $\alpha \in d^{-1}(A)$. Note that we have

$$d(\Phi(\alpha)) = \Phi(d(\alpha)) = d(\alpha)$$

and $r(\Phi(\alpha)) = r(\alpha)$. Thus $(\Phi(\alpha)^{-1}, \alpha)$ is a composable pair and we have $\Phi(\alpha)^{-1}\alpha \in G_{d(\alpha)} \cap G^{d(\alpha)}$. Since $d(\alpha) \in A$, we obtain $\Phi(\alpha) = \alpha$. Since $d^{-1}(A) \subset G$ is dense, it follows that $\Phi(\alpha) = \alpha$ holds for all $\alpha \in G$. \blacksquare

Corollary 3.2.5. *Let G be a topologically principal étale groupoid and $\Phi_1, \Phi_2: G \rightarrow G$ be continuous groupoid automorphisms. If $\Phi_1|_{G^{(0)}} = \Phi_2|_{G^{(0)}}$, then $\Phi_1 = \Phi_2$ holds.*

Proof. Since $\Phi_1 \circ \Phi_2^{-1}(x) = x$ holds for all $x \in G^{(0)}$, we obtain $\Phi_1 \circ \Phi_2^{-1} = \text{id}$ by Proposition 3.2.4. Hence we obtain $\Phi_1 = \Phi_2$. ■

Corollary 3.2.6. *Let G be a topologically principal étale groupoid. Fix $(\Phi, c) \in \text{Aut}(G) \ltimes Z(G, \mathbb{T})$. Then $\Phi = \text{id}_G$ holds if and only if $\varphi_{\Phi, c} \in \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$ holds. In particular, the restriction of Ψ in Corollary 3.2.2 gives a group isomorphism*

$$\Psi|_{Z(G, \mathbb{T})}: Z(G, \mathbb{T}) \rightarrow \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$$

and $\text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$ is an abelian group.

Proof. It is clear that $\varphi_{\Phi, c} \in \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$ holds if $\Phi = \text{id}_G$. Assume $\varphi_{\Phi, c} \in \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$. Then we have

$$f(\Phi^{-1}(x)) = \varphi_{\Phi, c}(f)(x) = f(x)$$

for all $f \in C_0(G^{(0)})$ and $x \in G^{(0)}$. Hence, we obtain $\Phi^{-1}|_{G^{(0)}} = \text{id}_{G^{(0)}}$ and therefore $\Phi|_{G^{(0)}} = \text{id}_{G^{(0)}}$. By Proposition 3.2.4, we obtain $\Phi = \text{id}_G$. Now, the last assertion is clear. ■

A *-homomorphism $\varphi: C_r^*(G) \rightarrow C_r^*(G)$ automatically becomes an automorphism if φ fixes $C_0(G^{(0)})$ pointwisely as follows.

Corollary 3.2.7. *Let G be a topologically principal étale groupoid and $\varphi: C_r^*(G) \rightarrow C_r^*(G)$ be a *-homomorphism. Assume that $\varphi(f) = f$ holds for all $f \in C_0(G^{(0)})$. Then $\varphi \in \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$.*

Proof. By Theorem 3.1.1, there exists a locally homeomorphic groupoid homomorphism $\Phi: G \rightarrow G$ and $c \in Z(G, \mathbb{T})$ such that

$$\varphi(f)(\delta) = \sum_{\alpha \in \Phi^{-1}(\{\delta\})} c(\alpha) f(\alpha)$$

holds for all $f \in C_c(G)$ and $\delta \in H$. Here, in this situation, the corresponding closed invariant subset $F \subset G^{(0)}$ is the whole set $G^{(0)}$ since φ is injective on $C_0(G^{(0)})$. In addition, note that $\Phi|_{G^{(0)}}: G^{(0)} \rightarrow G^{(0)}$ is injective. Since G is effective and $\Phi|_{G^{(0)}}: G^{(0)} \rightarrow G^{(0)}$ is injective, $\Phi: G \rightarrow G$ is injective by the fundamental theorem on homomorphisms [10, Proposition 2.2]. Now, for all $f \in C_0(G^{(0)})$ and $x \in G^{(0)}$, we have

$$f(\Phi(x)) = \varphi(f)(\Phi(x)) = f(x).$$

Hence we obtain $\Phi|_{G^{(0)}} = \text{id}_{G^{(0)}}$ and therefore $\Phi = \text{id}_G$ by Proposition 3.2.4. Thus, for all $f \in C_c(G)$ and $\alpha \in G$, we obtain

$$\varphi(f)(\alpha) = c(\alpha) f(\alpha)$$

and therefore $\varphi \in \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$. ■

For a topological group H , we denote the abelianization of H by H^{ab} . Recall that H^{ab} is the quotient group of H by the closure of commutator subgroup $[\overline{H}, \overline{H}]$. In addition, we denote the quotient map by $\pi: H \rightarrow H^{\text{ab}}$ in the next corollary.

Corollary 3.2.8. *Let G be a topologically principal étale groupoid, H be a topological group and $\sigma: H \curvearrowright C_r^*(G)$ be an action such that*

$$\ker \sigma := \{s \in H \mid \alpha_s = \text{id}_{C_r^*(G)}\}$$

is closed in H . Assume that the fixed point algebra

$$C_r^*(G)^\sigma := \bigcap_{s \in H} \{x \in C_r^*(G) \mid \sigma_s(x) = x\}$$

contains $C_0(G^{(0)})$. Then there exists an action $\tilde{\sigma}: H^{\text{ab}} \curvearrowright C_r^(G)$ such that $\sigma_s = \tilde{\sigma}_{\pi(s)}$ holds for all $s \in H$.*

Proof. Since we assume $C_0(G^{(0)}) \subset C_r^*(G)^\sigma$, we have

$$\sigma_s \in \text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$$

for all $s \in H$. Hence, it follows $[H, H] \subset \ker \sigma$ since $\text{FAut}_{C_0(G^{(0)})}(C_r^*(G))$ is an abelian group by Corollary 3.2.6. Since we assume $\ker \sigma$ is closed, we obtain $\overline{[H, H]} \subset \ker \sigma$. Now, the existence of $\tilde{\sigma}$ follows from the fundamental group theory. ■

The assumption that $\ker \sigma \subset H$ is closed in Corollary 3.2.8 holds if we assume some continuity of σ . For example, if σ is a strongly continuous action, then $\ker \sigma \subset H$ is closed.

Acknowledgments. The author would like to thank Prof. Takeshi Katsura for his support and encouragement. The author is also grateful to Yuki Arano for fruitful discussions. In addition, the author would like to express his gratitude to the anonymous reviewer for the constructive comments and suggestions.

Funding. This work was supported by JST CREST Grant Number JPMJCR1913 and RIKEN Special Postdoctoral Researcher Program.

References

- [1] C. Anantharaman-Delaroche and J. Renault, *Amenable groupoids*. Monogr. Enseign. Math. 36, L'Enseignement Mathématique, Geneva, 2000 Zbl 0960.43003 MR 1799683
- [2] J. Brown, L. O. Clark, C. Farthing, and A. Sims, [Simplicity of algebras associated to étale groupoids](#). *Semigroup Forum* **88** (2014), no. 2, 433–452 Zbl 1304.46046 MR 3189105
- [3] J. H. Brown, R. Exel, A. H. Fuller, D. R. Pitts, and S. A. Reznikoff, [Intermediate \$C^*\$ -algebras of Cartan embeddings](#). *Proc. Amer. Math. Soc. Ser. B* **8** (2021), 27–41 Zbl 1471.46054 MR 4199728
- [4] N. P. Brown and N. Ozawa, *C^* -algebras and finite-dimensional approximations*. Grad. Stud. Math. 88, American Mathematical Society, Providence, RI, 2008 Zbl 1160.46001 MR 2391387

- [5] A. P. Donsig and D. R. Pitts, Coordinate systems and bounded isomorphisms. *J. Operator Theory* **59** (2008), no. 2, 359–416 Zbl [1199.46116](#) MR [2411050](#)
- [6] R. Exel, [Inverse semigroups and combinatorial \$C^*\$ -algebras](#). *Bull. Braz. Math. Soc. (N.S.)* **39** (2008), no. 2, 191–313 Zbl [1173.46035](#) MR [2419901](#)
- [7] R. Exel, Noncommutative Cartan subalgebras of C^* -algebras. *New York J. Math.* **17** (2011), 331–382 Zbl [1228.46061](#) MR [2811068](#)
- [8] J. Feldman and C. C. Moore, [Ergodic equivalence relations, cohomology, and von Neumann algebras. II](#). *Trans. Amer. Math. Soc.* **234** (1977), no. 2, 325–359 Zbl [0369.22010](#) MR [0578730](#)
- [9] F. Komura, [Quotients of étale groupoids and the abelianizations of groupoid \$C^*\$ -algebras](#). *J. Aust. Math. Soc.* **111** (2021), no. 1, 56–75 Zbl [1481.46051](#) MR [4282027](#)
- [10] F. Komura, [Invariant sets and normal subgroupoids of universal étale groupoids induced by congruences of inverse semigroups](#). *J. Aust. Math. Soc.* **113** (2022), no. 1, 99–118 Zbl [1512.20210](#) MR [4450914](#)
- [11] F. Komura, [Submodules of normalisers in groupoid \$C^*\$ -algebras and discrete group coactions](#). *Forum Math.* **37** (2025), no. 1, 121–133 Zbl [1557.46045](#) MR [4846653](#)
- [12] M. V. Lawson, *Inverse semigroups. The theory of partial symmetries*. World Scientific, River Edge, NJ, 1998 Zbl [1079.20505](#) MR [1694900](#)
- [13] K. Matsumoto and H. Matui, [Continuous orbit equivalence of topological Markov shifts and Cuntz–Krieger algebras](#). *Kyoto J. Math.* **54** (2014), no. 4, 863–877 Zbl [1320.46055](#) MR [3276420](#)
- [14] H. Matui, [Homology and topological full groups of étale groupoids on totally disconnected spaces](#). *Proc. Lond. Math. Soc. (3)* **104** (2012), no. 1, 27–56 Zbl [1325.19001](#) MR [2876963](#)
- [15] A. L. T. Paterson, *Groupoids, inverse semigroups, and their operator algebras*. Progr. Math. 170, Birkhäuser, Boston, MA, 1999 Zbl [0913.22001](#) MR [1724106](#)
- [16] J. Renault, *A groupoid approach to C^* -algebras*. Lecture Notes in Math. 793, Springer, Berlin, 1980 Zbl [0433.46049](#) MR [0584266](#)
- [17] J. Renault, [Cartan subalgebras in \$C^*\$ -algebras](#). *Irish Math. Soc. Bull.* (2008), no. 61, 29–63 Zbl [1175.46050](#) MR [2460017](#)
- [18] A. Sims, Hausdorff étale groupoids and their C^* -algebras. In *Operator algebras and dynamics: groupoids, crossed products, and Rokhlin dimension*, pp. 59–120, Adv. Courses Math. CRM Barcelona, Birkhäuser, Cham, 2020 Zbl [1495.46055](#)
- [19] R. Willett, [A non-amenable groupoid whose maximal and reduced \$C^*\$ -algebras are the same](#). *Münster J. Math.* **8** (2015), no. 1, 241–252 Zbl [1369.46064](#) MR [3549528](#)

Communicated by Wilhelm Winter

Received 1 April 2024; revised 10 July 2024.

Fuyuta Komura

Department of Basic Sciences, Faculty of Engineering, Kyushu Institute of Technology, 1-1 Sensui-cho, Tobata-ku, Kitakyushu-shi, Fukuoka, Kitakyushu 804-8550, Japan;
komura@mns.kyutech.ac.jp