



Giovanni Catino · Dario Daniele Monticelli

# Semilinear elliptic equations on manifolds with nonnegative Ricci curvature

Received 2 April 2022; revised 26 January 2024

**Abstract.** In this paper, we prove classification results for solutions to subcritical and critical semilinear elliptic equations with a nonnegative potential on noncompact manifolds with nonnegative Ricci curvature. We show in the subcritical case that all nonnegative solutions vanish identically. Moreover, under some natural assumptions, in the critical case we prove a strong rigidity result, namely we classify all nontrivial solutions showing that they exist only if the potential is constant and the manifold is isometric to the Euclidean space.

**Keywords:** semilinear elliptic equations, manifolds with nonnegative Ricci curvature, rigidity of solutions to PDEs.

## 1. Introduction

Let  $(M^n, g)$ ,  $n \geq 2$ , be a smooth complete (with no boundary), noncompact,  $n$ -dimensional Riemannian manifold with nonnegative Ricci curvature. In this paper, we consider nonnegative solutions to the semilinear elliptic equation

$$-\Delta_g u = Ku^p, \quad (1.1)$$

where  $\Delta_g$  is the Laplace–Beltrami operator and  $K$  is a smooth nonnegative function on  $M$ . We restrict our attention to the superlinear and subcritical case

$$1 < p < p_c, \quad \text{where} \quad p_c = \begin{cases} +\infty & \text{if } n = 2, \\ \frac{n+2}{n-2} & \text{if } n \geq 3, \end{cases}$$

and to the critical case,  $p = p_c = \frac{n+2}{n-2}$  when  $n \geq 3$ . In the case  $n = 2$ , we deal with solutions to the critical equation with exponential nonlinearity

$$-\Delta_g u = Ke^u. \quad (1.2)$$

---

Giovanni Catino: Dipartimento di Matematica, Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133 Milano, Italy; [giovanni.catino@polimi.it](mailto:giovanni.catino@polimi.it)

Dario Daniele Monticelli (corresponding author): Dipartimento di Matematica, Politecnico di Milano, Piazza Leonardo da Vinci 32, 20133 Milano, Italy; [dario.monticelli@polimi.it](mailto:dario.monticelli@polimi.it)

*Mathematics Subject Classification 2020:* 35J91 (primary); 35B33, 58J05, 53C21, 53C24 (secondary).

If we denote by  $R = R_g$  the scalar curvature of the metric  $g$ , it is known that these critical equations arise in the problem of prescribing the scalar curvature of a conformal metric when the original metric has zero scalar curvature. More precisely, if  $n \geq 3$ , then the scalar curvature of the metric  $\tilde{g} = u^{\frac{4}{n-2}} g$ ,  $u > 0$ , is given by

$$-\Delta_g u + \frac{n-2}{4(n-1)} R_g u = R_{\tilde{g}} u^{\frac{n+2}{n-2}}$$

while in dimension  $n = 2$ , the corresponding equation for the conformal change  $\tilde{g} = e^u g$  reads as

$$-\Delta_g u + R_g = R_{\tilde{g}} e^u.$$

It is natural to expect stronger classification results when  $K \geq 0$  which will be the case we study in this paper. In particular, we will first state our results in the simpler but geometrically relevant case  $K \equiv 1$ . This choice of  $K$  corresponds to the so called Yamabe problem, when  $(M^n, g)$  is scalar flat and hence Ricci flat. It is clear that solutions to (1.1) or (1.2) are trivial when  $M$  is compact.

In the Euclidean setting, problem (1.1) with  $K \equiv 1$  is now well understood. Gidas and Spruck [19] showed that the only nonnegative solution when  $1 < p < p_c$  is  $u \equiv 0$  on  $\mathbb{R}^n$ , via test functions argument. Indeed, this is a consequence of the following general result that they proved in the case  $n \geq 3$ .

**Theorem 1.1** ([19]). *Let  $(M^n, g)$  be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative solution of*

$$-\Delta u = u^p \quad \text{in } M \text{ with } 1 < p < p_c.$$

*Then*

$$u \equiv 0 \quad \text{on } M.$$

It is known that the same result holds also when  $n = 2$ . In Theorem 1.7, we provide a simpler proof of a more general result which includes Theorem 1.1 for every  $n \geq 2$ .

The explicit positive solutions to the critical equation

$$-\Delta u = u^{\frac{n+2}{n-2}} \quad \text{on } \mathbb{R}^n, \tag{1.3}$$

with  $n \geq 3$ , are given by

$$u(x) = \left( \frac{1}{a + b|x - x_0|^2} \right)^{\frac{n-2}{2}} \tag{1.4}$$

with  $a, b > 0$ ,  $1 = n(n-2)ab$  and  $x_0 \in \mathbb{R}^n$ . These functions were constructed by Aubin [1] and Talenti [26] as minimizers of

$$S_g(M) := \inf_{0 \neq u \in D^{1,2}(M)} \frac{\int_M |\nabla u|^2 dV_g}{(\int_M u^{\frac{2n}{n-2}} dV_g)^{\frac{n-2}{n}}},$$

where  $dV_g$  is the canonical volume element and

$$D^{1,2}(M) = \{u \in L^{\frac{2n}{n-2}}(M) : |\nabla u| \in L^2(M)\},$$

with  $M = \mathbb{R}^n$ . We note that if  $S_g(M) > 0$ , then it is the best constant in the Sobolev embedding. Caffarelli, Gidas and Spruck [6] (see also [13, 22]) proved that any solution to (1.3) is radial and hence given by (1.4). Essential tools in their proof are the moving planes method and the Kelvin transform that allow to reduce the problem to the study of (singular) solutions that have nice decaying properties at infinity. Previous results were proved by Gidas, Ni and Nirenberg [18] and Obata [24] under the additional assumption that  $u$  decays as  $|x|^{-(n-2)}$  at infinity. In the case  $n = 2$ , solutions to

$$\begin{cases} -\Delta u = e^u & \text{in } \mathbb{R}^2, \\ \int_{\mathbb{R}^2} e^u < +\infty \end{cases}$$

were classified by Chen and Li [13], who showed that

$$u(x) = \log \frac{1}{(a + b|x - x_0|^2)^2}$$

for some  $a, b > 0$  with  $1 = 8ab$  and  $x_0 \in \mathbb{R}^2$ . Their method relies on the moving plane method and on a previous result by Brezis and Merle [5] on the upper boundedness of solutions.

Extending such classification results for the critical equations to the case of more general Riemannian manifolds requires the introduction of different techniques than those used in the Euclidean setting, which strongly rely on the conformal invariance of the problem and the rich structure of the conformal group of the ambient space. In this paper, we study the problem in the natural setting of a complete noncompact Riemannian manifold  $(M, g)$  with nonnegative Ricci curvature, also allowing for the presence of a nonnegative potential function  $K$ .

Here and in the rest of the paper, we will denote by  $\text{Ric} = \text{Ric}_g$ ,  $R = R_g$ ,  $dV_g$  and  $r(\cdot)$  the Ricci curvature, the scalar curvature, the canonical Riemannian volume form and the geodesic distance from a fixed reference point of  $M$ , respectively.

The novelty of our approach consists in using a careful test functions argument which starts from the classical Bochner formula

$$\frac{1}{2} \Delta |\nabla f|^2 = |\nabla^2 f|^2 + \text{Ric}(\nabla f, \nabla f) + \langle \nabla f, \nabla \Delta f \rangle,$$

through which we are able to prove integral estimates involving the squared norm of the traceless Hessian

$$\mathring{\nabla}^2 f = \nabla^2 f - \frac{\Delta f}{n} g$$

of a suitable power of the solution, the Ricci tensor in the direction of the gradient of the solution and, in the subcritical case, some integral norms of the gradient of the solution. We show that all these quantities can be controlled by integrals on large annuli of

first-order derivatives of the solution which, in turn, can be controlled using the main equation. Under very general assumptions, we can then show that these quantities must vanish identically on  $M$  and this leads, in the subcritical case, to the triviality of the solution while, in the critical case, it yields the classification of nontrivial solutions and the rigidity of the ambient manifold, by using a characterization of conformal gradient vector fields. The starting point of our approach is partly reminiscent of the method used by Gidas and Spruck [19] for subcritical equations ( $n \geq 3$ ), Bidaut-Véron and Raoux [3] for subcritical systems and the first author with Castorina and Mantegazza [8] for subcritical parabolic equations.

In the first theorem, we deal with the case  $n \geq 3$  and finite energy solutions, i.e.,  $u \in D^{1,2}(M)$ .

**Theorem 1.2.** *Let  $(M^n, g)$ ,  $n \geq 3$ , be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative finite energy solution, i.e.,  $u \in D^{1,2}(M)$ , of*

$$-\Delta u = u^{\frac{n+2}{n-2}} \quad \text{in } M. \quad (1.5)$$

*Then either  $u \equiv 0$  on  $M$  or  $(M^n, g)$  is isometric to  $\mathbb{R}^n$  with the Euclidean metric and*

$$u(x) = \left( \frac{1}{a + b|x - x_0|^2} \right)^{\frac{n-2}{2}}$$

*for some  $a, b > 0$  with  $1 = n(n-2)ab$  and  $x_0 \in \mathbb{R}^n$ .*

An immediate consequence of this result is the following.

**Corollary 1.3.** *A complete noncompact nonflat  $n$ -dimensional ( $n \geq 3$ ) Riemannian manifold with nonnegative Ricci curvature does not admit any Sobolev minimizer of  $S_g(M)$ .*

In particular, if the manifold is also Ricci flat, it does not admit any Yamabe minimizer, i.e., a smooth function attaining the Yamabe constant

$$Y(M, [g]) = \inf_{0 \neq u \in C_0^\infty(M)} \frac{\int_M |\nabla u|^2 dV_g + \frac{n-2}{4(n-1)} \int_M R_g u^2 dV_g}{\left( \int_M u^{\frac{2n}{n-2}} dV_g \right)^{\frac{n-2}{n}}}.$$

An alternative proof of Corollary 1.3 can be recovered using a recent result by Brendle [4]. In fact, if  $S_g(M) = 0$ , then clearly it cannot be attained by any function in  $D^{1,2}(M)$ . If  $S_g(M) > 0$ , then  $(M^n, g)$  supports the Sobolev inequality and  $S_g(M)$  is the best constant. Hence, by a result of Carron [7] (see also [30, Proposition 2.4]),  $(M^n, g)$  has maximal volume growth, i.e., there exists  $C > 0$  such that

$$\text{Vol}_g B_\rho(x_0) \geq C\rho^n$$

for every  $x_0 \in M$ ,  $\rho > 0$ , and then one concludes using [4].

A result similar to Corollary 1.3 in the setting of Cartan–Hadamard manifolds (simply connected manifolds with nonpositive sectional curvature) has been recently obtained

assuming the validity of an optimal isoperimetric inequality on  $M$  (see [23]). We also point out a recent result obtained by Ciraolo, Figalli and Roncoroni [15] concerning the classification of finite energy solutions to the critical (anisotropic)  $p$ -Laplace equation on convex cones of  $\mathbb{R}^n$ , obtained via integral estimates and sharp a priori bounds.

In the second theorem, we consider the case  $n \geq 3$  and solutions which may not have finite energy, but satisfy a suitable condition at infinity.

**Theorem 1.4.** *Let  $(M^n, g)$ ,  $n \geq 3$ , be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative solution of*

$$-\Delta u = u^{\frac{n+2}{n-2}} \quad \text{in } M.$$

*If  $n \geq 4$ , also assume that, outside a compact set of  $M$ ,*

$$u(x) \leq Cr(x)^\alpha \quad \text{for some } \alpha < -\frac{(n-2)(n-6)}{2(n-4)}.$$

*Then either  $u \equiv 0$  on  $M$  or  $(M^n, g)$  is isometric to  $\mathbb{R}^n$  with the Euclidean metric and*

$$u(x) = \left( \frac{1}{a + b|x - x_0|^2} \right)^{\frac{n-2}{2}}$$

*for some  $a, b > 0$  with  $1 = n(n-2)ab$  and  $x_0 \in \mathbb{R}^n$ .*

We explicitly note that no assumption on the behavior of  $u$  at  $\infty$  is needed in Theorem 1.4 if  $n = 3$ . In particular, on  $\mathbb{R}^3$  we recover the full result by Caffarelli, Gidas and Spruck [6]. Moreover, we have the following.

**Corollary 1.5.** *A complete noncompact nonflat three-dimensional Riemannian manifold with nonnegative Ricci curvature does not admit any nonnegative, nontrivial solution of the critical equation*

$$-\Delta u = u^5.$$

In the case  $n = 4$ , Theorem 1.4 only needs that  $u$  is bounded above by  $r(x)$  to any power, i.e.,  $u$  has at most algebraic growth (with respect to the distance function), as  $r(x)$  tends to  $\infty$ . We also note that  $\alpha > -\frac{n-2}{2}$  for every  $n \geq 4$ , thus obtaining on a Riemannian manifold with nonnegative Ricci curvature an improvement of the classical results in  $\mathbb{R}^n$  by Gidas–Ni–Nirenberg [18] (see also Obata [24] in the case of manifolds and Caffarelli–Gidas–Spruck [6] in the case of  $\mathbb{R}^n$ ) in any dimension  $n \geq 3$ , where the authors assume that the solution decays as  $|x|^{-(n-2)}$  at infinity.

In our third theorem, we deal with the case  $n = 2$ . We have the following result.

**Theorem 1.6.** *Let  $(M^2, g)$  be a complete noncompact Riemannian surface with nonnegative scalar curvature. Let  $u \in C^2(M)$  be a solution of*

$$\begin{cases} -\Delta u = e^u, \\ \int_M e^u < +\infty. \end{cases}$$

Assume that, outside a compact set of  $M$ ,

$$u(x) \geq -4 \log r(x) - 2\gamma \log \log r(x), \quad \gamma \in [0, 1).$$

Then  $(M^2, g)$  is isometric to  $\mathbb{R}^2$  with the Euclidean metric and

$$u(x) = \log \frac{1}{(a + b|x - x_0|^2)^2}$$

for some  $a, b > 0$  with  $1 = 8ab$  and  $x_0 \in \mathbb{R}^2$ .

In contrast to the Euclidean case, on a general Riemannian surface with nonnegative curvature, to the best of our knowledge, there is no result concerning the behavior of a solution  $u$  at infinity. For this reason, we have to assume an a priori lower bound. A better lower bound implying our condition was proved on  $\mathbb{R}^2$  by Chen and Li [14], relying on [5] and the explicit expression of the Green's function. In particular, Theorem 1.6 generalizes the result obtained in  $\mathbb{R}^2$  by Chen and Li [13].

Our previous results are particular instances of more general theorems where we can allow for the presence of a nonnegative potential function  $K$ . The following theorem contains Theorem 1.1 as a particular case.

**Theorem 1.7.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative solution of*

$$-\Delta u = Ku^p \quad \text{in } M$$

with  $1 < p < p_c$ , where

$$p_c := \begin{cases} +\infty & \text{if } n = 2, \\ \frac{n+2}{n-2} & \text{if } n \geq 3 \end{cases}$$

and  $0 \not\equiv K \in C^2(M)$ ,  $K \geq 0$ . Moreover, if  $n \geq 3$ , we also assume  $\Delta K \geq 0$  on  $M$ , and if  $n \geq 4$ , we also assume  $K(x) \geq \frac{C}{r(x)^\sigma}$  outside a compact set of  $M$  for some  $C > 0$ ,  $\sigma < \frac{2}{n-3}$ . Then

$$u \equiv 0 \quad \text{on } M.$$

Theorem 1.7 improves the results by Gidas and Spruck [19, Theorems 4.1 and 6.1], removing an assumption on  $K$  and including the case  $n = 2$ . Moreover, as it is clear from the proof, when  $n \geq 4$ , the lower bound on  $K$  can be relaxed to

$$K(x) \geq \frac{C}{r(x)^\sigma} \quad \text{with } \sigma < \sigma^*(n, p)$$

for some explicit exponent  $\sigma^*(n, p) \geq \frac{2}{n-3}$ , see (3.7).

The next two theorems extend Theorems 1.2 and 1.4, respectively. In the first one, we consider finite energy solutions, i.e., such that

$$u \in D_K^{1,2}(M) := \{u : Ku^{\frac{2n}{n-2}} \in L^1(M), |\nabla u| \in L^2(M)\},$$

while in the second we deal with solutions with prescribed behavior at infinity.

**Theorem 1.8.** *Let  $(M^n, g)$ ,  $n \geq 3$ , be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative finite energy solution, i.e.,  $u \in D_K^{1,2}(M)$ , of*

$$-\Delta u = Ku^{\frac{n+2}{n-2}} \quad \text{in } M$$

*with  $0 \not\equiv K \in C^2(M)$ ,  $K \geq 0$  and  $\Delta K \geq 0$ . Assume that, outside a compact set of  $M$ ,*

(i)  $K(x) \leq C(1 + r(x)^2)$ , or

(ii)  $|\nabla K(x)| \leq \frac{C}{r(x)} K(x)$ .

*Then either  $u \equiv 0$  on  $M$  or  $(M^n, g)$  is isometric to  $\mathbb{R}^n$  with the Euclidean metric and*

$$u(x) = \left( \frac{1}{a + b|x - x_0|^2} \right)^{\frac{n-2}{2}}, \quad K \equiv n(n-2)ab$$

*for some  $a, b > 0$  and  $x_0 \in \mathbb{R}^n$ .*

We note that, under suitable conditions on the potential  $K$ , assuming  $Ku^{\frac{2n}{n-2}} \in L^1(M)$  is sufficient to deduce  $u \in D_K^{1,2}(M)$ , i.e.,  $u$  has finite energy. This is the case, in particular, if  $K \equiv 1$ . See Lemma 2.9.

Similarly to Corollary 1.3, we see that a complete noncompact nonflat  $n$ -dimensional ( $n \geq 3$ ) Riemannian manifold with nonnegative Ricci curvature does not admit any minimizer in  $D_K^{1,2}(M)$  of the corresponding weighted Sobolev quotient, with weight  $K$  satisfying the assumptions of Theorem 1.8.

**Theorem 1.9.** *Let  $(M^n, g)$ ,  $n \geq 3$ , be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative solution of*

$$-\Delta u = Ku^{\frac{n+2}{n-2}} \quad \text{in } M$$

*with  $0 \not\equiv K \in C^2(M)$ ,  $K \geq 0$  and  $\Delta K \geq 0$ . Assume that*

$$|\nabla K(x)| \leq \frac{C}{r(x)} K(x)$$

*outside a compact set of  $M$ . If  $n \geq 4$ , also assume that*

$$u \leq Cr^\alpha \quad \text{and} \quad K(x) \geq \frac{C}{r(x)^\sigma}$$

*outside a compact set of  $M$ , for some*

$$\alpha < \max \left\{ -\frac{(n-2)(n-4)}{2(n-3)}, -\frac{(n-2)^2\sigma + 2(n-2)(n-6)}{4(n-4)} \right\}.$$

*Then either  $u \equiv 0$  on  $M$  or  $(M^n, g)$  is isometric to  $\mathbb{R}^n$  with the Euclidean metric and*

$$u(x) = \left( \frac{1}{a + b|x - x_0|^2} \right)^{\frac{n-2}{2}}, \quad K \equiv n(n-2)ab$$

*for some  $a, b > 0$  and  $x_0 \in \mathbb{R}^n$ .*

From this theorem, we have the following.

**Corollary 1.10.** *A complete noncompact nonflat three-dimensional Riemannian manifold with nonnegative Ricci curvature does not admit any nonnegative, nontrivial solution of the critical equation*

$$-\Delta u = Ku^5$$

*with  $0 \not\equiv K \in C^2(M)$ ,  $K \geq 0$ ,  $\Delta K \geq 0$  and  $|\nabla K(x)| \leq \frac{C}{r(x)} K(x)$  outside a compact set of  $M$ .*

In the proofs of Theorems 1.7 and 1.9, we use the Bishop–Gromov volume estimate which ensures that the volume of geodesic balls of radius  $\rho$  grows at most as  $C\rho^n$  as  $\rho$  tends to infinity. As it is clear from the proofs, a slower rate of growth allows for weaker assumptions on  $u$  and/or  $K$ . We leave the details to the interested reader.

Our last theorem deals with the case  $n = 2$  and generalizes Theorem 1.6.

**Theorem 1.11.** *Let  $(M^2, g)$  be a complete noncompact Riemannian surface with nonnegative scalar curvature. Let  $u \in C^2(M)$  be a solution of*

$$\begin{cases} -\Delta u = Ke^u, \\ \int_M Ke^u < +\infty \end{cases}$$

*with  $0 \not\equiv K \in C^2(M)$ ,  $K \geq 0$  and  $\Delta K \geq 0$ . Assume that, outside a compact set of  $M$ ,*

$$u(x) \geq -4 \log r(x) - 2\gamma \log \log r(x), \quad \gamma \in [0, 1),$$

*and, for some  $C > 0$ ,*

- (i)  $K(x) \leq C(1 + r(x)^2)$  and  $u(x) \leq C \log r(x)$ , or
- (ii)  $|\nabla K(x)| \leq \frac{C}{r(x)} K(x)$ .

*Then  $(M^2, g)$  is isometric to  $\mathbb{R}^2$  with the Euclidean metric and*

$$u(x) = \log \frac{1}{(a + b|x - x_0|^2)^2}, \quad K \equiv 8ab$$

*for some  $a, b > 0$  and  $x_0 \in \mathbb{R}^2$ .*

It is interesting to note that Chen and Li in [14] exhibited the explicit radial solutions

$$u_\alpha(r) = (2 + \alpha) \log \frac{4}{4 + r^2} \quad \text{in } \mathbb{R}^2 \text{ for } K(r) = (2 + \alpha) \left( \frac{4 + r^2}{4} \right)^\alpha,$$

for every  $\alpha \in \mathbb{R}$ . Note that in these examples  $\Delta K \geq 0$  in  $\mathbb{R}^2$  if and only if  $\alpha \geq 0$ , while our lower bound on  $u_\alpha$  is satisfied if and only if  $\alpha \leq 0$ . In the same paper, the authors provide some sufficient conditions on  $K$  ensuring the validity of upper and lower bounds for solutions on  $\mathbb{R}^2$ .

Finally, to the best of our knowledge, Theorems 1.8, 1.9 and 1.11 are new even in the Euclidean setting. In the more general Riemannian setting, we point out some existence

results of variational solutions for the Yamabe equation under conditions on the Yamabe constant and the Yamabe constant at infinity, see [21, 31]. We explicitly note that the Yamabe equation reduces to (1.5) when  $n \geq 3$  and the manifold is scalar flat. Of course, the conditions in the cited references cannot hold on manifolds with nonnegative Ricci curvature.

The paper is organized as follows: in Section 2, we collect all the technical lemmas that we will need in the proofs of our main theorems; in Section 3, we prove Theorem 1.7 concerning the nonexistence of nontrivial solutions in the subcritical case; in Sections 4 and 5, we prove Theorems 1.8 and 1.9 which deal with the critical case, when  $n \geq 3$ ; finally, in Section 6, we prove Theorem 1.11 on the critical equation on surfaces.

*Added note:* After this paper was submitted for publication (on March 2022), several very interesting results on this topic appeared in the literature. We explicitly mention Fogagnolo–Malchiodi–Mazzieri [17], where the authors studied the same problem with a different technique and Catino–Monticelli–Roncoroni [11], Ou [25] and Vétois [28], improving this method in order to deal with the  $p$ -Laplacian. We also cite Catino–Li–Monticelli–Roncoroni [9] and Flynn–Vétois [16], where similar techniques are used to study the uniqueness of positive solutions of the critical semilinear equation for the sublaplacian on the Heisenberg group.

## 2. Preliminary lemmas

### 2.1. Part I

We collect here all technical lemmas that we will need in the study of equation (1.1). We start showing the following key technical identity. We note that in the Euclidean case, Bidaut-Véron and Raoux in [3, Lemma 3.1] showed an estimate without including two extra terms that are crucial in proving our result in the critical case.

**Lemma 2.1.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a Riemannian manifold. For any positive function  $w \in C^2(M)$ , any nonnegative  $\eta \in C_c^2(M)$  and any real numbers  $d, m \in \mathbb{R}$  such that  $d \neq m + 2$ , the following identity holds:*

$$\begin{aligned} & \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n} \int_M \eta w^{m-2} |\nabla w|^4 - \frac{n-1}{n} \int_M \eta w^m (\Delta w)^2 \\ & - \frac{2(n-1)m + (n+2)d}{2n} \int_M \eta w^{m-1} |\nabla w|^2 \Delta w \\ & + \frac{4}{(m+2-d)^2} \int_M \eta w^d |\overset{\circ}{\nabla}^2 w^{\frac{m+2-d}{2}}|^2 \\ & + \frac{4}{(m+2-d)^2} \int_M \eta w^d \text{Ric}(\nabla w^{\frac{m+2-d}{2}}, \nabla w^{\frac{m+2-d}{2}}) \\ & = \frac{m+d}{2} \int_M w^{m-1} |\nabla w|^2 \langle \nabla w, \nabla \eta \rangle + \int_M w^m \Delta w \langle \nabla w, \nabla \eta \rangle + \frac{1}{2} \int_M w^m |\nabla w|^2 \Delta \eta. \end{aligned}$$

*Proof.* Let  $f := w^t$  for some  $t \neq 0$ . Then

$$\nabla f = tw^{t-1}\nabla w, \quad \Delta f = tw^{t-1}\Delta w + t(t-1)w^{t-2}|\nabla w|^2.$$

Now we use the Bochner formula

$$\begin{aligned} \frac{1}{2}\Delta|\nabla f|^2 &= |\nabla^2 f|^2 + \text{Ric}(\nabla f, \nabla f) + \langle \nabla f, \nabla \Delta f \rangle \\ &= |\overset{\circ}{\nabla}^2 f|^2 + \frac{1}{n}(\Delta f)^2 + \text{Ric}(\nabla f, \nabla f) + \langle \nabla f, \nabla \Delta f \rangle \\ &= |\overset{\circ}{\nabla}^2 w^t|^2 + \text{Ric}(\nabla w^t, \nabla w^t) + \frac{t^2}{n}w^{2t-2}(\Delta w)^2 + \frac{t^2(t-1)^2}{n}w^{2t-4}|\nabla w|^4 \\ &\quad + \frac{2t^2(t-1)}{n}w^{2t-3}|\nabla w|^2\Delta w + t^2(t-1)w^{2t-3}|\nabla w|^2\Delta w \\ &\quad + t^2w^{2t-2}\langle \nabla w, \nabla \Delta w \rangle + t^2(t-1)w^{2t-3}\langle \nabla w, \nabla |\nabla w|^2 \rangle \\ &\quad + t^2(t-1)(t-2)w^{2t-4}|\nabla w|^4. \end{aligned}$$

Let  $d \in \mathbb{R}$ . Multiplying the above equation by  $t^{-2}\eta w^d$  and integrating over  $M$ , we obtain the following:

$$\begin{aligned} &\left( \frac{(t-1)^2}{n} + (t-1)(t-2) \right) \int_M \eta w^{d+2t-4}|\nabla w|^4 \\ &\quad + \frac{(t-1)(n+2)}{n} \int_M \eta w^{d+2t-3}|\nabla w|^2\Delta w + \int_M \eta w^{d+2t-2}\langle \nabla w, \nabla \Delta w \rangle \\ &\quad + \frac{1}{n} \int_M \eta w^{d+2t-2}(\Delta w)^2 + (t-1) \int_M \eta w^{d+2t-3}\langle \nabla w, \nabla |\nabla w|^2 \rangle \\ &\quad - \frac{1}{2} \int_M \eta w^d \Delta(w^{2t-2}|\nabla w|^2) + t^{-2} \int_M \eta w^d |\overset{\circ}{\nabla}^2 w^t|^2 \\ &\quad + t^{-2} \int_M \eta w^d \text{Ric}(\nabla w^t, \nabla w^t) = 0. \end{aligned}$$

Integrating by parts, we obtain

$$\begin{aligned} \int_M \eta w^{d+2t-2}\langle \nabla w, \nabla \Delta w \rangle &= - \int_M w^{d+2t-2}\Delta w \langle \nabla w, \nabla \eta \rangle \\ &\quad - (d+2t-2) \int_M \eta w^{d+2t-3}|\nabla w|^2\Delta w \\ &\quad - \int_M \eta w^{d+2t-2}(\Delta w)^2, \\ \int_M \eta w^{d+2t-3}\langle \nabla w, \nabla |\nabla w|^2 \rangle &= - \int_M w^{d+2t-3}|\nabla w|^2 \langle \nabla w, \nabla \eta \rangle \\ &\quad - (d+2t-3) \int_M \eta w^{d+2t-4}|\nabla w|^4 \\ &\quad - \int_M \eta w^{d+2t-3}|\nabla w|^2\Delta w, \end{aligned}$$

$$\begin{aligned} \int_M \eta w^d \Delta(w^{2t-2} |\nabla w|^2) &= \int_M w^{d+2t-2} |\nabla w|^2 \Delta \eta + 2d \int_M w^{d+2t-3} |\nabla w|^2 \langle \nabla w, \nabla \eta \rangle \\ &\quad + d(d-1) \int_M \eta w^{d+2t-4} |\nabla w|^4 \\ &\quad + d \int_M \eta w^{d+2t-3} |\nabla w|^2 \Delta w. \end{aligned}$$

Substituting in the above identity, rearranging terms and setting  $m := d + 2t - 2$ , we conclude.  $\blacksquare$

In the next lemma, we apply the previous identity to positive solutions of equation (1.1).

**Lemma 2.2.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a Riemannian manifold. For any positive solution  $u \in C^2(M)$  of*

$$-\Delta u = Ku^p$$

*with  $p \in \mathbb{R}$ ,  $K \in C^2(M)$ , any nonnegative  $\psi \in C_c^2(M)$  and any real numbers  $q \geq 2$ ,  $d, m \in \mathbb{R}$  such that  $d \neq m + 2$ ,  $m + p + 1 \neq 0$ , the following identity holds:*

$$\begin{aligned} \alpha \int_M \psi^q u^{m-2} |\nabla u|^4 + \beta \int_M \psi^q Ku^{m+p-1} |\nabla u|^2 + \frac{n-1}{n(m+p+1)} \int_M \psi^q \Delta Ku^{m+p+1} \\ + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d |\overset{\circ}{\nabla}^2 u^{\frac{m+2-d}{2}}|^2 \\ + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d \text{Ric}(\nabla u^{\frac{m+2-d}{2}}, \nabla u^{\frac{m+2-d}{2}}) \\ = \frac{n-2}{n} \int_M Ku^{m+p} \langle \nabla u, \nabla \psi^q \rangle + \frac{m+d}{2} \int_M u^{m-1} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle \\ + \frac{1}{2} \int_M u^m |\nabla u|^2 \Delta \psi^q + \frac{n-1}{n(m+p+1)} \int_M \Delta \psi^q Ku^{m+p+1}, \end{aligned}$$

where  $\alpha = \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n}$  and  $\beta = \frac{(n+2)d - 2(n-1)p}{2n}$ .

*Proof.* Applying Lemma 2.1 to  $w = u$  and  $\eta = \psi^q$ , for any real numbers  $d, m$  with  $d \neq m + 2$ , we get

$$\begin{aligned} \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n} \int_M \psi^q u^{m-2} |\nabla u|^4 - \frac{n-1}{n} \int_M \psi^q u^m (\Delta u)^2 \\ - \frac{2(n-1)m + (n+2)d}{2n} \int_M \psi^q u^{m-1} |\nabla u|^2 \Delta u \\ + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d |\overset{\circ}{\nabla}^2 u^{\frac{m+2-d}{2}}|^2 \\ + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d \text{Ric}(\nabla u^{\frac{m+2-d}{2}}, \nabla u^{\frac{m+2-d}{2}}) \\ = \frac{m+d}{2} \int_M u^{m-1} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle + \int_M u^m \Delta u \langle \nabla u, \nabla \psi^q \rangle + \frac{1}{2} \int_M u^m |\nabla u|^2 \Delta \psi^q. \end{aligned}$$

Hence, using the equation  $\Delta u = -Ku^p$ , we obtain

$$\begin{aligned}
& \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n} \int_M \psi^q u^{m-2} |\nabla u|^4 \\
& + \frac{n-1}{n} \int_M \psi^q Ku^{m+p} \Delta u + \frac{2(n-1)m + (n+2)d}{2n} \int_M \psi^q Ku^{m+p-1} |\nabla u|^2 \\
& + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d |\overset{\circ}{\nabla}^2 u^{\frac{m+2-d}{2}}|^2 \\
& + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d \operatorname{Ric}(\nabla u^{\frac{m+2-d}{2}}, \nabla u^{\frac{m+2-d}{2}}) \\
& = \frac{m+d}{2} \int_M u^{m-1} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle - \int_M Ku^{m+p} \langle \nabla u, \nabla \psi^q \rangle \\
& + \frac{1}{2} \int_M u^m |\nabla u|^2 \Delta \psi^q. \tag{2.1}
\end{aligned}$$

Integrating by parts in the first integral in the second line above, we get

$$\begin{aligned}
\frac{n-1}{n} \int_M \psi^q Ku^{m+p} \Delta u & = - \frac{n-1}{n} \int_M Ku^{m+p} \langle \nabla u, \nabla \psi^q \rangle \\
& - \frac{(m+p)(n-1)}{n} \int_M K \psi^q u^{m+p-1} |\nabla u|^2 \\
& - \frac{n-1}{n} \int_M \psi^q u^{m+p} \langle \nabla K, \nabla u \rangle,
\end{aligned}$$

and

$$\begin{aligned}
& - \frac{n-1}{n} \int_M \psi^q u^{m+p} \langle \nabla K, \nabla u \rangle \\
& = - \frac{n-1}{n(m+p+1)} \int_M \psi^q \langle \nabla K, \nabla u^{m+p+1} \rangle \\
& = \frac{n-1}{n(m+p+1)} \int_M \langle \nabla \psi^q, \nabla K \rangle u^{m+p+1} + \frac{n-1}{n(m+p+1)} \int_M \psi^q \Delta K u^{m+p+1} \\
& = - \frac{n-1}{n(m+p+1)} \int_M \Delta \psi^q Ku^{m+p+1} - \frac{n-1}{n} \int_M Ku^{m+p} \langle \nabla \psi^q, \nabla u \rangle \\
& + \frac{n-1}{n(m+p+1)} \int_M \psi^q \Delta K u^{m+p+1}.
\end{aligned}$$

Thus, substituting and setting

$$\begin{aligned}
I_1 & = \int_M Ku^{m+p} \langle \nabla u, \nabla \psi^q \rangle, \\
I_2 & = \int_M u^{m-1} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle, \\
I_3 & = \int_M u^m |\nabla u|^2 \Delta \psi^q,
\end{aligned}$$

equality (2.1) becomes

$$\begin{aligned}
& \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n} \int_M \psi^q u^{m-2} |\nabla u|^4 \\
& + \frac{n-1}{n(m+p+1)} \int_M \psi^q \Delta K u^{m+p+1} \\
& - 2 \frac{n-1}{n} I_1 - \frac{(m+p)(n-1)}{n} \int_M \psi^q K u^{m+p-1} |\nabla u|^2 \\
& + \frac{2(n-1)m + (n+2)d}{2n} \int_M \psi^q K u^{m+p-1} |\nabla u|^2 \\
& - \frac{n-1}{n(m+p+1)} \int_M \Delta \psi^q K u^{m+p+1} + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d |\overset{\circ}{\nabla}{}^2 u^{\frac{m+2-d}{2}}|^2 \\
& + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d \operatorname{Ric}(\nabla u^{\frac{m+2-d}{2}}, \nabla u^{\frac{m+2-d}{2}}) = \frac{m+d}{2} I_2 - I_1 + \frac{1}{2} I_3,
\end{aligned}$$

Hence, rearranging and simplifying, we conclude

$$\begin{aligned}
& \alpha \int_M \psi^q u^{m-2} |\nabla u|^4 + \beta \int_M \psi^q K u^{m+p-1} |\nabla u|^2 \\
& + \frac{n-1}{n(m+p+1)} \int_M \psi^q \Delta K u^{m+p+1} + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d |\overset{\circ}{\nabla}{}^2 u^{\frac{m+2-d}{2}}|^2 \\
& + \frac{4}{(m+2-d)^2} \int_M \psi^q u^d \operatorname{Ric}(\nabla u^{\frac{m+2-d}{2}}, \nabla u^{\frac{m+2-d}{2}}) \\
& = \frac{n-2}{n} I_1 + \frac{m+d}{2} I_2 + \frac{1}{2} I_3 + \frac{n-1}{n(m+p+1)} \int_M \Delta \psi^q K u^{m+p+1},
\end{aligned}$$

where  $\alpha = \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n}$  and  $\beta = \frac{(n+2)d - 2(n-1)p}{2n}$ .  $\blacksquare$

In the subcritical case  $1 < p < p_c = \frac{n+2}{n-2}$ , we neglect some nonnegative terms in Lemma 2.2, thus obtaining the following integral gradient estimate.

**Corollary 2.3.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a Riemannian manifold with  $\operatorname{Ric} \geq 0$ . For any positive solution  $u \in C^2(M)$  of*

$$-\Delta u = K u^p$$

*with  $p \in \mathbb{R}$ ,  $K \in C^2(M)$ ,  $\Delta K \geq 0$ , any nonnegative  $\psi \in C_c^2(M)$  and any real numbers  $q \geq 2$ ,  $d, m \in \mathbb{R}$  such that  $m + p + 1 > 0$  the following estimate holds:*

$$\begin{aligned}
& \alpha \int_M \psi^q u^{m-2} |\nabla u|^4 + \beta \int_M \psi^q K u^{m+p-1} |\nabla u|^2 \\
& \leq \frac{n-2}{n} \int_M K u^{m+p} \langle \nabla u, \nabla \psi^q \rangle + \frac{m+d}{2} \int_M u^{m-1} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle \\
& + \frac{1}{2} \int_M u^m |\nabla u|^2 \Delta \psi^q + \frac{n-1}{n(m+p+1)} \int_M \Delta \psi^q K u^{m+p+1},
\end{aligned}$$

where  $\alpha = \frac{2(n-m)d - (n-1)(m^2 + d^2)}{4n}$  and  $\beta = \frac{(n+2)d - 2(n-1)p}{2n}$ .

For  $n \geq 3$ , in the critical case  $p = p_c = \frac{n+2}{n-2}$ , by choosing

$$m = -\frac{2}{n-2}, \quad d = \frac{2(n-1)}{n-2}$$

in Lemma 2.2, we get  $\alpha = \beta = 0$ , thus obtaining the following identity.

**Corollary 2.4.** *Let  $(M^n, g)$ ,  $n \geq 3$ , be a Riemannian manifold. For any positive solution  $u \in C^2(M)$  of*

$$-\Delta u = Ku^{\frac{n+2}{n-2}}$$

*with  $K \in C^2(M)$ , any nonnegative  $\psi \in C_c^2(M)$  and any real number  $q \geq 2$ , the following identity holds:*

$$\begin{aligned} & \frac{(n-2)^2}{4} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 \\ & + \frac{(n-2)^2}{4} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) + \frac{n-2}{2n} \int_M \psi^q \Delta K u^{2\frac{n-1}{n-2}} \\ & = \frac{n-2}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^q \rangle + \int_M u^{-\frac{n}{n-2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle \\ & + \frac{1}{2} \int_M u^{-\frac{2}{n-2}} |\nabla u|^2 \Delta \psi^q + \frac{n-2}{2n} \int_M \Delta \psi^q K u^{2\frac{n-1}{n-2}}, \end{aligned}$$

In the next lemma, we estimate some of the terms in the right-hand side of the above identity in terms of controlled quantities. This inequality will be useful in the critical case.

**Lemma 2.5.** *Let  $(M^n, g)$ ,  $n \geq 3$ , be a Riemannian manifold. For any positive solution  $u \in C^2(M)$  of*

$$-\Delta u = Ku^{\frac{n+2}{n-2}}$$

*with  $K \in C^0(M)$ , any nonnegative  $\psi \in C_c^2(M)$  and any real numbers  $q \geq 2$ ,  $\varepsilon > 0$  the following estimate holds:*

$$\begin{aligned} & -\frac{1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^q \rangle + \int_M u^{-\frac{n}{n-2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle + \frac{1}{2} \int_M u^{-\frac{2}{n-2}} |\nabla u|^2 \Delta \psi^q \\ & \leq \frac{q(n-2)^2 \varepsilon}{8} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 + \frac{q}{2\varepsilon} \int_M \psi^{q-2} u^{-\frac{2}{n-2}} |\nabla u|^2 |\nabla \psi|^2. \end{aligned}$$

*Proof.* Let

$$\begin{aligned} I_1 &= \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^q \rangle, \\ I_2 &= \int_M u^{-\frac{n}{n-2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle, \\ I_3 &= \int_M u^{-\frac{2}{n-2}} |\nabla u|^2 \Delta \psi^q. \end{aligned}$$

Integrating by parts, we obtain

$$\begin{aligned}
I_2 &= \frac{(n-2)^3}{8(n-1)} \int_M |\nabla u^{-\frac{2}{n-2}}|^2 \langle \nabla u^{\frac{2(n-1)}{n-2}}, \nabla \psi^q \rangle \\
&= -\frac{(n-2)^3}{8(n-1)} \int_M u^{\frac{2(n-1)}{n-2}} \langle \nabla |\nabla u^{-\frac{2}{n-2}}|^2, \nabla \psi^q \rangle \\
&\quad - \frac{(n-2)^3}{8(n-1)} \int_M u^{\frac{2(n-1)}{n-2}} |\nabla u^{-\frac{2}{n-2}}|^2 \Delta \psi^q \\
&= -\frac{(n-2)^3}{4(n-1)} \int_M u^{\frac{2(n-1)}{n-2}} \nabla^2 u^{-\frac{2}{n-2}} \langle \nabla u^{-\frac{2}{n-2}}, \nabla \psi^q \rangle - \frac{n-2}{2(n-1)} I_3 \\
&= -\frac{(n-2)^3}{4(n-1)} \int_M u^{\frac{2(n-1)}{n-2}} \overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}} \langle \nabla u^{-\frac{2}{n-2}}, \nabla \psi^q \rangle \\
&\quad - \frac{(n-2)^3}{4n(n-1)} \int_M u^{\frac{2(n-1)}{n-2}} \Delta u^{-\frac{2}{n-2}} \langle \nabla u^{-\frac{2}{n-2}}, \nabla \psi^q \rangle - \frac{n-2}{2(n-1)} I_3.
\end{aligned}$$

Since

$$\Delta u^{-\frac{2}{n-2}} = \frac{2}{n-2} K u^{\frac{2}{n-2}} + \frac{2n}{(n-2)^2} u^{-\frac{2(n-1)}{n-2}} |\nabla u|^2,$$

using Young's inequality on the first integral in the last equality, we obtain

$$\begin{aligned}
I_2 &\leq \frac{q(n-2)^3 \varepsilon}{8(n-1)} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 \\
&\quad + \frac{q(n-2)^3}{8(n-1)\varepsilon} \int_M u^{\frac{2(n-1)}{n-2}} \psi^{q-2} |\nabla u^{-\frac{2}{n-2}}|^2 |\nabla \psi|^2 \\
&\quad - \frac{(n-2)^3}{4n(n-1)} \int_M u^{\frac{2(n-1)}{n-2}} \left( \frac{2}{n-2} K u^{\frac{2}{n-2}} + \frac{2n}{(n-2)^2} u^{-\frac{2(n-1)}{n-2}} |\nabla u|^2 \right) \\
&\quad \times \langle \nabla u^{-\frac{2}{n-2}}, \nabla \psi^q \rangle - \frac{n-2}{2(n-1)} I_3 \\
&= \frac{q(n-2)^3 \varepsilon}{8(n-1)} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} \left| \overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}} \right|^2 \\
&\quad + \frac{q(n-2)^3}{8(n-1)\varepsilon} \int_M u^{\frac{2(n-1)}{n-2}} \psi^{q-2} |\nabla u^{-\frac{2}{n-2}}|^2 |\nabla \psi|^2 \\
&\quad + \frac{n-2}{n(n-1)} I_1 + \frac{1}{n-1} I_2 - \frac{n-2}{2(n-1)} I_3.
\end{aligned}$$

Rearranging terms, we conclude. ■

## 2.2. Part II

We collect here all technical lemmas that we will need in the study of equation (1.2). We start showing the following key technical identity, which is the counterpart of Lemma 2.1.

**Lemma 2.6.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a Riemannian manifold. For any function  $w \in C^2(M)$ , any nonnegative  $\eta \in C_c^2(M)$  and any real numbers  $d, m \in \mathbb{R} \setminus \{0\}$ , the following identity holds:*

$$\begin{aligned}
& - \left( \frac{(n-1)m^2}{n} + md + \frac{d^2}{2} \right) \int_M \eta e^{(2m+d)w} |\nabla w|^4 \\
& - \frac{n-1}{n} \int_M \eta e^{(2m+d)w} (\Delta w)^2 \\
& - \left( \frac{2(n-1)m}{n} + \frac{3}{2}d \right) \int_M \eta e^{(2m+d)w} |\nabla w|^2 \Delta w \\
& + \frac{1}{m^2} \int_M \eta e^{dw} |\overset{\circ}{\nabla}{}^2 e^{mw}|^2 + \frac{1}{m^2} \int_M \eta e^{dw} \operatorname{Ric}(\nabla e^{mw}, \nabla e^{mw}) \\
& = (m+d) \int_M e^{(2m+d)w} |\nabla w|^2 \langle \nabla w, \nabla \eta \rangle + \int_M e^{(2m+d)w} \Delta w \langle \nabla w, \nabla \eta \rangle \\
& + \frac{1}{2} \int_M e^{(2m+d)w} |\nabla w|^2 \Delta \eta.
\end{aligned}$$

*Proof.* For  $f = e^{mw}$ , we have

$$\nabla f = m e^{mw} \nabla w, \quad \Delta f = m e^{mw} \Delta w + m^2 e^{mw} |\nabla w|^2.$$

We use the Bochner formula

$$\begin{aligned}
\frac{1}{2} \Delta |\nabla f|^2 &= |\nabla^2 f|^2 + \operatorname{Ric}(\nabla f, \nabla f) + \langle \nabla f, \nabla \Delta f \rangle \\
&= |\overset{\circ}{\nabla}{}^2 f|^2 + \frac{1}{n} (\Delta f)^2 + \operatorname{Ric}(\nabla f, \nabla f) + \langle \nabla f, \nabla \Delta f \rangle \\
&= |\overset{\circ}{\nabla}{}^2 e^{mw}|^2 + \operatorname{Ric}(\nabla e^{mw}, \nabla e^{mw}) + \frac{m^2}{n} e^{2mw} (\Delta w)^2 + \frac{m^4}{n} e^{2mw} |\nabla w|^4 \\
&+ \frac{2m^3}{n} e^{2mw} |\nabla w|^2 \Delta w + m^3 e^{2mw} \Delta w |\nabla w|^2 + m^2 e^{2mw} \langle \nabla w, \nabla \Delta w \rangle \\
&+ m^4 e^{2mw} |\nabla w|^4 + m^3 e^{2mw} \langle \nabla w, \nabla |\nabla w|^2 \rangle.
\end{aligned}$$

Now we multiply by  $m^{-2} \eta e^{dw}$  and integrate over  $M$  to obtain

$$\begin{aligned}
& m^{-2} \int_M \eta e^{dw} |\overset{\circ}{\nabla}{}^2 e^{mw}|^2 + m^{-2} \int_M \eta e^{dw} \operatorname{Ric}(\nabla e^{mw}, \nabla e^{mw}) \\
& + \frac{1}{n} \int_M \eta e^{(2m+d)w} (\Delta w)^2 + \frac{(n+1)m^2}{n} \int_M \eta e^{(2m+d)w} |\nabla w|^4 \\
& + \frac{(2+n)m}{n} \int_M \eta e^{(2m+d)w} |\nabla w|^2 \Delta w + \int_M \eta e^{(2m+d)w} \langle \nabla w, \nabla \Delta w \rangle \\
& + m \int_M \eta e^{(2m+d)w} \langle \nabla w, \nabla |\nabla w|^2 \rangle - \frac{1}{2} \int_M \eta e^{dw} \Delta(e^{2mw} |\nabla w|^2) = 0. \quad (2.2)
\end{aligned}$$

Integrating by parts,

$$\begin{aligned}
\int_M \eta e^{(2m+d)w} \langle \nabla w, \nabla \Delta w \rangle &= - \int_M e^{(2m+d)w} \Delta w \langle \nabla \eta, \nabla w \rangle - \int_M \eta e^{(2m+d)w} (\Delta w)^2 \\
&\quad - (2m+d) \int_M \eta e^{(2m+d)w} \Delta w |\nabla w|^2, \\
\int_M \eta e^{(2m+d)w} \langle \nabla w, \nabla |\nabla w|^2 \rangle &= - \int_M e^{(2m+d)w} |\nabla w|^2 \langle \nabla \eta, \nabla w \rangle \\
&\quad - \int_M \eta e^{(2m+d)w} |\nabla w|^2 \Delta w \\
&\quad - (2m+d) \int_M \eta e^{(2m+d)w} |\nabla w|^4, \\
\int_M \eta e^{dw} \Delta (e^{2mw} |\nabla w|^2) &= \int_M e^{(2m+d)w} \Delta \eta |\nabla w|^2 \\
&\quad + 2d \int_M e^{(2m+d)w} |\nabla w|^2 \langle \nabla w, \nabla \eta \rangle \\
&\quad + d^2 \int_M \eta e^{(2m+d)w} |\nabla w|^4 \\
&\quad + d \int_M \eta e^{(2m+d)w} |\nabla w|^2 \Delta w.
\end{aligned}$$

Substituting these identities in (2.2) and rearranging terms, we obtain the result.  $\blacksquare$

We apply the previous identity to solutions of equation (1.2).

**Corollary 2.7.** *Let  $(M^2, g)$  be a Riemannian surface, with scalar curvature  $R$ . For any solution  $w \in C^2(M)$  of*

$$-\Delta u = K e^u$$

*with  $K \in C^2(M)$ , any nonnegative  $\psi \in C_c^2(M)$  and any real number  $q \geq 2$  the following identity holds:*

$$\begin{aligned}
&8 \int_M \psi^q e^{\frac{u}{2}} |\mathring{\nabla}^2 e^{-\frac{u}{2}}|^2 + 4 \int_M \psi^q e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 + 2 \int_M \psi^q e^{\frac{u}{2}} \Delta K \\
&= 2 \int_M K e^{\frac{u}{2}} \Delta \psi^q + \int_M e^{-\frac{u}{2}} |\nabla u|^2 \Delta \psi^q.
\end{aligned}$$

*Proof.* We start from Lemma 2.6 with  $w = u$ ,  $\eta = \psi^q$ . Integrating by parts, using the equation on  $u$ , we obtain

$$\begin{aligned}
\int_M \psi^q e^{(2m+d)u} (\Delta u)^2 &= - \int_M \psi^q K e^{(2m+d+1)u} \Delta u \\
&= \int_M K e^{(2m+d+1)u} \langle \nabla \psi^q, \nabla u \rangle + \int_M \psi^q e^{(2m+d+1)u} \langle \nabla K, \nabla u \rangle \\
&\quad + (2m+d+1) \int_M \psi^q K e^{(2m+d+1)u} |\nabla u|^2.
\end{aligned}$$

Assuming  $2m + d + 1 \neq 0$ , we have

$$\begin{aligned}
\int_M \psi^q e^{(2m+d+1)u} \langle \nabla K, \nabla u \rangle \\
&= \frac{1}{2m+d+1} \int_M \psi^q \langle \nabla K, \nabla e^{(2m+d+1)u} \rangle \\
&= -\frac{1}{2m+d+1} \int_M \psi^q e^{(2m+d+1)u} \Delta K - \frac{1}{2m+d+1} \int_M e^{(2m+d+1)u} \langle \nabla K, \nabla \psi^q \rangle \\
&= -\frac{1}{2m+d+1} \int_M \psi^q e^{(2m+d+1)u} \Delta K + \frac{1}{2m+d+1} \int_M K e^{(2m+d+1)u} \Delta \psi^q \\
&\quad + \int_M K e^{(2m+d+1)u} \langle \nabla u, \nabla \psi^q \rangle.
\end{aligned}$$

Hence,

$$\begin{aligned}
&- \left( \frac{(n-1)m^2}{n} + md + \frac{d^2}{2} \right) \int_M \psi^q e^{(2m+d)u} |\nabla u|^4 \\
&\quad + \left( \frac{n+2}{2n} d - \frac{n-1}{n} \right) \int_M \psi^q K e^{(2m+d+1)u} |\nabla u|^2 \\
&\quad + \frac{1}{m^2} \int_M \psi^q e^{du} |\overset{\circ}{\nabla}{}^2 e^{mu}|^2 + \frac{1}{m^2} \int_M \psi^q e^{du} \text{Ric}(\nabla e^{mu}, \nabla e^{mu}) \\
&\quad + \frac{n-1}{n(2m+d+1)} \int_M \psi^q e^{(2m+d+1)u} \Delta K \\
&= (m+d) \int_M e^{(2m+d)u} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle + \frac{n-2}{n} \int_M K e^{(2m+d+1)u} \langle \nabla u, \nabla \psi^q \rangle \\
&\quad + \frac{1}{2} \int_M e^{(2m+d)u} |\nabla u|^2 \Delta \psi^q + \frac{n-1}{n(2m+d+1)} \int_M K e^{(2m+d+1)u} \Delta \psi^q.
\end{aligned}$$

We have  $n = 2$ , hence  $\text{Ric} = \frac{1}{2} R g$ , and we choose  $m = -\frac{1}{2}$ ,  $d = \frac{1}{2}$  so that

$$\begin{aligned}
&4 \int_M \psi^q e^{\frac{u}{2}} |\overset{\circ}{\nabla}{}^2 e^{-\frac{u}{2}}|^2 + 2 \int_M \psi^q e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 + \int_M \psi^q e^{\frac{u}{2}} \Delta K \\
&= \int_M K e^{\frac{u}{2}} \Delta \psi^q + \frac{1}{2} \int_M e^{-\frac{u}{2}} |\nabla u|^2 \Delta \psi^q.
\end{aligned}$$

■

### 2.3. Part III

We collect here some general lemmas that we will need in the proofs of our main theorems. The first is a lower bound for positive superharmonic functions on the Riemannian manifold with nonnegative Ricci curvature. We include a proof for completeness.

**Lemma 2.8.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a complete Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a positive superharmonic function outside a compact set of  $M$ , i.e.,  $\Delta u \leq 0$  on  $M \setminus \Omega$  with  $\Omega$  compact. Then there exist positive constants  $\rho, A > 0$  such that*

$$u(x) \geq \frac{A}{r(x)^{n-2}} \quad \text{on } B_\rho^c.$$

*Proof.* Let  $\rho > 0$  be such that  $\Delta u \leq 0$  on  $B_\rho^c$ . On  $B_\rho^c$  let

$$v(x) := u(x) - \frac{A}{r(x)^\alpha}$$

with  $\alpha = n - 2$  if  $n \geq 3$ ,  $\alpha > 0$  if  $n = 2$  and where  $A := \rho^\alpha \min_{\partial B_\rho} u > 0$ . Then  $v \geq 0$  on  $\partial B_\rho$  and  $\Delta v \leq 0$ , since  $\text{Ric} \geq 0$  implies  $\Delta r \leq \frac{n-1}{r}$  weakly on  $B_\rho^c$  by classical Laplacian comparison. Since  $\liminf_{r(x) \rightarrow \infty} v(x) \geq 0$ , if  $\inf_{B_\rho^c} v < 0$ , then  $v$  attains its negative absolute minimum at a point in  $\bar{B}_\rho^c$ . By the strong maximum principle,  $v$  must be constant and negative on its domain, a contradiction. Thus,  $v \geq 0$  in  $B_\rho^c$ . The proof is complete if  $n \geq 3$ . If  $n = 2$ , for every  $x \in B_\rho^c$ , we have

$$u(x) \geq \frac{\rho^\alpha \min_{\partial B_\rho} u}{r(x)^\alpha}$$

for every  $\alpha > 0$ , and we conclude passing to the limit as  $\alpha$  tends to 0.  $\blacksquare$

The next lemma shows that, under suitable assumptions on the potential  $K$ , subsolutions of equation (1.1) with  $u^{p+1}|K| \in L^1(M)$  automatically have finite energy.

**Lemma 2.9.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a Riemannian manifold, and let  $u \in C^2(M)$  be a nonnegative solution of*

$$-\Delta u \leq Ku^p$$

*with  $1 < p \leq p_c$  and  $K \in C^0(M)$ . If  $K > 0$  almost everywhere outside a compact set of  $M$ ,  $|K|u^{p+1} \in L^1(M)$ , and*

$$\int_{B_{2\rho} \setminus B_\rho} K^{-\frac{2}{p-1}} \leq C\rho^{2\frac{p+1}{p-1}} \quad \text{for every } \rho \text{ large enough,}$$

*then  $|\nabla u| \in L^2(M)$ , i.e.,  $u \in D_{|K|}^{1,2}(M)$ .*

*Proof.* For any  $\rho > 1$ , let  $\psi \in C_c^2(M)$  be such that  $\psi \equiv 1$  in  $B_\rho$ ,  $\psi \equiv 0$  in  $B_{2\rho}^c$ ,  $0 \leq \psi \leq 1$  on  $M$  and  $\psi$  satisfies

$$|\nabla \psi|^2 \leq C\rho^{-2} \quad \text{in } B_{2\rho} \setminus B_\rho.$$

Then, for every  $0 < \varepsilon < 1$ , we obtain

$$\begin{aligned} \int_M K\psi^2 u^{p+1} &\geq - \int_M \psi^2 u \Delta u \geq \int_M \psi^2 |\nabla u|^2 - 2 \int_M \psi u |\nabla u| |\nabla \psi| \\ &\geq (1 - \varepsilon) \int_M \psi^2 |\nabla u|^2 - \frac{1}{\varepsilon} \int_M u^2 |\nabla \psi|^2. \end{aligned}$$

For every large enough  $\rho$ , we have

$$\begin{aligned} \int_M u^2 |\nabla \psi|^2 &\leq \frac{C}{\rho^2} \left( \int_{B_{2\rho}} |K|u^{p+1} \right)^{\frac{2}{p+1}} \left( \int_{B_{2\rho} \setminus B_\rho} K^{-\frac{2}{p-1}} \right)^{\frac{p-1}{p+1}} \\ &\leq C \left( \int_{B_{2\rho}} |K|u^{p+1} \right)^{\frac{2}{p+1}}. \end{aligned}$$

By the previous estimate, we obtain

$$\int_{B_\rho} |\nabla u|^2 \leq C \left( \int_{B_{2\rho}} |K| u^{p+1} \right)^{\frac{2}{p+1}} + C \int_{B_{2\rho}} |K| u^{p+1}.$$

Passing to the limit as  $\rho \rightarrow \infty$ , since  $u^{p+1}|K| \in L^1(M)$ , we obtain the result.  $\blacksquare$

Note that if

$$K \geq \frac{C}{(1+r(x))^\lambda} \quad \text{on } M$$

for some  $\lambda \leq \frac{(n+2)-p(n-2)}{2}$ , then by the Bishop–Gromov volume comparison theorem, we have

$$\int_{B_{2\rho} \setminus B_\rho} K^{-\frac{2}{p-1}} \leq C \rho^{n+\lambda \frac{2}{p-1}} \leq C \rho^{2 \frac{p+1}{p-1}}$$

for every  $\rho > 0$ . In particular, for the critical equation with  $p = p_c = \frac{n+2}{n-2}$ , the condition on  $K$  reads as

$$K \geq C > 0 \quad \text{on } M.$$

Note that these conditions are satisfied if  $K$  is constant and positive on  $M$ .

We finally recall the following characterization of conformal gradient vector fields which can be deduced from general results in [10, 27] and which will be used when dealing with critical equations.

**Lemma 2.10.** *Let  $(M^n, g)$ ,  $n \geq 2$ , be a complete noncompact Riemannian manifold with nonnegative Ricci curvature, and assume there exists a nonconstant function  $f$  on  $M$  such that*

$$\mathring{\nabla}^2 f \equiv 0, \quad \text{Ric}(\nabla f, \nabla f) \equiv 0 \quad \text{on } M.$$

*Then  $(M^n, g)$  is isometric to either*

- (i) *a direct product  $(\mathbb{R} \times N^{n-1}, dr^2 + g_N)$ , where  $(N^{n-1}, g_N)$  is an  $(n-1)$ -dimensional complete Riemannian manifold with nonnegative Ricci curvature and  $f = f(r) = ar + b$  for some  $a, b \in \mathbb{R}$  with  $a \neq 0$ ;*
- (ii)  *$\mathbb{R}^n$  with the Euclidean metric and  $f = f(x) = a|x - x_0|^2 + b$  for some  $x_0 \in \mathbb{R}^n$  and  $a, b \in \mathbb{R}$  with  $a \neq 0$ .*

### 3. Proof of Theorem 1.7

We start considering the case of a Riemannian surface  $(M^2, g)$  with nonnegative scalar curvature  $R$ . As we already observed, by the Bishop–Gromov volume estimate  $(M^2, g)$  must be parabolic (see, for instance, [20]) and therefore every positive superharmonic function is constant. Thus, any solution  $u \geq 0$  of (1.1) is zero. We provide here a self-contained proof. Suppose by contradiction that  $u$  is not identically 0, then by the strong

maximum principle  $u > 0$  on  $M$ . Let  $\psi \in \text{Lip}_c(M)$  be a nonnegative cutoff function,  $\delta \in (-p, -1)$  and  $\varepsilon = |\delta|$ , then we have

$$\begin{aligned} \int_M K\psi^2 u^{p+\delta} &= - \int_M \psi^2 u^\delta \Delta u = 2 \int_M \psi \langle \nabla \psi, \nabla u \rangle u^\delta + \delta \int_M \psi^2 u^{\delta-1} |\nabla u|^2 \\ &\leq \frac{1}{\varepsilon} \int_M |\nabla \psi|^2 u^{\delta+1} + (\varepsilon + \delta) \int_M \psi^2 u^{\delta-1} |\nabla u|^2 \\ &= \frac{1}{\varepsilon} \int_M |\nabla \psi|^2 u^{\delta+1}. \end{aligned} \quad (3.1)$$

Let  $\Phi \in C^1(\mathbb{R})$  be a nonnegative function such that  $0 \leq \Phi \leq 1$ ,  $\Phi' \leq 0$ ,  $\Phi = 1$  on  $(-\infty, 1]$ ,  $\Phi = 0$  on  $[2, -\infty)$ , and  $|\Phi'| \leq C$  for some positive constant  $C$ . For any  $\rho > 1$ , define  $\psi = \Phi(\frac{2 \log r}{\log \rho})$ , then we have  $\psi \equiv 1$  on  $B_{\sqrt{\rho}}$ ,  $\psi \equiv 0$  on  $B_\rho^c$ , and  $|\nabla \psi| \leq \frac{C}{r \log \rho}$  on  $B_\rho \setminus B_{\sqrt{\rho}}$ . With this choice of  $\psi$ , (3.1) and Lemma 2.8 yield

$$\int_M K\psi u^{p+\delta} \leq \frac{C}{(\log \rho)^2} \int_{B_\rho \setminus B_{\sqrt{\rho}}} \frac{1}{r^2}.$$

Let  $V(r) = \text{Vol}(B_r)$  and  $S(r) = \text{meas}(\partial B_r)$  be the area of  $B_r$  and the length of  $\partial B_r$  respectively. Then by the coarea formula  $V' = S$ , and using the coarea formula, an integration by parts and the Bishop–Gromov volume comparison theorem, we obtain

$$\begin{aligned} \int_{B_\rho \setminus B_{\sqrt{\rho}}} \frac{1}{r^2} &= \int_{\sqrt{\rho}}^\rho \frac{1}{r^2} S(r) dr = \left[ \frac{1}{r^2} V(r) \right]_{\sqrt{\rho}}^\rho + 2 \int_{\sqrt{\rho}}^\rho \frac{1}{r^3} V(r) dr \\ &\leq C \left( 1 + \int_{\sqrt{\rho}}^\rho \frac{1}{r} \right) \leq C(1 + \log \rho). \end{aligned}$$

Hence, we have

$$\int_M K\psi u^{p+\delta} \leq \frac{C}{\log \rho},$$

and passing to the limit as  $\rho$  tends to  $\infty$ , we deduce that  $u = 0$  on the set  $\{x \in M : K(x) > 0\}$ . This is a contradiction, since we assumed  $u > 0$  on  $M$ .

Now we consider the case when  $(M^n, g)$ ,  $n \geq 3$ , is a Riemannian manifold with nonnegative Ricci curvature. Let  $u \in C^2(M)$  be a nonnegative solution of (1.1) with  $1 < p < p_c = \frac{n+2}{n-2}$ . By the strong maximum principle,  $u \equiv 0$  or  $u > 0$  on  $M$ . We assume by contradiction that  $u > 0$  on  $M$ , and we apply Corollary 2.3 with  $\psi \in C_c^2(M)$ ,  $0 \leq \psi \leq 1$  on  $M$ ,  $q > 4$  and with  $m, d$  satisfying

$$m = -2 \quad \text{and} \quad d = 2 + \frac{2}{5}p \quad \text{if } n = 3,$$

or

$$2 \frac{n-1}{n+2} p < d < 2 \frac{n-1}{n-2}, \quad (3.2)$$

and

$$\frac{-d - \sqrt{dn(2(n-1) - d(n-2))}}{n-1} < m < \frac{-d + \sqrt{dn(2(n-1) - d(n-2))}}{n-1}$$

when  $n \geq 4$ . Note that the condition on  $d$  is meaningful, since  $p < p_c = \frac{n+2}{n-2}$ , and that with these choices  $m > -2$  when  $n \geq 4$ . Note also that, with this choices, the constants  $\alpha, \beta$  in Corollary 2.3 are positive for every  $n \geq 3$ . Then we have

$$\begin{aligned} & \alpha \int_M \psi^q u^{m-2} |\nabla u|^4 + \beta \int_M \psi^q K u^{m+p-1} |\nabla u|^2 \\ & \leq \frac{n-2}{n} I_1 + \frac{m+d}{2} I_2 + \frac{1}{2} I_3 + \frac{(n-1)}{n(m+p+1)} \int_M \Delta \psi^q K u^{m+p+1}, \end{aligned} \quad (3.3)$$

where

$$\begin{aligned} I_1 &= \int_M K u^{m+p} \langle \nabla u, \nabla \psi^q \rangle, \\ I_2 &= \int_M u^{m-1} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle, \\ I_3 &= \int_M u^m |\nabla u|^2 \Delta \psi^q. \end{aligned}$$

We now bound the integrals  $I_1$ ,  $I_2$  and  $I_3$ . By using Young's inequality, for any  $\varepsilon > 0$ , there exists  $C(\varepsilon) > 0$  such that

$$\begin{aligned} I_1 &\leq \varepsilon \int_M \psi^q K u^{m+p-1} |\nabla u|^2 + C(\varepsilon) \int_M \psi^{q-2} K u^{m+p+1} |\nabla \psi|^2, \\ I_2 &\leq \varepsilon \int_M \psi^q u^{m-2} |\nabla u|^4 + C(\varepsilon) \int_M \psi^{q-4} u^{m+2} |\nabla \psi|^4, \\ I_3 &= q(q-1) \int_M \psi^{q-2} u^m |\nabla u|^2 |\nabla \psi|^2 + q \int_M \psi^{q-1} u^m |\nabla u|^2 \Delta \psi \\ &\leq \varepsilon \int_M \psi^q u^{m-2} |\nabla u|^4 + C(\varepsilon) \int_U \psi^{q-4} u^{m+2} |\nabla \psi|^4 \\ &\quad + C(\varepsilon) \int_M \psi^{q-2} u^{m+2} |\Delta \psi|^2 \\ &\leq \varepsilon \int_M \psi^q u^{m-2} |\nabla u|^4 + C(\varepsilon) \int_U \psi^{q-4} u^{m+2} |\nabla \psi|^4 \\ &\quad + C(\varepsilon) \int_U \psi^{q-4} u^{m+2} |\Delta \psi|^2 \\ &= \varepsilon \int_M \psi^q u^{m-2} |\nabla u|^4 + C(\varepsilon) \int_M \psi^{q-4} u^{m+2} (|\nabla \psi|^4 + |\Delta \psi|^2), \end{aligned} \quad (3.4)$$

since  $0 \leq \psi \leq 1$  everywhere. Substituting the above inequalities for  $\varepsilon$  small enough into (3.3), we obtain

$$\begin{aligned} & \int_M \psi^q u^{m-2} |\nabla u|^4 + \int_M \psi^q K u^{m+p-1} |\nabla u|^2 \\ & \leq C \int_M \psi^{q-2} K u^{m+p+1} (|\nabla \psi|^2 + |\Delta \psi|) \\ & \quad + C \int_M \psi^{q-4} u^{m+2} (|\nabla \psi|^4 + |\Delta \psi|^2) \end{aligned} \quad (3.5)$$

for some positive constant  $C$ . Multiplying equation (1.1) by  $Ku^{m+p}\psi^q$  and integrating by parts, we get

$$\begin{aligned}
& \int_M \psi^q K^2 u^{2p+m} \\
&= I_1 + (m+p) \int_M \psi^q Ku^{m+p-1} |\nabla u|^2 + \int_M \psi^q u^{p+m} \langle \nabla K, \nabla u \rangle \\
&= I_1 + (m+p) \int_M \psi^q Ku^{m+p-1} |\nabla u|^2 + \frac{1}{p+m+1} \int_M \psi^q \langle \nabla K, \nabla u^{p+m+1} \rangle \\
&= I_1 + (m+p) \int_M \psi^q Ku^{m+p-1} |\nabla u|^2 - \frac{1}{p+m+1} \int_M \psi^q \Delta K u^{p+m+1} \\
&\quad - \frac{1}{p+m+1} \int_M \langle \nabla \psi^q, \nabla K \rangle u^{p+m+1} \\
&= 2I_1 + (m+p) \int_M \psi^q Ku^{m+p-1} |\nabla u|^2 - \frac{1}{p+m+1} \int_M \psi^q \Delta K u^{p+m+1} \\
&\quad + \frac{1}{p+m+1} \int_M \Delta \psi^q K u^{p+m+1}.
\end{aligned}$$

By (3.4) and (3.5) and since  $\Delta K \geq 0$ , we have for some  $C > 0$

$$\begin{aligned}
\int_M \psi^q K^2 u^{2p+m} &\leq C \int_M \psi^{q-2} Ku^{m+p+1} (|\nabla \psi|^2 + |\Delta \psi|) \\
&\quad + C \int_M \psi^{q-4} u^{m+2} (|\nabla \psi|^4 + |\Delta \psi|^2).
\end{aligned}$$

We explicitly note that by our choice of  $m$ , we have  $2p+m > p+m+1 > 0$ . By Young's inequality, if  $q \geq \frac{2(2p+m)}{p-1}$ , we have

$$\int_M \psi^q K^2 u^{2p+m} \leq C \int_M \psi^{q-\frac{2(2p+m)}{p-1}} K^{-\frac{m+2}{p-1}} (|\nabla \psi|^2 + |\Delta \psi|)^{\frac{2p+m}{p-1}}.$$

Since  $(M^n, g)$  has nonnegative Ricci curvature, it is possible to construct cutoff functions such that  $\psi \equiv 1$  in  $B_\rho$ ,  $\psi \equiv 0$  in  $B_{2\rho}^c$ ,  $0 \leq \psi \leq 1$  on  $M$  and satisfying (see [29, Lemma 1.5] and [12], for instance)

$$|\Delta \psi| + |\nabla \psi|^2 \leq C\rho^{-2} \quad \text{in } B_{2\rho} \setminus B_\rho.$$

Therefore, we deduce

$$\int_{B_\rho} K^2 u^{2p+m} \leq C\rho^{-\frac{2(2p+m)}{p-1}} \int_{B_{2\rho} \setminus B_\rho} K^{-\frac{m+2}{p-1}}. \quad (3.6)$$

If  $n = 3$ , by our choice of  $m = -2$  and by the Bishop–Gromov volume comparison theorem, we obtain

$$\int_{B_\rho} K^2 u^{2(p-1)} \leq C\rho^{-4} \text{Vol}(B_{2\rho}) \leq \frac{C}{\rho}$$

and thus, passing to the limit as  $\rho$  tends to  $\infty$ , we see that  $u = 0$  on the set  $\{x \in M : K(x) > 0\}$ , a contradiction since we assumed  $u > 0$  on  $M$ .

If  $n \geq 4$ , from (3.6), our assumption on  $K$  and the Bishop–Gromov volume comparison theorem, we have

$$\int_{B_\rho} K^2 u^{2p+m} \leq C \rho^{n-\frac{2(2p+m)}{p-1}+\sigma \frac{m+2}{p-1}} = C \rho^\gamma,$$

and if  $\gamma < 0$ , passing to the limit as  $\rho$  tends to  $\infty$ , we reach a contradiction with our assumption  $u > 0$  on  $M$ , as in the previous case. Now note that we have  $\gamma < 0$  if and only if

$$\sigma < \frac{(-(n-4)p + 2m + n)}{m + 2}$$

since  $m > -2$ . Since this expression is increasing in  $m$ , the optimal choice is

$$\sigma < \frac{(-(n-4)p + 2m + n)}{m + 2} \quad \text{with } m = \frac{-d + \sqrt{dn(2(n-1) - d(n-2))}}{n-1},$$

that is,

$$\sigma < \sigma^*(n, p, d) := \frac{-n^2(p-1) - 4p + 5np - 2d + n + 2\sqrt{dn(2(n-1) - d(n-2))}}{2n - 2 - d + \sqrt{dn(2(n-1) - d(n-2))}}.$$

Since  $d$  satisfies (3.2), choosing  $d < \frac{2(n-1)}{(n-2)}$  arbitrarily close to  $\frac{2(n-1)}{(n-2)}$ , we see that it is sufficient to have

$$\sigma < \sigma^*(n, p) := \sigma^*\left(n, p, \frac{2(n-1)}{(n-2)}\right) = \frac{n(n-2 + 6p - np) - 4 - 8p}{2(n-3)}. \quad (3.7)$$

A simple computation shows that

$$\sigma^*(n, p) \geq \frac{2}{n-3} \Leftrightarrow p \leq \frac{n+2}{n-2},$$

and hence the result follows. We explicitly note that  $\sigma^*(n, p) = \frac{2}{n-3}$  if  $n = 4$  or if  $p = p_c = \frac{n+2}{n-2}$ . The proof of Theorem 1.7 is complete.

#### 4. Proof of Theorem 1.8

Let  $(M^n, g)$ ,  $n \geq 3$ , be a Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative solution of (1.1) with  $p = p_c = \frac{n+2}{n-2}$ . We assume  $u$  is nontrivial, and hence by the strong maximum principle  $u$  is positive on  $M$ . We now use Corollary 2.4 and Lemma 2.5 with  $q = 2$  and  $\varepsilon = \frac{1}{2}$ , and we obtain that for any nonnegative  $\psi \in C_c^2(M)$ ,

$$\begin{aligned} & \frac{(n-2)^2}{8} \int_M \psi^2 u^{\frac{2(n-1)}{n-2}} |\nabla^2 u^{-\frac{2}{n-2}}|^2 + \frac{(n-2)^2}{4} \int_M \psi^2 u^{\frac{2(n-1)}{n-2}} \text{Ric}(\nabla u^{-\frac{2}{n-2}}, u^{-\frac{2}{n-2}}) \\ & + \frac{n-2}{2n} \int_M \psi^2 \Delta K u^{\frac{2(n-1)}{n-2}} \\ & \leq 2 \int_M u^{-\frac{2}{n-2}} |\nabla u|^2 |\nabla \psi|^2 + \frac{n-1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^2 \rangle + \frac{n-2}{2n} \int_M \Delta \psi^2 K u^{\frac{2(n-1)}{n-2}}. \end{aligned}$$

For any  $\rho > 1$ , let  $\psi \in C_c^2(M)$  be such that  $\psi \equiv 1$  in  $B_\rho$ ,  $\psi \equiv 0$  in  $B_{2\rho}^c$ ,  $0 \leq \psi \leq 1$  on  $M$  and  $\psi$  satisfies

$$|\nabla \psi|^2 + |\Delta \psi| \leq C\rho^{-2} \quad \text{in } B_{2\rho} \setminus B_\rho.$$

Then, using Lemma 2.8, for some constant  $C > 0$  we have

$$\int_M u^{-\frac{2}{n-2}} |\nabla u|^2 |\nabla \psi|^2 \leq C \int_{B_\rho^c} |\nabla u|^2.$$

If  $0 \leq K \leq C(1 + r^2)$  outside a compact set of  $M$ , using Lemma 2.8 we have

$$\int_M |\Delta \psi|^2 K u^{\frac{2(n-1)}{n-2}} \leq \frac{C}{\rho^2} \int_{B_{2\rho} \setminus B_\rho} K u^{\frac{2n}{n-2}} u^{-\frac{2}{n-2}} \leq C \int_{B_\rho^c} K u^{\frac{2n}{n-2}}$$

and

$$\begin{aligned} \int_M K u^{\frac{n}{n-2}} |\nabla u| |\nabla \psi|^2 &\leq C \left( \int_{B_{2\rho} \setminus B_\rho} K^2 u^{\frac{2n}{n-2}} |\nabla \psi|^2 \right)^{\frac{1}{2}} \left( \int_{B_\rho^c} |\nabla u|^2 \right)^{\frac{1}{2}} \\ &\leq C \left( \int_{B_{2\rho} \setminus B_\rho} K u^{\frac{2n}{n-2}} \right)^{\frac{1}{2}} \left( \int_{B_\rho^c} |\nabla u|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

On the other hand, if  $|\nabla K(x)| \leq \frac{C}{r(x)} K(x)$  outside a compact set of  $M$ , we have

$$\begin{aligned} \frac{n-1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^2 \rangle &= \frac{n-2}{2n} \int_M K \langle \nabla u^{\frac{2(n-1)}{n-2}}, \nabla \psi^2 \rangle \\ &= -\frac{n-2}{2n} \int_M \Delta \psi^2 K u^{\frac{2(n-1)}{n-2}} \\ &\quad - \frac{n-2}{2n} \int_M u^{\frac{2(n-1)}{n-2}} \langle \nabla K, \nabla \psi^2 \rangle, \end{aligned}$$

and using Lemma 2.8, for  $\rho > 1$  large enough, we obtain

$$\begin{aligned} &\left| \frac{n-1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^2 \rangle + \frac{n-2}{2n} \int_M \Delta \psi^2 K u^{\frac{2(n-1)}{n-2}} \right| \\ &\leq C \int_{B_{2\rho} \setminus B_\rho} |\nabla K| |\nabla \psi| u^{\frac{2n}{n-2}} u^{-\frac{2}{n-2}} \leq C \int_{B_\rho^c} K u^{\frac{2n}{n-2}}. \end{aligned}$$

In either case, since  $\Delta K \geq 0$  on  $M$ , for every  $\rho > 1$  large enough we have

$$\begin{aligned} &\int_{B_\rho} u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 + \int_{B_\rho} u^{\frac{2(n-1)}{n-2}} \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) \\ &\leq C \left( \int_{B_\rho^c} |\nabla u|^2 + \int_{B_\rho^c} K u^{\frac{2n}{n-2}} \right) \end{aligned}$$

for some constant  $C > 0$ . Passing to the limit as  $\rho$  tends to  $\infty$ , since  $\text{Ric} \geq 0$ , we conclude that

$$\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}} \equiv 0, \quad \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) \equiv 0 \quad \text{on } M.$$

From Lemma 2.10, we deduce that  $(M^n, g)$  is either

- (i) a direct product  $(\mathbb{R} \times N^{n-1}, dr^2 + g_N)$ , where  $(N^{n-1}, g_N)$  is an  $(n-1)$ -dimensional complete Riemannian manifold with nonnegative Ricci curvature and  $u^{-\frac{2}{n-2}} = f = f(r) = ar + b$  for some  $a, b \in \mathbb{R}$  with  $a \neq 0$ , or
- (ii)  $\mathbb{R}^n$  with the Euclidean metric and  $u^{-\frac{2}{n-2}} = f = f(x) = a|x - x_0|^2 + b$  for some  $a, b \in \mathbb{R}$  with  $a \neq 0$ .

Since

$$\nabla f = -\frac{2}{n-2}u^{-\frac{n}{n-2}}\nabla u, \quad \Delta f = -\frac{2}{n-2}u^{-\frac{n}{n-2}}\Delta u + \frac{2n}{(n-2)^2}u^{-\frac{2(n-1)}{n-2}}|\nabla u|^2,$$

a simple computation shows that in the first case,

$$0 = \Delta f = \frac{2}{n-2}Ku^{\frac{2}{n-2}} + \frac{2n}{(n-2)^2}u^{-\frac{2(n-1)}{n-2}}|\nabla u|^2 \geq 0 \quad \text{on } M,$$

which is in contradiction with our assumption  $u > 0$  on  $M$ . Then  $(M^n, g)$  is  $\mathbb{R}^n$  with the Euclidean metric and  $u^{-\frac{2}{n-2}} = f = f(x) = a|x - x_0|^2 + b$  for some  $x_0 \in \mathbb{R}^n, a, b \in \mathbb{R}$  with  $a \neq 0$ . Since  $u, f$  are positive functions, we must have  $a, b > 0$  and thus

$$u(x) = (a|x - x_0|^2 + b)^{-\frac{n-2}{2}}.$$

Inserting this expression into the equation, we find that we must have  $K \equiv n(n-2)ab \in (0, \infty)$ , and thus we conclude.

## 5. Proof of Theorem 1.9

Let  $(M^n, g)$ ,  $n \geq 3$ , be a Riemannian manifold with nonnegative Ricci curvature, and let  $u \in C^2(M)$  be a nonnegative solution of (1.1) with  $p = p_c = \frac{n+2}{n-2}$ ,  $K \in C^2(M)$ ,  $0 \neq K \geq 0$  and  $\Delta K \geq 0$ . We assume  $u$  is nontrivial, and hence by the strong maximum principle,  $u$  is positive on  $M$ . We now use Corollary 2.4 and Lemma 2.5 with  $\varepsilon = \frac{1}{q}$  and obtain that for any nonnegative  $\psi \in C_c^2(M)$ ,

$$\begin{aligned} & \frac{(n-2)^2}{8} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 \\ & + \frac{(n-2)^2}{4} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) + \frac{n-2}{2n} \int_M \psi^q \Delta K u^{\frac{2(n-1)}{(n-2)}} \\ & \leq \frac{q^2}{2} \int_M \psi^{q-2} u^{-\frac{2}{n-2}} |\nabla u|^2 |\nabla \psi|^2 + \frac{n-1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla \psi^q, \nabla u \rangle \\ & + \frac{n-2}{2n} \int_M \Delta \psi^q K u^{\frac{2(n-1)}{(n-2)}}. \end{aligned}$$

We have

$$\begin{aligned} & \frac{n-1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla \psi^q, \nabla u \rangle = \frac{n-2}{2n} \int_M K \langle \nabla \psi^q, \nabla u^{\frac{2(n-1)}{(n-2)}} \rangle \\ & = -\frac{n-2}{2n} \int_M K \Delta \psi^q u^{\frac{2(n-1)}{(n-2)}} - \frac{n-2}{2n} \int_M \langle \nabla \psi^q, \nabla K \rangle u^{\frac{2(n-1)}{(n-2)}}. \end{aligned}$$

Hence, by our assumptions on  $K$  and since  $0 \leq \psi \leq 1$ ,

$$\begin{aligned} & \left| \frac{n-1}{n} \int_M K u^{\frac{n}{n-2}} \langle \nabla u, \nabla \psi^q \rangle + \frac{n-2}{2n} \int_M \Delta \psi^q K u^{\frac{2(n-1)}{n-2}} \right| \\ & \leq C \int_{B_{2\rho} \setminus B_\rho} |\nabla K| \psi^{q-1} |\nabla \psi| u^{\frac{2(n-1)}{n-2}} \leq \frac{\bar{C}}{\rho^2} \int_{B_{2\rho} \setminus B_\rho} \psi^{q-2} K u^{\frac{2(n-1)}{n-2}}, \end{aligned}$$

and thus

$$\begin{aligned} & \frac{(n-2)^2}{8} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 + \frac{(n-2)^2}{4} \int_M \psi^q u^{\frac{2(n-1)}{n-2}} \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) \\ & \leq 8 \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 u^{\frac{\gamma}{n-2}} |\nabla \psi|^2 + \frac{\bar{C}}{\rho^2} \int_{B_{2\rho} \setminus B_\rho} \psi^{q-2} K u^{\frac{2(n-1)}{n-2}} \end{aligned}$$

for any  $\gamma \geq 0$ . Multiplying the equation for  $u$  by  $\psi^{q-2} u^{\frac{n-4-\gamma}{n-2}}$  and integrating by parts, we obtain

$$\begin{aligned} & \frac{n-4-\gamma}{n-2} \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 + (q-2) \int_M \psi^{q-3} u^{\frac{n-4-\gamma}{n-2}} \langle \nabla u, \nabla \psi \rangle \\ & = \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}}. \end{aligned} \tag{5.1}$$

Hence, if  $n = 3$ , we choose  $\gamma = 0, q = 4$  thus obtaining

$$\begin{aligned} \int_M \psi^2 u^{-2} |\nabla u|^2 & = 2 \int_M \psi u^{-1} \langle \nabla u, \nabla \psi \rangle - \int_M \psi^2 K u^4 \\ & \leq \frac{1}{2} \int_M \psi^2 u^{-2} |\nabla u|^2 + 2 \int_M |\nabla \psi|^2 - \int_M \psi^2 K u^4. \end{aligned}$$

For any  $\rho > 1$ , let  $\psi \in C_c^2(M)$  be such that  $\psi \equiv 1$  in  $B_\rho$ ,  $\psi \equiv 0$  in  $B_{2\rho}^c$ ,  $0 \leq \psi \leq 1$  on  $M$  and  $\psi$  satisfies

$$|\nabla \psi|^2 \leq C \rho^{-2} \quad \text{in } B_{2\rho} \setminus B_\rho.$$

By the Bishop–Gromov volume estimate, we conclude

$$\int_{B_{2\rho}} \psi^2 u^{-2} |\nabla u|^2 \leq C \rho - 2 \int_M \psi^2 K u^4 \leq C_1 \left( C \rho - 2 \int_M \psi^2 K u^4 \right)$$

for every  $C_1 \geq 1$ , and thus

$$\begin{aligned} & \int_M \psi^4 u^4 |\overset{\circ}{\nabla}^2 u^{-2}|^2 + \int_M \psi^4 u^4 \text{Ric}(\nabla u^{-2}, \nabla u^{-2}) \\ & \leq C \int_M \psi^2 u^{-2} |\nabla u|^2 |\nabla \psi|^2 + \frac{\bar{C}}{\rho^2} \int_{B_{2\rho} \setminus B_\rho} \psi^2 K u^4 \\ & \leq \frac{C C_1}{\rho^2} \left( C \rho - 2 \int_M \psi^2 K u^4 \right) + \frac{\bar{C}}{\rho^2} \int_{B_{2\rho} \setminus B_\rho} \psi^2 K u^4 \leq \frac{C}{\rho} \end{aligned}$$

if  $C_1 \geq 1$  is chosen large enough. Letting  $\rho \rightarrow \infty$ , we deduce that

$$\tilde{\nabla}^2 u^{-2} \equiv 0, \quad \text{Ric}(\nabla u^{-2}, \nabla u^{-2}) \equiv 0 \quad \text{on } M,$$

and we conclude as in the proof of Theorem 1.8. If  $n \geq 4$ , for every  $n-4 < \gamma < 2n-6$  and every  $\varepsilon > 0$  by (5.1), we have

$$\begin{aligned} & \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 \\ & \leq (q-2)C \int_M \psi^{q-3} u^{\frac{n-4-\gamma}{n-2}} \langle \nabla u, \nabla \psi \rangle - C \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} \\ & \leq C \varepsilon \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 + \frac{C}{\varepsilon} \int_M \psi^{q-4} u^{\frac{2n-6-\gamma}{n-2}} |\nabla \psi|^2 \\ & \quad - C \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}}, \end{aligned}$$

and thus for some  $C > 0$  depending on  $\gamma$ , we have

$$\begin{aligned} & \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 \\ & \leq C \int_M \psi^{q-4} u^{\frac{2n-6-\gamma}{n-2}} |\nabla \psi|^2 - C \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}}. \end{aligned} \quad (5.2)$$

Now for every  $\varepsilon > 0$ , since  $\gamma < 2n-6$ , we have

$$\begin{aligned} & \int_M \psi^{q-4} u^{\frac{2n-6-\gamma}{n-2}} |\nabla \psi|^2 \\ & \leq \frac{\varepsilon}{2} \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} + \frac{1}{2\varepsilon} \int_M \psi^{q-1-n+\frac{\gamma}{2}} K^{-\frac{2n-6-\gamma}{4}} |\nabla \psi|^{\frac{2n-2-\gamma}{2}}. \end{aligned}$$

Choosing  $\varepsilon > 0$  suitably small and substituting in (5.2), we find

$$\begin{aligned} & \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 \\ & \leq C \int_M \psi^{q-1-n+\frac{\gamma}{2}} |\nabla \psi|^{\frac{2n-2-\gamma}{2}} K^{-\frac{2n-6-\gamma}{4}} - \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} \\ & \leq C_1 \left( C \int_M \psi^{q-1-n+\frac{\gamma}{2}} |\nabla \psi|^{\frac{2n-2-\gamma}{2}} K^{-\frac{2n-6-\gamma}{4}} - \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} \right) \end{aligned}$$

for large enough  $C > 0$  and every  $C_1 \geq 1$ . For any  $\rho > 1$ , we choose again  $\psi \in C_c^2(M)$  such that  $\psi \equiv 1$  in  $B_\rho$ ,  $\psi \equiv 0$  in  $B_{2\rho}^c$ ,  $0 \leq \psi \leq 1$  on  $M$  and such that  $\psi$  satisfies

$$|\nabla \psi|^2 \leq C \rho^{-2} \quad \text{in } B_{2\rho} \setminus B_\rho.$$

Thus, by our assumptions on  $u$ ,  $K$  and using the Bishop–Gromov volume comparison theorem, choosing

$$q \geq 1 + n - \frac{\gamma}{2} \quad \text{and} \quad C_1 \geq 1$$

large enough, we obtain

$$\begin{aligned}
& \int_M \psi^q u^{\frac{2(n-1)}{n-2}} |\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}}|^2 + \int_M \psi^q u^{\frac{2(n-1)}{n-2}} \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) \\
& \leq C(\gamma) \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 u^{\frac{\gamma}{n-2}} |\nabla \psi|^2 + \frac{\bar{C}}{\rho^2} \int_{B_{2\rho} \setminus B_\rho} \psi^{q-2} K u^{\frac{2(n-1)}{n-2}} \\
& \leq \frac{1}{\rho^2} (\sup_{B_{2\rho} \setminus B_\rho} u)^{\frac{\gamma}{n-2}} \left( C \int_M \psi^{q-2} u^{-\frac{2+\gamma}{n-2}} |\nabla u|^2 + \bar{C} \int_{B_{2\rho} \setminus B_\rho} \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} \right) \\
& \leq \frac{1}{\rho^2} (\sup_{B_{2\rho} \setminus B_\rho} u)^{\frac{\gamma}{n-2}} \left( C C_1 \int_M \psi^{q-1-n+\frac{\gamma}{2}} |\nabla \psi|^{\frac{2n-2-\gamma}{2}} K^{-\frac{2n-6-\gamma}{4}} \right. \\
& \quad \left. - C_1 \int_M \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} + \bar{C} \int_{B_{2\rho} \setminus B_\rho} \psi^{q-2} K u^{\frac{2n-2-\gamma}{n-2}} \right) \\
& \leq \frac{C}{\rho^2} (\sup_{B_{2\rho} \setminus B_\rho} u)^{\frac{\gamma}{n-2}} \int_M |\nabla \psi|^{\frac{2n-2-\gamma}{2}} K^{-\frac{2n-6-\gamma}{4}} \\
& \leq C \rho^{\gamma(\frac{\alpha}{n-2} + \frac{1}{2} - \frac{\sigma}{4}) + \frac{n-3}{2}\sigma - 1}.
\end{aligned}$$

Now if

$$\lambda = \lambda(\gamma) := \gamma \left( \frac{\alpha}{n-2} + \frac{1}{2} - \frac{\sigma}{4} \right) + \frac{n-3}{2}\sigma - 1 < 0$$

for some  $\gamma \in (n-4, 2n-6)$ , passing to the limit as  $\rho$  tends to  $\infty$  yields

$$\overset{\circ}{\nabla}^2 u^{-\frac{2}{n-2}} \equiv 0, \quad \text{Ric}(\nabla u^{-\frac{2}{n-2}}, \nabla u^{-\frac{2}{n-2}}) \equiv 0 \quad \text{on } M,$$

and again we can conclude as in the proof of Theorem 1.8. Since  $\lambda(\gamma)$  is monotone in  $\gamma$ , we have that  $\lambda(\gamma) < 0$  for some  $\gamma \in (n-4, 2n-6)$  if and only if  $\lambda(n-4) < 0$  or  $\lambda(2n-6) < 0$ . These conditions are equivalent to, respectively,

$$\alpha < -\frac{(n-2)^2\sigma + 2(n-2)(n-6)}{4(n-4)} \quad \text{or} \quad \alpha < -\frac{(n-2)(n-4)}{2(n-3)}.$$

Then by our assumptions on  $\alpha, \sigma$  there exists  $\gamma \in (n-4, 2n-6)$  such that  $\lambda(\gamma) < 0$ , and the proof is complete.

## 6. Proof of Theorem 1.11

By Corollary 2.7 for every nonnegative  $\psi \in C_c^2(M)$  and every  $q \geq 2$ , since  $\Delta K \geq 0$ , we have

$$\begin{aligned}
& 8 \int_M \psi^q e^{\frac{u}{2}} |\overset{\circ}{\nabla}^2 e^{-\frac{u}{2}}|^2 + 4 \int_M \psi^q e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 \\
& \leq 2 \int_M K e^{\frac{u}{2}} \Delta \psi^q + \int_M e^{-\frac{u}{2}} |\nabla u|^2 \Delta \psi^q.
\end{aligned} \tag{6.1}$$

Integrating by parts and using Young's inequality, for every  $\varepsilon > 0$ , we have

$$\begin{aligned}
\frac{1}{4} \int_M e^{-\frac{u}{2}} |\nabla u|^2 \Delta \psi^q &= \int_M e^{\frac{u}{2}} |\nabla e^{-\frac{u}{2}}|^2 \Delta \psi^q \\
&= -\frac{1}{2} \int_M e^{\frac{u}{2}} |\nabla e^{-\frac{u}{2}}|^2 \langle \nabla u, \nabla \psi^q \rangle - \int_M e^{\frac{u}{2}} \langle \nabla |\nabla e^{-\frac{u}{2}}|^2, \nabla \psi^q \rangle \\
&= -\frac{1}{8} \int_M e^{-\frac{u}{2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle - 2 \int_M e^{\frac{u}{2}} \nabla^2 e^{-\frac{u}{2}} \langle \nabla e^{-\frac{u}{2}}, \nabla \psi^q \rangle \\
&= -\frac{1}{8} \int_M e^{-\frac{u}{2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle - 2 \int_M e^{\frac{u}{2}} \overset{\circ}{\nabla}{}^2 e^{-\frac{u}{2}} \langle \nabla e^{-\frac{u}{2}}, \nabla \psi^q \rangle \\
&\quad - \int_M e^{\frac{u}{2}} \Delta e^{-\frac{u}{2}} \langle \nabla e^{-\frac{u}{2}}, \nabla \psi^q \rangle \\
&\leq -\frac{1}{8} \int_M e^{-\frac{u}{2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle + \varepsilon \int_M \psi^q e^{\frac{u}{2}} |\overset{\circ}{\nabla}{}^2 e^{-\frac{u}{2}}|^2 \\
&\quad + \frac{C}{\varepsilon} \int_M \psi^{q-2} e^{-\frac{u}{2}} |\nabla u|^2 |\nabla \psi|^2 \\
&\quad + \frac{1}{8} \int_M e^{-\frac{u}{2}} |\nabla u|^2 \langle \nabla u, \nabla \psi^q \rangle + \frac{1}{4} \int_M K e^{\frac{u}{2}} \langle \nabla u, \nabla \psi^q \rangle.
\end{aligned}$$

Then, from (6.1) choosing  $\varepsilon = 1, q = 4$ , we obtain

$$\begin{aligned}
&4 \int_M \psi^4 e^{\frac{u}{2}} |\overset{\circ}{\nabla}{}^2 e^{-\frac{u}{2}}|^2 + 4 \int_M \psi^4 e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 \\
&\leq 2 \int_M K e^{\frac{u}{2}} \Delta \psi^4 + C \int_M \psi^2 e^{-\frac{u}{2}} |\nabla u|^2 |\nabla \psi|^2 + \int_M K e^{\frac{u}{2}} \langle \nabla u, \nabla \psi^4 \rangle. \quad (6.2)
\end{aligned}$$

*Case 1:* Suppose that

$$0 \leq K(x) \leq C r(x)^2$$

outside a compact set of  $M$ . Since  $M$  has nonnegative Ricci curvature, by using [2], one can see that there exist  $C > 0$  and  $\theta \in (0, \frac{1}{2})$  such that, for any  $\rho$  large enough, there exists a cut-off function  $\psi \in C^2(M)$  such that  $\psi \equiv 1$  on  $B_{\rho^\theta}$ ,  $\psi \equiv 0$  on  $B_\rho^c$  and

$$|\nabla \psi| \leq \frac{C}{r \log \rho}, \quad |\Delta \psi| \leq \frac{C}{r^2 \log \rho} \quad \text{on } B_\rho \setminus B_{\rho^\theta}.$$

Note also that, up to increasing the constant  $C > 0$ , we have

$$|\nabla \psi| \leq \frac{C}{(r+1) \log \rho}, \quad \text{on } B_\rho \setminus B_{\rho^\theta}.$$

With this choice of  $\psi$ , by our assumption on  $u$ , for  $\rho > 1$  large enough

$$\int_M \psi^2 e^{-\frac{u}{2}} |\nabla u|^2 |\nabla \psi|^2 \leq \frac{C}{(\log \rho)^2} \int_M \frac{\psi^2}{(r+1)^2} e^{-\frac{u}{2}} |\nabla u|^2 \quad (6.3)$$

and

$$\begin{aligned}
\int_M \frac{\psi^2}{(r+1)^2} e^{-\frac{u}{2}} |\nabla u|^2 &= -2 \int_M \frac{\psi^2}{(r+1)^2} \langle \nabla e^{-\frac{u}{2}}, \nabla u \rangle \\
&= 2 \int_M \frac{1}{(r+1)^2} e^{-\frac{u}{2}} \langle \nabla \psi^2, \nabla u \rangle + 2 \int_M \frac{\psi^2}{(r+1)^2} e^{-\frac{u}{2}} \Delta u \\
&\quad - 4 \int_M \frac{\psi^2}{(r+1)^3} e^{-\frac{u}{2}} \langle \nabla r, \nabla u \rangle \\
&\leq \frac{1}{4} \int_M \frac{\psi^2}{(r+1)^2} e^{-\frac{u}{2}} |\nabla u|^2 + C \int_M \frac{1}{(r+1)^2} e^{-\frac{u}{2}} |\nabla \psi|^2 \\
&\quad - 2 \int_M \frac{\psi^2}{(r+1)^2} K e^{\frac{u}{2}} + \frac{1}{4} \int_M \frac{\psi^2}{(r+1)^2} e^{-\frac{u}{2}} |\nabla u|^2 \\
&\quad + C \int_M \frac{\psi^2}{(r+1)^4} e^{-\frac{u}{2}}.
\end{aligned}$$

Thus, by our assumptions on  $u$ , the choice of  $\psi$  and the Bishop–Gromov volume comparison, arguing as in the proof of Theorem 1.7 when  $n = 2$ , we have

$$\begin{aligned}
\int_M \frac{\psi^2}{(r+1)^2} e^{-\frac{u}{2}} |\nabla u|^2 &\leq C \int_M \frac{1}{(r+1)^2} e^{-\frac{u}{2}} |\nabla \psi|^2 + C \int_M \frac{\psi^2}{(r+1)^4} e^{-\frac{u}{2}} \\
&\leq \frac{C}{(\log \rho)^2} \int_{B_\rho \setminus B_{\rho^\theta}} \frac{(\log r)^\gamma}{r^2} + C \int_{B_\rho} \frac{r^2 (\log r)^\gamma}{(r+1)^4} \\
&\leq C (\log \rho)^{1+\gamma}.
\end{aligned}$$

From (6.3) for every  $\rho > 1$  large enough, we obtain

$$\int_M \psi^2 e^{-\frac{u}{2}} |\nabla u|^2 |\nabla \psi|^2 \leq \frac{C}{(\log \rho)^{1-\gamma}}. \quad (6.4)$$

Moreover,

$$\int_M K e^{\frac{u}{2}} \Delta \psi^4 \leq C \int_{B_\rho \setminus B_{\rho^\theta}} K e^u \frac{e^{-\frac{u}{2}}}{r^2 \log \rho} \leq C \int_{(B_{\rho^\theta})^c} K e^u.$$

Finally,

$$\begin{aligned}
\int_M K e^{\frac{u}{2}} \langle \nabla u, \nabla \psi^4 \rangle &\leq \frac{1}{2} \int_{(B_{\rho^\theta})^c} K e^u + \frac{1}{2} \int_{B_\rho \setminus B_{\rho^\theta}} K |\nabla u|^2 |\nabla \psi|^2 \psi^2 \\
&\leq \frac{1}{2} \int_{(B_{\rho^\theta})^c} K e^u + \frac{C}{(\log \rho)^2} \int_{B_\rho} |\nabla u|^2 \psi^2.
\end{aligned}$$

We multiply the equation satisfied by  $u$ , i.e., (1.2), by  $\psi^2 u$  and integrate by parts. Since  $K \geq 0$ ,  $0 \leq \psi \leq 1$  on  $M$ ,  $K e^u \in L^1(M)$ , and

$$-4 \log r - \gamma \log \log r \leq u \leq C \log r$$

for  $r$  large enough, for every large enough  $\rho > 1$ , we have

$$\begin{aligned} \int_{B_\rho} |\nabla u|^2 \psi^2 &= \int_M \psi^2 K u e^u - 2 \int_M \psi u \langle \nabla u, \nabla \psi \rangle \\ &\leq \int_{B_\rho} \psi^2 K u e^u + \frac{1}{2} \int_{B_\rho} |\nabla u|^2 \psi^2 + 2 \int_{B_\rho \setminus B_{\rho^\theta}} u^2 |\nabla \psi|^2 \\ &\leq \frac{1}{2} \int_{B_\rho} |\nabla u|^2 \psi^2 + C \log \rho + C, \end{aligned}$$

where we also used the Bishop–Gromov volume comparison theorem. Then

$$\int_{B_\rho} |\nabla u|^2 \psi^2 \leq C \log \rho + C,$$

and therefore

$$\int_M K e^{\frac{u}{2}} \langle \nabla u, \nabla \psi^4 \rangle \leq \frac{1}{2} \int_{(B_{\rho^\theta})^c} K e^u + \frac{C}{\log \rho}.$$

Inserting these estimates into (6.2), we obtain for every  $\rho > 1$  large enough

$$\begin{aligned} 4 \int_M \psi^4 e^{\frac{u}{2}} |\overset{\circ}{\nabla}^2 e^{-\frac{u}{2}}|^2 + 4 \int_M \psi^4 e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 \\ \leq C \int_{(B_{\rho^\theta})^c} K e^u + \frac{C}{(\log \rho)} + \frac{C}{(\log \rho)^{1-\gamma}}. \end{aligned}$$

Passing to the limit as  $\rho$  tends to  $\infty$ , we obtain

$$\overset{\circ}{\nabla}^2 e^{-\frac{u}{2}} \equiv 0, \quad R |\nabla e^{-\frac{u}{2}}|^2 \equiv 0 \quad \text{on } M,$$

and again we can conclude as in the proof of Theorem 1.8.

*Case 2:* If outside a compact set  $|\nabla K(x)| \leq \frac{C}{r(x)} K(x)$ , integrating by parts we have

$$\begin{aligned} \int_M K e^{\frac{u}{2}} \langle \nabla u, \nabla \psi^4 \rangle &= 2 \int_M K \langle \nabla e^{\frac{u}{2}}, \nabla \psi^4 \rangle \\ &= -2 \int_M e^{\frac{u}{2}} \langle \nabla K, \nabla \psi^4 \rangle - 2 \int_M K e^{\frac{u}{2}} \Delta \psi^4 \\ &\leq C \int_M \frac{K}{r} e^{\frac{u}{2}} |\nabla \psi| - 2 \int_M K e^{\frac{u}{2}} \Delta \psi^4. \end{aligned}$$

Substituting in (6.2), we obtain

$$\begin{aligned} 4 \int_M \psi^4 e^{\frac{u}{2}} |\overset{\circ}{\nabla}^2 e^{-\frac{u}{2}}|^2 + 4 \int_M \psi^4 e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 \\ \leq C \int_M \psi^2 e^{-\frac{u}{2}} |\nabla u|^2 |\nabla \psi|^2 + C \int_M \frac{K}{r} e^{\frac{u}{2}} |\nabla \psi|. \end{aligned} \tag{6.5}$$

Let  $\Phi \in C^1(\mathbb{R})$  be a nonnegative function such that  $0 \leq \Phi \leq 1$ ,  $\Phi' \leq 0$ ,  $\Phi = 1$  on  $(-\infty, 1]$ ,  $\Phi = 0$  on  $[2, -\infty)$ , and  $|\Phi'| \leq C$  for some positive constant  $C$ . For any  $\rho > 1$ , define  $\psi =$

$\Phi(\frac{2\log r}{\log \rho})$ , then we have  $\psi \equiv 1$  on  $B_{\sqrt{\rho}}$ ,  $\psi \equiv 0$  on  $B_\rho^c$ , and  $|\nabla \psi| \leq \frac{C}{r \log \rho}$  on  $B_\rho \setminus B_{\sqrt{\rho}}$ . With this choice of  $\psi$ , by our assumption on  $u$ , we get

$$\int_M \frac{K}{r} e^{\frac{u}{2}} |\nabla \psi| \leq \frac{C}{\log \rho} \int_{B_\rho \setminus B_{\sqrt{\rho}}} K e^u \left( \frac{e^{-\frac{u}{2}}}{r^2} \right) \leq C \int_{B_{\sqrt{\rho}}^c} K e^u.$$

On the other hand, arguing as in case 1, we obtain inequality (6.4). Finally, using (6.5), we have

$$4 \int_M \psi^4 e^{\frac{u}{2}} |\overset{\circ}{\nabla}^2 e^{-\frac{u}{2}}|^2 + 4 \int_M \psi^4 e^{\frac{u}{2}} R |\nabla e^{-\frac{u}{2}}|^2 \leq \frac{C}{(\log \rho)^{1-\gamma}} + C \int_{B_{\sqrt{\rho}}^c} K e^u.$$

Passing to the limit as  $\rho$  tends to  $\infty$ , we obtain

$$\overset{\circ}{\nabla}^2 e^{-\frac{u}{2}} \equiv 0, \quad R |\nabla e^{-\frac{u}{2}}|^2 \equiv 0 \quad \text{on } M,$$

and again we can conclude as in the proof of Theorem 1.8.

*Acknowledgments.* The first author is a member of the *GNSAGA, Gruppo Nazionale per le Strutture Algebriche, Geometriche e le loro Applicazioni* of Indam. The second author is a member of *GNAMPA, Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni* of Indam.

## References

- [1] Aubin, T.: [Problèmes isopérimétriques et espaces de Sobolev](#). J. Differential Geometry **11**, 573–598 (1976) Zbl 0371.46011 MR 448404
- [2] Bianchi, D., Setti, A. G.: [Laplacian cut-offs, porous and fast diffusion on manifolds and other applications](#). Calc. Var. Partial Differential Equations **57**, article no. 4 (2018) Zbl 1502.53057 MR 3735744
- [3] Bidaut-Véron, M.-F., Raoux, T.: [Asymptotics of solutions of some nonlinear elliptic systems](#). Comm. Partial Differential Equations **21**, 1035–1086 (1996) Zbl 0932.35075 MR 1399190
- [4] Brendle, S.: [Sobolev inequalities in manifolds with nonnegative curvature](#). Comm. Pure Appl. Math. **76**, 2192–2218 (2023) Zbl 1527.53030 MR 4612577
- [5] Brezis, H., Merle, F.: [Uniform estimates and blow-up behavior for solutions of  \$-\Delta u = V\(x\)e^u\$  in two dimensions](#). Comm. Partial Differential Equations **16**, 1223–1253 (1991) Zbl 0746.35006 MR 1132783
- [6] Caffarelli, L. A., Gidas, B., Spruck, J.: [Asymptotic symmetry and local behavior of semilinear elliptic equations with critical Sobolev growth](#). Comm. Pure Appl. Math. **42**, 271–297 (1989) Zbl 0702.35085 MR 982351
- [7] Carron, G.: Inégalités isopérimétriques de Faber–Krahn et conséquences. In: Actes de la Table Ronde de Géométrie Différentielle (Luminy, 1992), Sémin. Congr. 1, Société Mathématique de France, Paris, 205–232 (1996) Zbl 0884.58088 MR 1427759
- [8] Castorina, D., Catino, G., Mantegazza, C.: [A triviality result for semilinear parabolic equations](#). Math. Eng. **4**, article no. 002 (2022) Zbl 1496.35234 MR 4231144
- [9] Catino, G., Li, Y., Monticelli, D. D., Roncoroni, A.: A Liouville theorem in the Heisenberg group. arXiv:2310.10469 (2023)
- [10] Catino, G., Mantegazza, C., Mazzieri, L.: [On the global structure of conformal gradient solitons with nonnegative Ricci tensor](#). Commun. Contemp. Math. **14**, article no. 1250045 (2012) Zbl 1253.53044 MR 2989649

- [11] Catino, G., Monticelli, D. D., Roncoroni, A.: [On the critical  \$p\$ -Laplace equation](#). Adv. Math. **433**, article no. 109331 (2023) Zbl 07759066 MR 4649289
- [12] Cheeger, J., Colding, T. H.: [Lower bounds on Ricci curvature and the almost rigidity of warped products](#). Ann. of Math. (2) **144**, 189–237 (1996) Zbl 0865.53037 MR 1405949
- [13] Chen, W. X., Li, C.: [Classification of solutions of some nonlinear elliptic equations](#). Duke Math. J. **63**, 615–622 (1991) Zbl 0768.35025 MR 1121147
- [14] Chen, W. X., Li, C.: [Qualitative properties of solutions to some nonlinear elliptic equations in  \$\mathbb{R}^2\$](#) . Duke Math. J. **71**, 427–439 (1993) Zbl 0923.35055 MR 1233443
- [15] Ciraolo, G., Figalli, A., Roncoroni, A.: [Symmetry results for critical anisotropic  \$p\$ -Laplacian equations in convex cones](#). Geom. Funct. Anal. **30**, 770–803 (2020) Zbl 1453.35104 MR 4135671
- [16] Flynn, J., Vétois, J.: Liouville-type results for the CR Yamabe equation in the Heisenberg group. arXiv:2310.14048 (2023)
- [17] Fogagnolo, M., Malchiodi, A., Mazzieri, L.: [A note on the critical Laplace equation and Ricci curvature](#). J. Geom. Anal. **33**, article no. 178 (2023) Zbl 1522.35163 MR 4567579
- [18] Gidas, B., Ni, W. M., Nirenberg, L.: Symmetry of positive solutions of nonlinear elliptic equations in  $\mathbb{R}^n$ . In: Mathematical analysis and applications, Part A, Adv. Math. Suppl. Stud. 7, Academic Press, New York, 369–402 (1981) Zbl 0469.35052 MR 634248
- [19] Gidas, B., Spruck, J.: [Global and local behavior of positive solutions of nonlinear elliptic equations](#). Comm. Pure Appl. Math. **34**, 525–598 (1981) Zbl 0465.35003 MR 615628
- [20] Grigor'yan, A.: [Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds](#). Bull. Amer. Math. Soc. (N.S.) **36**, 135–249 (1999) Zbl 0927.58019 MR 1659871
- [21] Kim, S.: [Scalar curvature on noncompact complete Riemannian manifolds](#). Nonlinear Anal. **26**, 1985–1993 (1996) Zbl 0858.53029 MR 1386128
- [22] Li, Y. Y., Zhang, L.: [Liouville-type theorems and Harnack-type inequalities for semilinear elliptic equations](#). J. Anal. Math. **90**, 27–87 (2003) Zbl 1173.35477 MR 2001065
- [23] Muratori, M., Soave, N.: [Some rigidity results for Sobolev inequalities and related PDEs on Cartan–Hadamard manifolds](#). Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **24**, 751–792 (2023) Zbl 1522.58012 MR 4630043
- [24] Obata, M.: [The conjectures on conformal transformations of Riemannian manifolds](#). J. Differential Geometry **6**, 247–258 (1971) Zbl 0236.53042 MR 303464
- [25] Ou, Q.: On the classification of entire solutions to the critical  $p$ -Laplace equation. arXiv:2210.05141 (2022)
- [26] Talenti, G.: [Best constant in Sobolev inequality](#). Ann. Mat. Pura Appl. (4) **110**, 353–372 (1976) Zbl 0353.46018 MR 463908
- [27] Tashiro, Y.: [Complete Riemannian manifolds and some vector fields](#). Trans. Amer. Math. Soc. **117**, 251–275 (1965) Zbl 0136.17701 MR 174022
- [28] Vétois, J.: [A note on the classification of positive solutions to the critical  \$p\$ -Laplace equation in  \$\mathbb{R}^n\$](#) . Adv. Nonlinear Stud. **24**, 543–552 (2024) Zbl 07857773 MR 4747988
- [29] Wang, F., Zhu, X.: [The structure of spaces with Bakry–Émery Ricci curvature bounded below](#). J. Reine Angew. Math. **757**, 1–50 (2019) Zbl 1429.53061 MR 4036569
- [30] Wang, Y. D.: [Harmonic maps from noncompact Riemannian manifolds with non-negative Ricci curvature outside a compact set](#). Proc. Roy. Soc. Edinburgh Sect. A **124**, 1259–1275 (1994) Zbl 0827.53035 MR 1313202
- [31] Wei, G.: [Yamabe equation on some complete noncompact manifolds](#). Pacific J. Math. **302**, 717–739 (2019) Zbl 1434.53042 MR 4036748