

A note on traces of operators on $L^\infty(\lambda)$

Marian Nowak

Abstract. Let $(\Omega, \Sigma, \lambda)$ be a finite measure space and $\tau(L^\infty(\lambda), L^1(\lambda))$ be a natural Mackey topology on $L^\infty(\lambda)$. Let $T : L^\infty(\lambda) \rightarrow L^\infty(\lambda)$ be a Bochner representable operator, that is, there exists $g \in L^1(\lambda, L^\infty(\lambda))$ so that $T(u) = \int_\Omega u(\omega)g(\omega) d\lambda(\omega)$ for $u \in L^\infty(\lambda)$. It is shown that T is a nuclear operator between the locally convex space $(L^\infty(\lambda), \tau(L^\infty(\lambda), L^1(\lambda)))$ and the Banach space $L^\infty(\lambda)$ and T has a well-defined trace: $\text{tr } T = \int_\Omega g(\omega)(\omega) d\lambda(\omega)$.

1. Introduction and preliminaries

Formulas for traces of kernel operators on Banach function spaces (in particular, $C(\Omega)$, $L^p(\lambda)$ -spaces) have been the object of much study (see [2, 3, 7, 9, 10, 13]). In particular, Grothendieck [9] showed that if Ω is a compact Hausdorff space with a positive Borel measure λ on Ω and $k(\cdot, \cdot) \in C(\Omega \times \Omega)$, then the kernel operator $T : C(\Omega) \rightarrow C(\Omega)$ defined by

$$T(u) := \int_\Omega u(\omega)k(\cdot, \omega) d\lambda(\omega) \quad \text{for } u \in C(\Omega),$$

is nuclear and has a well-defined trace $\text{tr } T = \int_\Omega k(\omega, \omega) d\lambda(\omega)$ (see [13, Theorem, p. 274]).

Grothendieck [8, Chap. I, p. 165] proved that the notion of “trace” can be defined for nuclear operators on Banach spaces with the approximation property (see [13, Lemma, pp. 210–211]).

From now on, let $(X, \|\cdot\|_X)$ be a real Banach space and X^* stand for its Banach dual. Let $\mathcal{L}(X, X)$ denote the Banach space of all bounded linear operators $T : X \rightarrow X$, equipped with the operator norm.

Assume that $T : X \rightarrow X$ is a nuclear operator, that is, there exist a bounded sequence (F_n) in X^* , a bounded sequence (x_n) in X and a sequence $(\alpha_n) \in \ell^1$ so

Mathematics Subject Classification 2020: 47B10 (primary); 46G10, 46E30 (secondary).

Keywords: Bochner representable operators, nuclear operators, Mackey topology, trace of operator, tensor products.

that

$$T = \sum_{n=1}^{\infty} \alpha_n F_n \otimes x_n \quad \text{in } \mathcal{L}(X, X), \tag{1.1}$$

where $(\alpha_n F_n \otimes x_n)(x) = \alpha_n F_n(x)x_n$ for $x \in X$.

If X has the approximation property, then the trace of T is given by

$$\text{tr } T = \sum_{n=1}^{\infty} \alpha_n F_n(x_n)$$

and it does not depend on the special choice of the nuclear representation (1.1) of T (see [7, Chap. 5, Theorem 1.2], [13, Lemma, pp. 210–211]).

Assume that $(\Omega, \Sigma, \lambda)$ is a complete finite measure space. Then the space

$$L^\infty(\lambda) := \{u \in L^0(\lambda) : \|u\|_\infty := \text{ess sup}_{\omega \in \Omega} |u(\omega)| < \infty\},$$

equipped with the norm $\|\cdot\|_\infty$ is a Dedekind complete Banach lattice. It is known that $L^\infty(\lambda)$ has the approximation property (see [5, p. 245], [14]).

Let $L^1(\lambda, X)$ stand for the Banach space of λ -equivalence classes of all Bochner integrable functions $g : \Omega \rightarrow X$, equipped with the norm $\|g\|_1 := \int_\Omega \|g(\omega)\|_X d\lambda(\omega)$.

Definition 1.1. A bounded linear operator $T : L^\infty(\lambda) \rightarrow X$ is said to be *Bochner representable*, if there exists a unique $g \in L^1(\lambda, X)$ such that

$$T(u) = \int_\Omega u(\omega)g(\omega) d\lambda(\omega) \quad \text{for } u \in L^\infty(\lambda).$$

The aim of this paper is to find a formula for the trace of a Bochner representable operator $T : L^\infty(\lambda) \rightarrow L^\infty(\lambda)$.

2. Traces of Bochner representable operators on $L^\infty(\lambda)$

Recall that a linear functional F on $L^\infty(\lambda)$ is said to be σ -order continuous if $F(u_n) \rightarrow 0$ whenever $u_n(\omega) \rightarrow 0$ μ -a.e. and $\sup_n \|u_n\|_\infty < \infty$ (see [11, Chap. 6]).

Let $L^\infty(\lambda)_c^*$ stand for the linear subspace of $L^\infty(\lambda)^*$ consisting of all σ -order continuous functionals. Then we have (see [11, Theorem 6.6.1])

$$L^\infty(\lambda)_c^* = \{F_v : v \in L^1(\lambda)\},$$

where $F_v(u) := \int_\Omega u(\omega)v(\omega) d\lambda(\omega)$ for $u \in L^\infty(\lambda)$ and $\|F_v\| = \|v\|_1$.

Let $\tau(L^\infty(\lambda), L^1(\lambda))$ denote the Mackey topology on $L^\infty(\lambda)$, with respect to the dual pair $\langle L^\infty(\lambda), L^\infty(\lambda)_c^* \rangle$ with the duality

$$\langle u, F_v \rangle := F_v(u) = \int_\Omega u(\omega)v(\omega) d\lambda(\omega) \quad \text{for } u \in L^\infty(\lambda), v \in L^1(\lambda).$$

This means that $\tau(L^\infty(\lambda), L^1(\lambda))$ is the topology of uniform convergence on the family of all convex, circled $\sigma(L^\infty(\lambda)_c^*, L^\infty(\lambda))$ -compact sets in $L^\infty(\lambda)_c^*$ (see [1, p. 143]). It follows that every convex, circled $\sigma(L^\infty(\lambda)_c^*, L^\infty(\lambda))$ -compact set in $L^\infty(\lambda)_c^*$ is $\tau(L^\infty(\lambda), L^1(\lambda))$ -equicontinuous. It is known that $\tau(L^\infty(\lambda), L^1(\lambda))$ is a locally convex-solid Hausdorff topology with the Lebesgue property (see [6, Corollary 83H]).

Nuclear operators $T : L^\infty(\lambda) \rightarrow X$ between the Banach spaces $L^\infty(\lambda)$ and X have been studied by Swartz [16] and Nowak [12].

Grothendieck [8,9] introduced the concept of nuclear operators for locally convex spaces (see [15, Chap. 3, § 7], [17, § 47]).

Definition 2.1. A linear operator $T : L^\infty(\lambda) \rightarrow X$ is said to be a *nuclear operator* between the locally convex space $(L^\infty(\lambda), \tau(L^\infty(\lambda), L^1(\lambda)))$ and the Banach space X if there exist a $\tau(L^\infty(\lambda), L^1(\lambda))$ -equicontinuous sequence (F_n) in $L^\infty(\lambda)_c^*$, a bounded sequence (x_n) in X and a sequence $(\alpha_n) \in \ell^1$ so that

$$T(u) = \sum_{n=1}^\infty \alpha_n F_n(u)x_n \quad \text{for } u \in L^\infty(\lambda),$$

that is,

$$T(u) = \sum_{n=1}^\infty \alpha_n F_n \otimes x_n \quad \text{for } \mathcal{L}(L^\infty(\lambda), X).$$

Now we can state our main result.

Theorem 2.1. Assume that $T : L^\infty(\lambda) \rightarrow L^\infty(\lambda)$ is a Bochner representable operator, that is, there exists a unique $g \in L^1(\lambda, L^\infty(\lambda))$ so that

$$T(u) = \int_\Omega u(\omega)g(\omega) d\lambda(\omega) \quad \text{for all } u \in L^\infty(\lambda).$$

Then T is a nuclear operator between the locally convex space $(L^\infty(\lambda), \tau(L^\infty(\lambda), L^1(\lambda)))$ and the Banach space $L^\infty(\lambda)$ and has a well-defined trace

$$\text{tr } T = \int_\Omega g(\omega)(\omega) d\lambda(\omega).$$

Proof. Let $L^1(\lambda) \widehat{\otimes}_\pi L^\infty(\lambda)$ denote the projective tensor product of the Banach spaces $L^1(\lambda)$ and $L^\infty(\lambda)$, equipped with the completed norm π (see [5, p. 227], [14, p. 17]). Note that for $z \in L^1(\lambda) \widehat{\otimes}_\pi L^\infty(\lambda)$, we have

$$\pi(z) = \inf \left\{ \sum_{n=1}^{\infty} \alpha_n \|v_n\|_1 \|w_n\|_\infty \right\},$$

where the infimum is taken over all sequences (v_n) in $L^1(\lambda)$ and (w_n) in $L^\infty(\lambda)$ with $\lim \|v_n\|_1 = 0 = \lim \|w_n\|_\infty$ and $(\alpha_n) \in \ell^1$ such that $z = \sum_{n=1}^{\infty} \alpha_n v_n \otimes w_n$ in the norm π (see [14, Proposition 2.8]).

Note that $L^1(\lambda) \widehat{\otimes}_\pi L^\infty(\lambda)$ is isometrically isomorphic to the Banach space $L^1(\lambda, L^\infty(\lambda))$ by the isometry J , defined by

$$J(v \otimes w) := v(\cdot)w \quad \text{for } v \in L^1(\lambda), w \in L^\infty(\lambda)$$

(see [5, Example 10, p. 228], [14, Example 2.19, p. 29]). Then there exist sequences (v_n) in $L^1(\lambda)$ and (w_n) in $L^\infty(\lambda)$ with $\lim \|v_n\|_1 = 0 = \lim \|w_n\|_\infty$ and $(\alpha_n) \in \ell^1$ so that

$$J^{-1}(g) = \sum_{n=1}^{\infty} \alpha_n v_n \otimes w_n \quad \text{in } L^1(\lambda) \widehat{\otimes}_\pi L^\infty(\lambda).$$

Thus it follows that

$$g = J\left(\sum_{n=1}^{\infty} \alpha_n v_n \otimes w_n\right) = \sum_{n=1}^{\infty} \alpha_n v_n(\cdot)w_n \quad \text{in } L^1(\lambda, L^\infty(\lambda)),$$

and hence

$$g(\omega) = \sum_{n=1}^{\infty} \alpha_n v_n(\omega)w_n \quad \text{for } \omega \in \Omega.$$

Then for every $u \in L^\infty(\lambda)$, we get

$$T(u) = \int_{\Omega} u(\omega) \left(\sum_{n=1}^{\infty} \alpha_n v_n(\omega)w_n \right) d\lambda(\omega) = \sum_{n=1}^{\infty} \alpha_n \left(\int_{\Omega} u(\omega)v_n(\omega) d\lambda(\omega) \right) w_n.$$

Since $\lim \|v_n\|_1 = 0$, the set $\{v_n : n \in \mathbb{N}\}$ is uniformly integrable and in view of the Dunford–Pettis theorem (see [4, Theorem, p. 93]), we obtain that $\{v_n : n \in \mathbb{N}\}$ is a relatively $\sigma(L^1(\lambda), L^\infty(\lambda))$ -compact subset of $L^1(\lambda)$. Then by the Krein–Smulian theorem (see [1, Theorem 10.15]), the convex, circled hull of the set $\{v_n : n \in \mathbb{N}\}$ is relatively $\sigma(L^1(\lambda), L^\infty(\lambda))$ -compact in $L^1(\lambda)$, and hence the convex, circled hull of the set $\{F_{v_n} : n \in \mathbb{N}\}$ is relatively $\sigma(L^\infty(\lambda)_c^*, L^\infty(\lambda))$ -compact in $L^\infty(\lambda)_c^*$. Hence the set $\{F_{v_n} : n \in \mathbb{N}\}$ is $\tau(L^\infty(\lambda), L^1(\lambda))$ -equicontinuous in $L^\infty(\lambda)_c^*$.

This means that T is a nuclear operator between the locally convex space $(L^\infty(\lambda), \tau(L^\infty(\lambda), L^1(\lambda)))$ and the Banach space $L^\infty(\lambda)$ (see Definition 2.1). It follows that T is a nuclear operator between the Banach spaces $L^\infty(\lambda)$ and $L^\infty(\lambda)$ and we have

$$T = \sum_{n=1}^{\infty} \alpha_n F_{v_n} \otimes w_n \quad \text{in } \mathcal{L}(L^\infty(\lambda), L^\infty(\lambda)).$$

Hence, since $L^\infty(\lambda)$ has the approximation property, we get

$$\text{tr } T = \sum_{n=1}^{\infty} \alpha_n F_{v_n}(w_n) = \sum_{n=1}^{\infty} \alpha_n \int_{\Omega} v_n(\omega) w_n(\omega) d\lambda(\omega). \tag{2.1}$$

For $n \in \mathbb{N}$, let $g_n := \sum_{i=1}^n \alpha_i v_i(\cdot) w_i$, that is, $g_n(\omega) = \sum_{i=1}^n \alpha_i v_i(\omega) w_i$ for $\omega \in \Omega$.

Then we have

$$\begin{aligned} & \left| \int_{\Omega} g(\omega)(\omega) d\lambda(\omega) - \sum_{i=1}^n \alpha_i \int_{\Omega} v_i(\omega) w_i(\omega) d\lambda(\omega) \right| \\ &= \left| \int_{\Omega} (g(\omega)(\omega) - g_n(\omega)(\omega)) d\lambda(\omega) \right| \\ &\leq \int_{\Omega} |g(\omega)(\omega) - g_n(\omega)(\omega)| d\lambda(\omega) \\ &\leq \int_{\Omega} \|g(\omega) - g_n(\omega)\|_{\infty} d\lambda(\omega) \\ &= \|g - g_n\|_1 \xrightarrow{n} 0. \end{aligned}$$

Thus

$$\int_{\Omega} g(\omega)(\omega) d\lambda(\omega) = \sum_{n=1}^{\infty} \alpha_n \int_{\Omega} v_n(\omega) w_n(\omega) d\lambda(\omega),$$

and in view of (2.1), we get

$$\text{tr } T = \int_{\Omega} g(\omega)(\omega) d\lambda(\omega). \quad \blacksquare$$

References

[1] C. D. Aliprantis and O. Burkinshaw, *Positive operators*. Pure Appl. Math. 119, Academic Press, Inc., Orlando, FL, 1985 MR 0809372

[2] C. Brislawn, [Kernels of trace class operators](#). *Proc. Amer. Math. Soc.* **104** (1988), no. 4, 1181–1190 Zbl 0695.47017 MR 0929421

- [3] J. Delgado, [A trace formula for nuclear operators on \$L^p\$](#) . In *Pseudo-differential operators: complex analysis and partial differential equations*, pp. 181–193, Oper. Theory Adv. Appl. 205, Birkhäuser, Basel, 2010 Zbl [1229.47051](#) MR [2664581](#)
- [4] J. Diestel, [Sequences and series in Banach spaces](#). Grad. Texts in Math. 92, Springer, New York, 1984 Zbl [0542.46007](#) MR [0737004](#)
- [5] J. Diestel and J. J. Uhl, Jr., [Vector measures](#). Math. Surv. 15, American Mathematical Society, Providence, RI, 1977 Zbl [0369.46039](#) MR [0453964](#)
- [6] D. H. Fremlin, [Topological Riesz spaces and measure theory](#). Cambridge University Press, London-New York, 1974 Zbl [0273.46035](#) MR [0454575](#)
- [7] I. Gohberg, S. Goldberg, and N. Krupnik, [Traces and determinants of linear operators](#). Oper. Theory Adv. Appl. 116, Birkhäuser, Basel, 2000 Zbl [0946.47013](#) MR [1744872](#)
- [8] A. Grothendieck, [Produits tensoriels topologiques et espaces nucléaires](#). Mem. Amer. Math. Soc. **16**, American Mathematical Society, 1955 Zbl [0064.35501](#) MR [0075539](#)
- [9] A. Grothendieck, [La théorie de Fredholm](#). *Bull. Soc. Math. France* **84** (1956), 319–384 MR [0088665](#)
- [10] C. W. Ha, On the trace of a class of nuclear operators. *Bull. Inst. Math. Acad. Sinica* **3** (1975), no. 1, 131–137 Zbl [0307.47024](#) MR [0394268](#)
- [11] L. V. Kantorovich and G. P. Akilov, [Functional analysis](#). Second edn., Pergamon Press, Oxford-Elmsford, New York, 1982 Zbl [0484.46003](#) MR [0664597](#)
- [12] M. Nowak, [Nuclear operators on Banach function spaces](#). *Positivity* **25** (2021), no. 3, 801–818 Zbl [1494.47058](#) MR [4274292](#)
- [13] A. Pietsch, [Eigenvalues and \$s\$ -numbers](#). *Mathematik und ihre Anwendungen in Physik und Technik* 43, Akademische Verlagsgesellschaft Geest & Portig K.-G., Leipzig, 1987 Zbl [0615.47019](#) MR [0917067](#)
- [14] R. A. Ryan, [Introduction to tensor products of Banach spaces](#). Springer Monogr. Math., Springer London, Ltd., London, 2002 Zbl [1090.46001](#) MR [1888309](#)
- [15] H. H. Schaefer, [Topological vector spaces](#). Graduate Texts in Mathematics, Vol 3, Springer, New York-Berlin, 1971 Zbl [0212.14001](#) MR [0342978](#)
- [16] C. Swartz, [An operator characterization of vector measures which have Radon–Nikodym derivatives](#). *Math. Ann.* **202** (1973), 77–84 Zbl [0236.46054](#) MR [0325917](#)
- [17] F. Trèves, [Topological vector spaces, distributions and kernels](#). Academic Press, New York-London, 1967 Zbl [0171.10402](#) MR [0225131](#)

Received 7 November 2024.

Marian Nowak

Institute of Mathematics, University of Zielona Góra, ul. Szafrana 4A, 65–516 Zielona Góra, Poland; m.nowak@wmie.uz.zgora.pl