

On the free boundary for thin obstacle problems with Sobolev variable coefficients

Giovanna Andreucci, Matteo Focardi, and Emanuele Spadaro

Abstract. We establish a quasi-monotonicity formula for an intrinsic frequency function related to solutions to thin obstacle problems with zero obstacle driven by quadratic energies with Sobolev $W^{1,p}$ coefficients, where p is bigger than the space dimension. From this, we deduce several regularity and structural properties of the corresponding free boundaries at those distinguished points with finite order of contact with the obstacle. In particular, we prove the rectifiability and the local finiteness of the Minkowski content of the whole free boundary in the case of Lipschitz coefficients.

1. Introduction

In this article, we consider a class of lower-dimensional obstacle problems with variable coefficients which has been extensively considered in the literature. In order to state the results, we introduce the following notation: for any subset $E \subset \mathbb{R}^{n+1}$, we set

$$E^+ := E \cap \{x \in \mathbb{R}^{n+1} : x_{n+1} > 0\} \quad \text{and} \quad E' := E \cap \{x_{n+1} = 0\}.$$

For any point $x \in \mathbb{R}^{n+1}$, we will write

$$x = (x', x_{n+1}) \in \mathbb{R}^n \times \mathbb{R}.$$

Moreover, $B_r(x) \subset \mathbb{R}^{n+1}$ denotes the open ball centered at $x \in \mathbb{R}^{n+1}$ with radius $r > 0$, and $\overline{B}_r(x)$ its closure (we omit to write the center of the ball x if $x = 0$).

We consider the problem of minimizing a variable coefficient quadratic (*Dirichlet*) energy with a unilateral constraint:

$$\min_{v \in \mathcal{A}} \int_{B_1^+} \langle \mathbb{A}(x) \nabla v, \nabla v \rangle \, dx, \tag{1.1}$$

where the class of competing functions is given by

$$\mathcal{A} := \{v \in H^1(B_1^+) : v = g \text{ in } (\partial B_1)^+ \text{ and } v \geq 0 \text{ in } B_1^+\},$$

with $g \in H^{\frac{1}{2}}((\partial B_1)^+)$ such that \mathcal{A} is not empty. (The boundary conditions are meant in

the sense of traces.) We assume the following hypotheses:

- (H1) $\mathbb{A} \in W^{1,p}(B_1, \mathbb{R}^{(n+1) \times (n+1)})$, $p \in (n + 1, \infty]$ (in particular, by Morrey’s embedding, we have $\mathbb{A} \in C^{0,\alpha}(B_1, \mathbb{R}^{(n+1) \times (n+1)})$, $\alpha := 1 - \frac{n+1}{p}$);
- (H2) $\mathbb{A}(x) = (a_{ij}(x))_{i,j=1}^{n+1}$ is a symmetric, bounded, and coercive matrix, i.e., for every $x \in B_1^+$, $i, j \in \{1, \dots, n + 1\}$, and $\xi \in \mathbb{R}^{n+1}$ $a_{ij}(x) = a_{ji}(x)$, and

$$\lambda|\xi|^2 \leq \langle \mathbb{A}(x)\xi, \xi \rangle \leq \Lambda|\xi|^2,$$

for some $0 < \lambda \leq \Lambda$,

By a well-known result due to Ural’tseva [38], the minimizers of (1.1) are $C^{1,\beta}$ regular for some $\beta > 0$ and satisfy an elliptic partial differential equation in the half-ball B_1^+ , whereas on the flat part of the boundary B_1' the unilateral constraint $u \geq 0$ leads to a free boundary problem. The Euler–Lagrange equation of (1.1) is

$$\begin{cases} \operatorname{div}(\mathbb{A}(x)\nabla u(x)) = 0 & \text{in } B_1^+, \\ u \geq 0, \quad \mathbb{A}\nabla u \cdot e_{n+1} \leq 0, \quad u(\mathbb{A}\nabla u \cdot e_{n+1}) = 0 & \text{in } B_1'. \end{cases} \tag{1.2}$$

The condition $u(\mathbb{A}\nabla u \cdot e_{n+1}) = 0$ in B_1' is the so-called *Signorini complementary* boundary condition. The behavior of u on B_1' is not prescribed and is characterized by the so-called *free boundary* $\Gamma(u)$, which is the relative boundary in B_1' of the contact set $B_1' \cap \{u = 0\}$, where the solution saturates the constraint.

This problem has been widely studied in the last decades and has become a very active field of research after the seminal papers by Athanasopoulos and Caffarelli [4] and Athanasopoulos, Caffarelli, and Salsa [5]. The key idea introduced in [5] is the use of *Almgren’s frequency function* in the study of both the regularity of the solution u and the properties of the free boundary $\Gamma(u)$. This has been the turning point for a long series of results for the constant coefficient case, leading to a detailed analysis of the free boundary (see, e.g., [5, 7, 8, 14–16, 19, 23, 26] and the references therein). The regularity of the solutions of the variable coefficients case has been considered since the works of Caffarelli [6] and Kinderlehrer [25] in the case of smooth coefficients, whereas problem (1.1) with Sobolev coefficients has been considered by Ural’tseva in [36–38]. The optimal regularity of the solutions and the regularity of a subset of the free boundary (points with selected orders of contact) have been proven more recently by Garofalo, and Smit Vega Garcia [22] and Garofalo, Petrosyan and Smit Vega Garcia in [20], respectively, for Lipschitz coefficients using a generalization of the frequency function (see also [21] for further results in the Lipschitz setting). In case of Sobolev coefficients, both topics have been addressed by Koch, Ruland, and Shi [27, 28] by means of Carleman inequalities, the optimal regularity of solutions is established by Ruland and Shi [34] for Holder coefficients, and for a more general notion of quasi-minimizers by Jeon, Petrosyan, and Smit Vega Garcia [24].

In this paper, we continue the analysis for quadratic energies with matrix field with Sobolev regularity. In particular, we address the question of the global structure of the

free boundary. In details, we need to consider only the points with finite order of contact: to this aim, we write

$$\Gamma(u) := \Gamma^{\text{finite}} \cup \Gamma^\infty,$$

where Γ^{finite} and Γ^∞ , to be properly defined in the next sections, represent the points with finite and infinite order of contact. For Γ^∞ , no structure at all is expected, in analogy with the case of nonzero obstacles studied in [10, 18], and the results on the lack of unique continuation by Pliš [33], Miller [30], Filonov [11], and Mandache [29] for solutions to second-order elliptic partial differential equations with Hölder coefficients. We show that the set of free boundary points with finite order of contact is *rectifiable*; i.e., can be stratified along submanifolds of dimension $n - 1$ and class C^1 .

Theorem 1.1. *Let $u \in \mathcal{A}$ be a solution to (1.1) under the hypotheses (H1) and (H2). The subset of points of the free boundary with finite order of contact $\Gamma^{\text{finite}}(u)$ is $(n - 1)$ -rectifiable, i.e., there exists a countable family of C^1 -submanifolds $M_i \subset B'_1$ of dimension $n - 1$ such that*

$$\mathcal{H}^{n-1}\left(\Gamma^{\text{finite}}(u) \setminus \bigcup_{i \in \mathbb{N}} M_i\right) = 0.$$

Furthermore, there exists a set $\Sigma(u) \subset \Gamma^{\text{finite}}(u)$ with Hausdorff dimension at most $n - 2$ such that for every $x \in \Gamma^{\text{finite}}(u) \setminus \Sigma(u)$

$$N_u(x, 0^+) \in \{2m, 2m - 1/2, 2m + 1\}_{m \in \mathbb{N} \setminus \{0\}}.$$

In the statement above, $N_u(x, 0^+)$ represents the intrinsic frequency of u at the free boundary point x (cf. Section 5 for its definition).

In addition, if \mathbb{A} is Lipschitz continuous, a more complete result holds for the whole free boundary $\Gamma(u)$, completely analogous to the case of the classical scalar Signorini problem for the Dirichlet energy, as shown in [15, 16].

Theorem 1.2. *Let $u \in \mathcal{A}$ be a solution to (1.1) under the hypotheses (H1) with $p = \infty$ and (H2). Then, the free boundary $\Gamma(u)$ is $(n - 1)$ -rectifiable with locally finite Minkowski content: for every $K \subset\subset B'_1$, there exists a constant $C(K) > 0$ such that*

$$\mathcal{L}^{n+1}(\mathcal{T}_r(\Gamma(u) \cap K)) \leq C(K)r^2, \quad \forall r \in (0, 1),$$

where $\mathcal{T}_r(\Gamma(u) \cap K) \subset \mathbb{R}^{n+1}$ is the r -tubular neighborhood of $\Gamma(u) \cap K$.

Furthermore, there exists a set $\Sigma(u) \subset \Gamma^{\text{finite}}(u)$ with Hausdorff dimension at most $n - 2$ such that for every $x \in \Gamma^{\text{finite}}(u) \setminus \Sigma(u)$

$$N_u(x, 0^+) \in \{2m, 2m - 1/2, 2m + 1\}_{m \in \mathbb{N} \setminus \{0\}}.$$

Theorems 1.1 and 1.2 are a natural development of a common trend of recent results, which are directed to the understanding the robustness of the techniques conceived for the Laplacian (i.e., constant coefficients operators) in more general cases (and more realistic from the point of view of applications). In the setting of the thin obstacle problem, we

recall the recent contributions on the structure of the regular set for Lipschitz coefficients [20], for Sobolev coefficients [28], and for Hölder continuous coefficients [24]; for the structure of the singular set with Lipschitz coefficients [21], and [24] for Hölder coefficients. The rectifiability of the whole free boundary has been addressed in [17] for the nonlinear case of the area functional.

One motivation for this study is related to the standard thin obstacle problem, provided the obstacle condition is assigned on a $C^{1,1}$ manifold rather than on a hyperplane. Indeed, a rectification of the manifold leads to a thin obstacle problem as the one stated in (1.1). A further motivation is contained in Section 7.1. Our results allow to deduce the global structure of the free boundary for solutions to some nonlinear thin obstacle problems, following the approach used for the area functional in [17] (cf. [1,9] for preliminary results on the regularity of the solutions).

1.1. New insights and main difficulties

The main ideas for this work stem from [16]. Starting from the groundbreaking papers by Naber and Valtorta [31, 32], it is well known that a monotone quantity of the type of the energy ratio for harmonic maps can be used to describe the structure of singularities: indeed, if the monotone quantity is able to detect homogeneous blowups at singular points and satisfies a suitable rigidity property, then general covering and rectifiability arguments lead to the estimate of the measure (actually the Minkowski content) and the rectifiability of the singular set.

This principle has been exploited in [16] in the case of thin and fractional obstacle problems with constant coefficients, using suitable variants of Almgren's frequency functions, which revealed itself to be a key tool for this class of problems since [5].

The main difficulties here are due to extension of such approach to the case of variable coefficients. Indeed, the monotonicity of the frequency is closely related to the linearity of the equations and is valid only for harmonic functions, while in the general case, one should find a suitable linear approximation of the local geometry prescribed by the matrix \mathbb{A} . This is clearly an issue for general nonlinear problems, and understanding this question for low regularity matrix fields \mathbb{A} is a first step towards such program.

We circumvent this difficulty by introducing an intrinsic frequency adapted to the coefficients matrix \mathbb{A} (as opposed to the natural frequency for variable coefficients, see Section 5 for more details). Actually, we need to use three different forms of the frequency, which, although different, can be suitably compared at the right scales. In particular, we show a quasi-monotonicity formula for a Dirichlet-type frequency for solutions to variable coefficient thin obstacle problems. This idea has been used for the analysis of the classical obstacle problem in [3, 13]. In the current setting, we couple it with the fundamental insight provided by Simon and Wickramasekera in the framework of 2-valued minimal graphs, that quasi-monotonicity of the frequency is actually equivalent to a doubling condition on the relevant quantities provided Schauder estimates hold (cf. [35, Lemma 6.1]).

A comment deserves the restriction to Sobolev coefficients as opposite to the more general Hölder setting, for which the analysis of regular and singular points is contained in [23, 24]. In the derivation of the basic estimate on the oscillation of the frequency, as well as for the monotonicity, we differentiate the matrix of coefficients and the gradient of the solutions, and therefore, we need enough regularity for \mathbb{A} .

2. Preliminaries on the thin obstacle problem

Here, we recall the hypotheses on the thin obstacle problem we address. We consider quadratic energies of the form

$$\int_{B_1^+} \langle \mathbb{A}(x) \nabla v(x), \nabla v(x) \rangle dx,$$

where the matrix field \mathbb{A} satisfies the following hypotheses:

(H1) $\mathbb{A} \in W^{1,p}(B_1, \mathbb{R}^{(n+1) \times (n+1)})$, $p \in (n + 1, \infty]$ (in particular, by Morrey’s embedding, we have $\mathbb{A} \in C^{0,\alpha}(B_1, \mathbb{R}^{(n+1) \times (n+1)})$, $\alpha := 1 - \frac{n+1}{p}$);

(H2) $\mathbb{A}(x) = (a_{ij}(x))_{i,j=1}^{n+1}$ is a symmetric, bounded, and coercive matrix, i.e., for every $x \in B_1^+$, $i, j \in \{1, \dots, n + 1\}$, and $\xi \in \mathbb{R}^{n+1}$ $a_{i,j}(x) = a_{j,i}(x)$, and

$$\lambda |\xi|^2 \leq \langle \mathbb{A}(x) \xi, \xi \rangle \leq \Lambda |\xi|^2,$$

for some $0 < \lambda \leq \Lambda$.

Following [37, Remark 1], by a means of a change of variables, it is not restrictive to additionally assume that

(H3) $a_{i,n+1}(x', 0) = 0$ for all $i = 1, \dots, n$ for every $x \in B'_1$.

Under these hypotheses, we can extend all the functions on B_1 by even symmetry: with a slight abuse of notation, we set

$$\mathbb{A}(x', x_{n+1}) = \mathbb{A}(x', -x_{n+1}), \quad u(x', x_{n+1}) = u(x', -x_{n+1}) \quad \forall x \in B_1^+.$$

In this way, it is equivalent to formulating the problem on B_1 with the symmetry condition

$$\min_{v \in \mathcal{A}} \int_{B_1} \langle \mathbb{A}(x) \nabla v(x), \nabla v(x) \rangle dx, \tag{2.1}$$

in the class of functions

$$\mathcal{A} := \{v \in g + H_0^1(B_1) : v(x', x_{n+1}) = v(x', -x_{n+1}), v(x', 0) \geq 0\},$$

with $g \in H^1(B_1)$ even symmetric and such that $g \geq 0$ on B'_1 .

By a result due to Ural'tseva [38, Theorem 3.1], the solution of the thin obstacle problem under assumptions (H1), (H2), and (H3) have Hölder continuous first derivative up to B'_1 (for the optimal Hölder exponent, see [27, 28]).

Theorem 2.1 ([38]). *For every $g \in H^1(B_1)$, even symmetric with $g \geq 0$ on B'_1 , there exists a unique solution $u \in \mathcal{A}$ to the thin obstacle problem (2.1). Moreover, $u \in H^2_{\text{loc}} \cap C^{1,\beta}_{\text{loc}}(B_1^+ \cup B'_1)$ for some $\beta \in (0, 1)$, and*

$$\|u\|_{H^2(B_{1/2}^+ \cup B'_{1/2})} + \|u\|_{C^{1,\beta}(B_{1/2}^+ \cup B'_{1/2})} \leq C \|u\|_{L^2(B_1)},$$

where $C = C(p, n, \beta, \|\mathbb{A}\|_{W^{1,p}}) > 0$.

The Euler–Lagrange equations satisfied by the solution u to the thin obstacle problem are then the following:

$$\begin{cases} \operatorname{div}(\mathbb{A}(x)\nabla u(x)) = 0 & \text{for } x \in B_1 \setminus \{(x', 0) : u(x', 0) = 0\}, \\ \operatorname{div}(\mathbb{A}(x)\nabla u(x)) \leq 0 & \text{in the sense of distribution in } B_1. \end{cases} \quad (2.2)$$

Moreover, in view of the assumptions (H2) and (H3), the Signorini complementary condition in (1.2) then reads as

$$u \partial_{n+1} u = 0 \quad \text{on } B'_1.$$

In the sequel, u will always denote a solution to the thin obstacle problem (2.1), unless otherwise stated.

3. The frequency function

In this section, we introduce a suitable version of Almgren’s frequency function. Let $\phi : [0, \infty) \rightarrow [0, \infty)$ be a decreasing $C^{1,1}$ function such that $\phi'(t) < 0$ if $\frac{1}{2} < t < 1$ and

$$\phi(t) = \begin{cases} 1 & \text{for } 0 \leq t \leq \frac{1}{2}, \\ > 0 & \text{for } \frac{1}{2} < t < 1, \\ 0 & \text{for } 1 \leq t. \end{cases}$$

For the sake of simplicity, we assume that

$$\phi(t) = 2(1 - t) \quad \forall t \in \left[\frac{5}{8}, \frac{7}{8}\right].$$

We define the frequency function of a function u at a point $x_0 \in B'_1$ by

$$I_u(x_0, r) := \frac{r D_u(x_0, r)}{H_u(x_0, r)} \quad \forall r < 1 - |x_0|,$$

where the Dirichlet energy is given by

$$D_u(x_0, r) := \int \phi\left(\frac{|x - x_0|}{r}\right) |\nabla u(x)|^2 dx,$$

and the “boundary” L^2 norm of u is given by

$$H_u(x_0, r) := - \int \phi' \left(\frac{|x - x_0|}{r} \right) \frac{u^2(x)}{|x - x_0|} dx.$$

Note that the frequency function is well defined as long as $H_u(x_0, r) > 0$. In what follows, when u is a solution to (2.1), we will tacitly assume that the latter condition is satisfied when writing $I_u(x_0, r)$. Indeed, $H_u(x_0, r) > 0$ if $x_0 \in \Gamma(u)$ by minimality of u and the uniqueness of minimizers. Otherwise, $u \equiv 0$ on $B_r(x_0) \setminus B_{r/2}(x_0)$, in turn implying $u \equiv 0$ on $B_r(x_0)$. This contradicts the fact that x_0 is a free boundary point. Analogously, if $x_0 \in \Gamma(u)$, then $D_u(x_0, r) > 0$.

Additionally, for later convenience, we introduce the following quantities:

$$G_u(x_0, r) := -r^{-1} \int \phi' \left(\frac{|x - x_0|}{r} \right) u(x) \nabla u(x) \cdot \frac{x - x_0}{|x - x_0|} dx$$

and

$$E_u(x_0, r) := - \int \phi' \left(\frac{|x - x_0|}{r} \right) \frac{|x - x_0|}{r^2} \left(\nabla u(x) \cdot \frac{x - x_0}{|x - x_0|} \right)^2 dx.$$

In particular, note that $E_u(x_0, r)H_u(x_0, r) - G_u^2(x_0, r) \geq 0$ by Cauchy–Schwarz inequality.

Finally, for every $x_0 \in \Gamma(u)$ and $r > 0$, the rescalings of a solution u are given by

$$u_{x_0, r}(y) := \frac{r^{n/2}}{H_u^{1/2}(x_0, r)} u(x_0 + ry) \quad \forall y \in B_{\frac{1-|x_0|}{r}}$$

so that

$$H_{u_{x_0, r}}(\underline{0}, 1) = 1.$$

We will always omit to write the base point x_0 in the notation of I_u, D_u, H_u, E_u, G_u when $x_0 = \underline{0}$.

By a simple corollary of Theorem 2.1, we have the following compactness: if $(u_j)_{j \in \mathbb{N}}$ are such that

$$\sup_{j \in \mathbb{N}} (D_{u_j}(1) + H_{u_j}(1)) < +\infty,$$

then u_j is uniformly bounded in $H^1(B_s)$ for every $s < 1$. Therefore, if u_j are solutions to the thin obstacle problem (2.1) satisfying the hypotheses (H1)–(H3) (holding uniformly in j for varying matrix fields \mathbb{A}_j , with the same constants p, λ, Λ), then by Theorem 2.1 there exist a function $u \in C^{1, \beta}(B_1^+)$ and a subsequence $(u_{j'}) \subset (u_j)$ such that

$$u_{j'} \rightarrow u \quad \text{in } C_{\text{loc}}^{1, \gamma}(B_1^+ \cup B_1') \quad \forall \gamma \in (0, \beta),$$

where β is the constant in Theorem 2.1.

In particular, the following compactness result holds for the rescaling of solutions.

Proposition 3.1. *Let $(u_j)_{j \in \mathbb{N}}$ be a sequence of solutions to the thin obstacle problem (2.1) for varying matrix fields \mathbb{A}_j under assumptions (H1)–(H3) holding uniformly in j , with $x_j \in \Gamma(u_j) \cap B'_{1/2}$ for every $j \in \mathbb{N}$. Assume that*

$$\sup_j I_{u_j}(x_j, \varrho_j) < \infty \quad \text{for some } \varrho_j \downarrow 0.$$

Then, there exist a subsequence (j') such that $x_{j'} \rightarrow x_\infty \in \bar{B}'_{1/2}$ and a function v_∞ such that $v_{j'} := (u_{j'})_{x_{j'}, \varrho_{j'}}$ satisfy as $j' \rightarrow \infty$

$$v_{j'} \rightarrow v_\infty \quad \text{in } C_{\text{loc}}^{1,\gamma}(B_1^+ \cup B_1') \quad \forall \gamma \in (0, \beta), \tag{3.1}$$

$$\underline{0} \in \{x \in \bar{B}_{1/2} : v_\infty(x) = |\nabla v_\infty(x)| = 0\}. \tag{3.2}$$

Moreover, v_∞ is a solution to the thin obstacle problem for the constant coefficients quadratic energy having density $\xi \mapsto \langle \mathbb{A}(x_\infty)\xi, \xi \rangle$.

Proof. From the assumption on the frequency, clearly, the functions $v_j := (u_j)_{x_j, \varrho_j}$ satisfy

$$\sup_{j \in \mathbb{N}} (D_{v_j}(1) + H_{v_j}(1)) < +\infty.$$

Therefore, by compactness, there exists a subsequence (j') such that the points $x_{j'}$ converge to some $x_\infty \in \bar{B}'_{1/2}$, and $v_{j'}$ converge to a limiting function v_∞ in the sense of (3.1). Since $v_{j'}(\underline{0}) = |\nabla v_{j'}(\underline{0})| = 0$ (because $x_{j'} \in \Gamma(u_{j'})$), by the convergence (3.1), we infer also (3.2).

Finally, the fact that v_∞ is a solution to a thin obstacle problem with fixed coefficients $\mathbb{A}(x_\infty)$ follows either by a Γ -convergence result or by passing into the limit in the weak formulation of the Euler–Lagrange equations (2.2) characterizing the solutions, thanks to the convergence (3.1) and the continuity of the matrix field \mathbb{A} . ■

3.1. Doubling estimates

To establish quasi-monotonicity of the frequency at distinguished points of the free boundary, we follow a perturbative approach developed for the classical obstacle in [13]. This approach is coupled with a fundamental insight present in the work of Simon and Wickramasekera [35] on minimal immersion, who highlighted that the doubling condition is equivalent to the quasi-monotonicity of the frequency.

More in detail, we will consider $x_0 \in \Gamma(u)$ satisfying the following hypothesis:

(H4) $x_0 \in \Gamma(u) \cap B'_{1/2}$ such that

$$m(x_0) := \sup_{r \in (0, 1/2)} I_u(x_0, r) < \infty. \tag{3.3}$$

As shown in Appendix A, condition (3.3) is equivalent to assuming that u has a finite order of contact with the null obstacle at $x_0 \in \Gamma(u)$ (cf. the Introduction).

A first step towards the monotonicity of the frequency is to establish a lower bound. A more refined version of the ensuing result will follow after showing the quasi-monotonicity of the frequency (cf. (3.26)). All the constants that will appear in the results below can depend on the parameters p, λ, Λ of (H1)–(H3), even though it will never be explicitly highlighted.

Lemma 3.2. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, then*

$$I_u(x_0, r) \geq C \quad \forall r \in (0, \varrho].$$

Proof. Assume by contradiction that there exist solutions u_j and points $x_j \in \Gamma(u_j) \cap B'_{1/2}$ with $m(x_j) \leq m_0$ for some $m_0 < \infty$ as in the statement such that for a suitable choice of radii $r_j \downarrow 0$ we have that

$$I_{u_j}(x_j, r_j) \leq 1/j.$$

For j sufficiently large $I_{u_j}(x_j, 2r_j) \leq m_0$, and therefore by Proposition 3.1, up to a subsequence not relabeled and keeping the notation introduced there, we conclude that the functions $v_j := (u_j)_{x_j, 2r_j}$ converges in $C^1(B_1)$ to some function v_∞ that minimizes

$$v \mapsto \int_{B_1} \langle \mathbb{A}(x_\infty) \nabla v, \nabla v \rangle dx,$$

among all functions $v \in v_\infty + H_0^1(B_1)$ with $v(x', 0) \geq 0$ on B'_1 . Moreover, by strong convergence, we infer that $H_{v_\infty}(1) = H_{(u_j)_{x_j, 2r_j}}(1) = 1$ and

$$I_{v_\infty}(1/2) = \lim_j I_{(u_j)_{x_j, 2r_j}}(1/2) = \lim_j I_{u_j}(x_j, r_j) = 0.$$

Therefore, $D_{v_\infty}(1/2) = 0$ and thus $v_\infty \equiv 0$ on $B_{1/2}$, a contradiction being v_∞ analytic on $B_1 \setminus B'_1$. ■

Next, we prove the above-mentioned doubling estimates.

Proposition 3.3. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, then*

$$C^{-1} \leq \frac{H_u(x_0, 2r)}{H_u(x_0, r)} \leq C, \quad 1 \leq \frac{D_u(x_0, 2r)}{D_u(x_0, r)} \leq C \quad \forall r \in (0, \varrho].$$

Proof. First, note that by the very definition $D_u(x_0, r) \leq D_u(x_0, 2r)$. Assume by contradiction that there exist solutions u_j , points $x_j \in \Gamma(u_j) \cap B'_{1/2}$ with $m(x_j) \leq m_0$ for some $m_0 < \infty$ as in the statement, and radii $r_j \downarrow 0$ such that

$$\lim_j \frac{H_{u_j}(x_j, 2r_j)}{H_{u_j}(x_j, r_j)} \in \{0, \infty\} \quad \text{and/or} \quad \lim_j \frac{D_{u_j}(x_j, 2r_j)}{D_{u_j}(x_j, r_j)} = \infty.$$

For j sufficiently large, $I_{u_j}(x_j, 2r_j) \leq m_0$; thus, by Proposition 3.1, up to a subsequence not relabeled, $v_j := (u_j)_{x_j, 2r_j}$ converges in $C^1(B_1)$ to some function v_∞ that minimizes

$$v \mapsto \int_{B_1} \langle \mathbb{A}(x_\infty) \nabla v, \nabla v \rangle dx;$$

among all functions, $v \in v_\infty + H_0^1(B_1)$, with $v(x', 0) \geq 0$ on B'_1 . In particular, we conclude that

$$\lim_{j \rightarrow \infty} \frac{H_{u_j}(x_j, 2r_j)}{H_{u_j}(x_j, r_j)} = \frac{H_{v_\infty}(1)}{H_{v_\infty}(1/2)} \in [C^{-1}, C]$$

and

$$\lim_{j \rightarrow \infty} \frac{D_{u_j}(x_j, 2r_j)}{D_{u_j}(x_j, r_j)} = \frac{D_{v_\infty}(1)}{D_{v_\infty}(1/2)} \in [1, C],$$

for some constant $C > 0$ (depending on m_0) because of the doubling estimates satisfied by v_∞ which is a solution to an obstacle problem with constant coefficients and frequency $I_{v_\infty}(1)$ bounded by m_0 (e.g., cf. [16, Corollary 2.8]). This gives the desired contradiction. ■

In a similar fashion, the frequency computed at nearby points can be compared at radii that are bigger than the distance between the points.

Lemma 3.4. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, then for all $r \in (0, \varrho]$, $x \in B'_{r/2}(x_0)$, and $t \in [r, \varrho]$*

$$\frac{H_u(x, t)}{H_u(x_0, t)}, \quad \frac{D_u(x, t)}{D_u(x_0, t)} \in [C^{-1}, C]. \tag{3.4}$$

In particular, the frequency function of u is well defined at every point $x \in B'_{r/2}(x_0)$ at the scales $t \in [r, \varrho]$ and

$$|I_u(x_0, t) - I_u(x, t)| \leq C. \tag{3.5}$$

Remark 3.5. We stress that the conclusions of Lemma 3.4 hold even for points x not necessarily in the free boundary.

Proof. The proof proceeds analogously to the previous ones by contradiction: assume that there exist functions u_j , points $z_j \in \Gamma(u_j) \cap B'_{1/2}$ with $m(z_j) \leq m_0 < \infty$ and $x_j \in B'_{r_j/2}(z_j)$ contradicting one of the two sets of inequalities in (3.4) for some sequence $t_j \in [r_j, \varrho_j]$ with $\varrho_j \downarrow 0$. As above, we can apply Proposition 3.1; thus (up to passing to a subsequence not relabeled), there exists v_∞ such that $v_j := (u_j)_{z_j, t_j} \rightarrow v_\infty$ in $C_{loc}^{1, \beta}(B_2)$ with v_∞ solution to the thin obstacle problem with matrix fields $\mathbb{A}(x_\infty)$. Clearly, we may also assume that $t_j^{-1}(x_j - z_j) \rightarrow y_\infty \in \bar{B}'_{1/2}$ (note that $r_j/t_j \leq 1$).

Assume now that the first set of inequalities in (3.4) is contradicted, i.e.,

$$\lim_j \frac{H_{u_j}(x_j, t_j)}{H_{u_j}(z_j, t_j)} \in \{0, \infty\}.$$

By the convergence of v_j to v_∞ , we then deduce that

$$H_{v_\infty}(y_\infty, 1) = \lim_j H_{v_j}(t_j^{-1}(x_j - z_j), 1) = \lim_j \frac{H_{u_j}(x_j, t_j)}{H_{u_j}(z_j, t_j)}.$$

Thus, $H_{v_\infty}(y_\infty, 1) \in \{0, \infty\} \cap \mathbb{R} = \{0\}$. Given that v_∞ is analytical in $B_2 \setminus \{x_{n+1} = 0\}$, by unique continuation, we conclude that $v_\infty \equiv 0$ in B_2 , against the assumption

$$H_{v_\infty}(1) = \lim_j H_{v_j}(1) = 1.$$

In case the second set of inequalities in (3.4) is contradicted, i.e.,

$$\lim_j \frac{D_{u_j}(x_j, t_j)}{D_{u_j}(z_j, t_j)} \in \{0, \infty\}.$$

we have on, one hand that $D_{v_\infty}(1) = \lim_j D_{v_j}(1) = \lim_j I_{u_j}(z_j, t_j) \in [C, m_0]$, where C is the constant of the lower bound found in Lemma 3.2 and m_0 is the upper bound for the $m(z_j)$. On the other hand, we have that

$$D_{v_\infty}(y_\infty, 1) = \lim_j D_{v_j}(t_j^{-1}(x_j - z_j), 1) = \lim_j t_j \frac{D_{u_j}(x_j, t_j)}{H_{u_j}(z_j, t_j)} = \lim_j I_{u_j}(z_j, t_j) \frac{D_{u_j}(x_j, t_j)}{D_{u_j}(z_j, t_j)}.$$

Therefore, under the contradiction assumption, we infer that $D_{v_\infty}(y_\infty, 1) \in \{0, \infty\} \cap \mathbb{R} = \{0\}$, and taking into account the analyticity of the solutions, the last equality implies $v_\infty \equiv 0$, which is a contradiction.

Finally, as a byproduct of the first estimate in (3.4), the frequency function is well defined for $t \in [r, \varrho]$, provided that $x \in B'_{r/2}(x_0)$. Moreover, (3.5) follows straightforwardly from (3.4):

$$\left| I_u(x_0, t) - I_u(x, t) \right| = \left| I_u(x_0, t) \left(1 - \frac{D_u(x, t)}{D_u(x_0, t)} \cdot \frac{H_u(x_0, t)}{H_u(x, t)} \right) \right| \leq C. \quad \blacksquare$$

As an immediate consequence, the doubling estimates hold not only for the points on the free boundary but also for nearby points at suitable scales. This information will be crucial to bound error terms in the almost monotonicity formulas in the sequel (cf. (3.12) and (3.13)).

Corollary 3.6. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, then for every $r \in (0, \varrho]$, $x \in B'_{r/2}(x_0)$, and $t \in [r, \varrho]$*

$$\frac{H_u(x, 2t)}{H_u(x, t)}, \frac{D_u(x, 2t)}{D_u(x, t)} \leq [C^{-1}, C]. \quad (3.6)$$

Proof. The proof follows straightforwardly from Proposition 3.3 and Lemma 3.4 once the constants are chosen in such a way to apply the above-mentioned results. \blacksquare

3.2. Almost monotonicity of the frequency

By means of the doubling estimates established in Proposition 3.3, and arguing similarly to [13, Theorem 2.2], we can conclude almost monotonicity of the frequency $I_u(x_0, \cdot)$ under the condition

$$\mathbb{A}(x_0) = \text{Id}.$$

In fact, we establish quasi-monotonicity for all points x_0 and radii $r \geq (\alpha(x_0))^{\frac{1}{\alpha}}$; recall that $\alpha = 1 - \frac{n+1}{p}$, provided that

$$\alpha(x_0) := |\mathbb{A}(x_0) - \text{Id}|.$$

Notice that $\alpha(x_0) = 0$ if and only if $\mathbb{A}(x_0) = \text{Id}$. We start off proving a couple of preliminary results. In what follows, differentiation with respect to the radius will be denoted by a prime, and we write

$$\epsilon_D(x, t) := G_u(x, t) - D_u(x, t), \tag{3.7}$$

$$\epsilon_{D'}(x, t) := tD'_u(x, t) - (n - 1)D_u(x, t) - 2tE_u(x, t). \tag{3.8}$$

Lemma 3.7. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$, depending on m_0 with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, then for every $r \in (0, \varrho]$, $x \in B'_{r/2}(x_0)$, and $t \in [r, \varrho]$*

$$|\epsilon_D(x, t)| \leq C([\mathbb{A}]_{0,\alpha}(|x - x_0| + t)^\alpha + \alpha(x_0)) \cdot (D_u(x, t) + t^{-1/2}H_u^{1/2}(x, t)D_u^{1/2}(x, t)) \tag{3.9}$$

and

$$|\epsilon_{D'}(x, t)| \leq C([\mathbb{A}]_{0,\alpha}(|x - x_0| + t)^\alpha + \alpha(x_0))D_u(x, t). \tag{3.10}$$

Proof. Without loss of generality, we suppose $x_0 = \underline{0}$. Fix a point $x \in B'_{1/2}$, and set $\mathbb{B}(z) := \mathbb{A}(z) - \text{Id}$ for every $z \in B'_1$. To infer (3.9) and (3.10), we consider the equation satisfied by u and test it with a suitable function.

We start noticing that

$$\begin{aligned} \epsilon_D(x, t) &= G_u(x, t) - D_u(x, t) \\ &= - \int \nabla u(z) \cdot \nabla \left(u(z) \phi \left(\frac{|z - x|}{t} \right) \right) dz \\ &= \int \mathbb{B}(z) \nabla u(z) \cdot \nabla \left(u(z) \phi \left(\frac{|z - x|}{t} \right) \right) dz, \end{aligned} \tag{3.11}$$

where in the last equality we use that u is a solution to (2.2) (tested with $u(z)\phi(\frac{|z-x|}{t})$) and Signorini ambiguous boundary conditions. In order to prove (3.9), we use (3.11): for

ϱ sufficiently small, by the Hölder inequality we get

$$\begin{aligned} |\epsilon_{\mathcal{D}}(x, t)| &\leq ([\mathbb{A}]_{0,\alpha}(|x| + t)^\alpha + \alpha(x_0)) \\ &\quad \cdot \left(D_u(x, t) - t^{-1} \int_{B_t(x) \setminus B_{t/2}(x)} \phi' \left(\frac{|z-x|}{t} \right) |u(z)| |\nabla u(z)| \, dz \right) \\ &\leq C([\mathbb{A}]_{0,\alpha}(|x| + t)^\alpha + \alpha(x_0)) (D_u(x, t) + t^{-1/2} H_u^{1/2}(x, t) D_u^{1/2}(x, 2t)), \end{aligned} \tag{3.12}$$

with $C = C(\|\phi'\|_\infty)$. We conclude the estimate in (3.9) in view of Corollary 3.6 (cf. (3.6)).

To prove (3.10), we argue similarly and test the equation with the function w defined as the even extension across B'_1 of the restriction to B_1^+ of $\phi\left(\frac{|z-x|}{t}\right)\nabla u(z) \cdot (z-x)$. Note that w is an admissible test in view of the $H_{\text{loc}}^2(B_1^\pm \cup B'_1)$ regularity of u . Additionally, (H3) and Signorini’s ambiguous boundary conditions imply that

$$(\mathbb{A}(z)\nabla u(z) \cdot e_{n+1})\nabla u(z) \cdot (z-x) = a_{n+1,n+1}(z)\partial_{n+1}u(z) = 0 \quad \text{on } B'_1.$$

In view of this, we compute explicitly using the divergence theorem:

$$\begin{aligned} &\int \mathbb{B}(z)\nabla u(z) \cdot \nabla w \, dz && (3.13) \\ &= \int \nabla u(z) \cdot \nabla w \, dz \\ &= D_u(x, t) + \frac{1}{2} \int \phi \left(\frac{|z-x|}{t} \right) \nabla(|\nabla u(z)|^2) \cdot (z-x) \, dz \\ &\quad + \frac{1}{t} \int \phi' \left(\frac{|z-x|}{t} \right) \frac{(\nabla u(z) \cdot (z-x))^2}{|z-x|} \, dz \\ &= -\frac{n-1}{2} D_u(x, t) + \frac{1}{t} \int \phi' \left(\frac{|z-x|}{t} \right) \left(-\frac{|\nabla u(z)|^2}{2} |z-x| + \frac{(\nabla u(z) \cdot (z-x))^2}{|z-x|} \right) \, dz \\ &= -\frac{n-1}{2} D_u(x, t) + \frac{t}{2} D'_u(x, t) - t E_u(x, t) = -\epsilon_{\mathcal{D}'}(x, t). \end{aligned}$$

To establish (3.10), we can then proceed similarly as above:

$$\begin{aligned} |\epsilon_{\mathcal{D}'}(x, t)| &\leq C([\mathbb{A}]_{0,\alpha}(|x| + t)^\alpha + \alpha(x_0)) \\ &\quad \cdot \left(D_u(x, t) + t \int \phi \left(\frac{|z-x|}{t} \right) |\nabla^2 u(z)| |\nabla u(z)| \, dz + \int_{B_t(x) \setminus B_{t/2}(x)} |\nabla u(z)|^2 \, dz \right) \\ &\leq C([\mathbb{A}]_{0,\alpha}(|x| + t)^\alpha + \alpha(x_0)) (D_u(x, t) + D_u^{1/2}(x, t) D_u^{1/2}(x, 2t) + D_u(x, 2t)), \end{aligned}$$

where we used the $H_{\text{loc}}^2(B_1^\pm)$ regularity estimates of u . The conclusion then follows from the doubling properties of $D_u(x, \cdot)$ (cf. Corollary 3.6). ■

We establish next a similar result for $H_u(x_0, \cdot)$ together with a quasi-monotonicity formula.

Lemma 3.8. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 and $[\mathbb{A}]_{0,\alpha}$ with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, if $(\alpha(x_0))^{1/\alpha} \leq \varrho$, then*

$$((\alpha(x_0))^{1/\alpha}, \varrho] \ni t \mapsto \frac{H_u(x_0, t)}{t^n} \cdot \exp(Ct^\alpha) \text{ is non-decreasing,} \tag{3.14}$$

and

$$((\alpha(x_0))^{1/\alpha}, \varrho] \ni t \mapsto \frac{H_u(x_0, t)}{t^{n+2m_0}} \cdot \exp(-Ct^\alpha) \text{ is non-increasing.} \tag{3.15}$$

In particular, for all $(\alpha(x_0))^{1/\alpha} \leq r \leq s \leq \varrho$,

$$\int_{B_s(x_0) \setminus B_r(x_0)} |u(x)|^2 \, dx \leq CsH_u(x_0, s). \tag{3.16}$$

Proof. First, note that by scaling and a direct differentiation we easily get

$$\begin{aligned} H'_u(x, t) &= \frac{n}{t} H_u(x, t) - 2t^{-1} \int \phi' \left(\frac{|z-x|}{t} \right) u(z) \nabla u(z) \cdot \frac{z-x}{|z-x|} \, dz \\ &= \frac{n}{t} H_u(x, t) + 2G_u(x, t). \end{aligned} \tag{3.17}$$

We employ equalities (3.7), (3.17) and estimate (3.9) (with $x = x_0$) to deduce that

$$\begin{aligned} &\left| \frac{d}{dt} \left(\ln \left(\frac{H_u(x_0, t)}{t^n} \right) + 2 \int_t^\varrho \frac{I_u(x_0, s)}{s} \, ds \right) \right| \\ &= \frac{2|\epsilon_D(x_0, t)|}{H_u(x_0, t)} \leq C([\mathbb{A}]_{0,\alpha} t^\alpha + \alpha(x_0)) t^{-1} (I_u(x_0, t) + I_u^{1/2}(x_0, t)) \leq Ct^{\alpha-1}, \end{aligned}$$

where $C = C([\mathbb{A}]_{0,\alpha}, m_0) > 0$ and we have used that $(\alpha(x_0))^{1/\alpha} < t < 1$. The conclusion in (3.14) then follows at once by direct integration. Similarly, using $I_u(x_0, s) \leq m_0$, we have

$$\frac{d}{dt} \ln \left(\frac{H_u(x_0, t)}{t^{n+2m_0}} \right) \geq \frac{d}{dt} \left(\ln \left(\frac{H_u(x_0, t)}{t^n} \right) + 2 \int_t^\varrho \frac{I_u(x_0, s)}{s} \, ds \right) \geq -Ct^{\alpha-1},$$

and (3.15) follows.

Finally, the proof of (3.16) is a simple consequence of Fubini theorem by taking advantage of (3.14). ■

Thanks to Lemmata 3.7 and 3.8, we are now ready to establish the quasi-monotonicity of the frequency function at free boundary points x_0 with $\alpha(x_0) = 0$.

Proposition 3.9. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 , $[\mathbb{A}]_{0,\alpha}$, and α with this property. If u is a solution to (2.1) under assumptions (H1)–(H3),*

for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, if $\max\{(\alpha(x_0))^{1/\alpha}, r\} \leq \varrho$ and $x \in B'_{r/2}(x_0)$, then

$$I'_u(x, t) = \frac{2t}{H_u^2(x, t)} (E_u(x, t)H_u(x, t) - G_u^2(x, t)) + R_u(x, t) \tag{3.18}$$

and

$$|R_u(x, t)| \leq Ct^{\alpha-1} I_u(x, t) \quad \forall t \in [\max\{(\alpha(x_0))^{1/\alpha}, r\}, \varrho]. \tag{3.19}$$

In particular, if $\alpha(x_0) = 0$, then for every $x \in B'_{r/2}(x_0)$, $r < \varrho$,

$$[r, \varrho] \ni t \mapsto e^{Ct^\alpha} I_u(x, t) \text{ is non-decreasing.} \tag{3.20}$$

Proof. Without loss of generality, we prove the result in case $x_0 = \underline{0}$. We take advantage of formulas (3.7)–(3.10) in Lemma 3.7 and (3.17) in Lemma 3.8 to deduce that

$$\begin{aligned} \frac{I'_u(x, t)}{I_u(x, t)} &= \frac{1}{t} + \frac{D'_u(x, t)}{D_u(x, t)} - \frac{H'_u(x, t)}{H_u(x, t)} \stackrel{(3.8)}{=} 2 \frac{E_u(x, t)}{D_u(x, t)} - 2 \frac{G_u(x, t)}{H_u(x, t)} + \frac{\epsilon_{D'}(x, t)}{tD_u(x, t)} \\ &= \frac{2}{D_u(x, t)H_u(x, t)} (E_u(x, t)H_u(x, t) - G_u(x, t)D_u(x, t)) + \frac{\epsilon_{D'}(x, t)}{tD_u(x, t)} \\ &\stackrel{(3.7)}{=} \frac{2}{D_u(x, t)H_u(x, t)} (E_u(x, t)H_u(x, t) - G_u^2(x, t)) \\ &\quad + \frac{\epsilon_{D'}(x, t)}{tD_u(x, t)} + 2 \frac{\epsilon_D(x, t)G_u(x, t)}{D_u(x, t)H_u(x, t)}. \end{aligned}$$

Thus, for $t \in (0, 1/2)$, we conclude equality (3.18), i.e.,

$$\begin{aligned} I'_u(x, t) - \frac{2t}{H_u^2(x, t)} (E_u(x, t)H_u(x, t) - G_u^2(x, t)) \\ &= \frac{\epsilon_{D'}(x, t)}{H_u(x, t)} + 2t \frac{G_u(x, t)}{H_u^2(x, t)} \epsilon_D(x, t) \\ &\stackrel{(3.7)}{=} \frac{\epsilon_{D'}(x, t)}{H_u(x, t)} + 2t \frac{\epsilon_D^2(x, t)}{H_u^2(x, t)} + 2I(x, t) \frac{\epsilon_D(x, t)}{H_u(x, t)} =: R_u(x, t). \end{aligned} \tag{3.21}$$

To estimate R_u , we note that if $t \geq \max\{(\alpha(x_0))^{1/\alpha}, r\}$ and $x \in B'_{r/2}(x_0)$, then

$$\left| \frac{\epsilon_{D'}(x, t)}{H_u(x, t)} \right| \stackrel{(3.10)}{\leq} Ct^{\alpha-1} I_u(x, t),$$

in turn implying that

$$\frac{t\epsilon_D^2(x, t)}{H_u^2(x, t)} + I_u(x, t) \frac{|\epsilon_D(x, t)|}{H_u(x, t)} \stackrel{(3.9)}{\leq} Ct^{\alpha-1} I_u(x, t),$$

with $C = C(m(x_0), [\mathbb{A}]_{0,\alpha})$, where we use the local uniform upper bound on $I_u(x, t)$ given by Lemma 3.4. Therefore, estimate (3.19) follows straightforwardly.

In particular, if $\alpha(x_0) = 0$, then (3.19) holds true for all $t \in [r, \varrho]$, and thus, (3.20) follows by direct integration of (3.18) by taking into account estimate (3.19) and Cauchy–Schwarz inequality that implies $E_u(x, t)H_u(x, t) - G_u^2(x, t) \geq 0$. ■

Remark 3.10. It is also convenient to highlight a different expression of the derivative of the frequency function for later purposes: from (3.21), we get

$$I'_u(x, t) = \frac{2t}{H_u^2(x, t)} (E_u(x, t)H_u(x, t) - D_u^2(x, t)) + \tilde{R}_u(x, t), \quad (3.22)$$

where

$$\tilde{R}_u(x, t) := \frac{\epsilon_{D'}(x, t)}{H_u(x, t)} - 2I_u(x, t) \frac{\epsilon_D(x, t)}{H_u(x, t)}. \quad (3.23)$$

In particular, if (H4) is satisfied in x_0 , $x \in B'_{r/2}(x_0)$ and $t \geq \max\{(\alpha(x_0))^{1/\alpha}, r\}$, then

$$|\tilde{R}_u(x, t)| \leq Ct^{\alpha-1} I_u(x, t). \quad (3.24)$$

An additive quasi-monotonicity formula is then easily deduced.

Corollary 3.11. *For every $m_0 > 0$, there exist constants $\varrho, C > 0$ depending on m_0 and $[\mathbb{A}]_{0,\alpha}$ with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, if $\max\{(\alpha(x_0))^{1/\alpha}, r\} < \varrho$ and $x \in B'_{r/2}(x_0)$, then*

$$[\max\{(\alpha(x_0))^{1/\alpha}, r\}, \varrho] \ni t \mapsto I_u(x, t) + Ct^\alpha \text{ is non-decreasing.} \quad (3.25)$$

In particular, if $\alpha(x_0) = 0$, then

$$I_u(x_0, 0^+) := \lim_{r \downarrow 0} I_u(x_0, r) \geq 3/2. \quad (3.26)$$

Proof. The proof of (3.25) is straightforward from inequality (3.18), estimate (3.19) in Proposition 3.9, and (3.5) in Lemma 3.4. Furthermore, if $\alpha(x_0) = 0$, then from [5, Lemma 1], Proposition 3.1, and Proposition 3.9, one deduces that $I_u(x_0, r) \geq \frac{3}{2}e^{-Cr^\alpha}$ for all $r \in (0, \varrho]$. Therefore, $I_u(x_0, 0^+) \geq 3/2$. ■

Remark 3.12. The $H_{\text{loc}}^2(B_1)$ regularity of a solution u has been exploited to infer the quasi-monotonicity property of the frequency function $I_u(x_0, \cdot)$ at points x_0 as in the statement of Corollary 3.11 in order to estimate the error term $\epsilon_{D'}$ (cf. Lemma 3.7). Different approaches, such as those in [23, 24], lead to quasi-monotonicity formulas holding in the less restrictive Hölder regularity scale for the matrix field. Despite this, in order to establish rectifiability of free boundary points with finite order of contact, we will crucially use the $W^{1,p}$ regularity of the matrix field as well as the already mentioned $H_{\text{loc}}^2(B_1)$ regularity of solutions (cf. Proposition 4.2).

4. Main estimates on the frequency

4.1. Oscillation estimate of the frequency

We introduce the following notation for the radial variations of the frequency at a point $x_0 \in \Gamma(u)$ with $m(x_0) < \infty$ and $\alpha(x_0) = 0$:

$$\Delta_r^x(x_0) := I_u(x_0, r) + Cr^\alpha - (I_u(x_0, \rho) + C\rho^\alpha), \quad 0 < \rho < r, \quad (4.1)$$

for $C > 0$ the constant in Corollary 3.11 so that $\Delta_\rho^r(x_0) \geq 0$ for r, ρ sufficiently small. The result in the ensuing Proposition 4.2 shows how the spatial oscillation of the frequency in two nearby points at a given scale is in turn controlled by the radial variations at comparable scales. We establish first a technical result. To this aim, it is convenient to define the parameter

$$\theta := \min\{[\mathbb{A}]_{0,\alpha}^{-1/\alpha}, 1\}. \tag{4.2}$$

Lemma 4.1. *For every $\mathfrak{m}_0 > 0$, there exist constants $\varrho, C > 0$ depending on \mathfrak{m}_0 and $[\mathbb{A}]_{0,\alpha}$ with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $\mathfrak{m}(x_0) \leq \mathfrak{m}_0$, and $\mathfrak{a}(x_0) = 0$, then, for all $r_1 \leq \varrho$, $r_0 \in (\frac{\theta}{16}r_1, r_1)$, and $x \in B'_{\frac{\theta}{32}r_1}(x_0)$, we have*

$$\int_{B_{r_1}(x) \setminus B_{r_0}(x)} (\nabla u(z) \cdot (z - x) - I_u(x, r_0)u(z))^2 \frac{1}{|z - x|} dz \leq CH_u(x, 2r_1)\Delta_{r_0}^{2r_1}(x).$$

Proof. By translation, it suffices to prove the lemma for $x_0 = \underline{0}$. We start off with the following computation that uses Remark 3.10:

$$\begin{aligned} & - \int \phi' \left(\frac{|z - x|}{t} \right) (\nabla u(z) \cdot (z - x) - I_u(x, t)u(z))^2 \frac{1}{|z - x|} dz \\ & = t^2 E_u(x, t) - 2t I_u(x, t)G_u(x, t) + I_u^2(x, t)H_u(x, t) \\ & = \frac{t^2}{H_u(x, t)} (E_u(x, t)H_u(x, t) - D_u^2(x, t)) - 2t \epsilon_D(x, t)I_u(x, t) \\ & \stackrel{(3.22)}{=} \frac{t}{2} H_u(x, t) (I_u'(x, t) - \bar{R}_u(x, t)), \end{aligned} \tag{4.3}$$

where

$$\bar{R}_u = \tilde{R}_u + \frac{4}{H_u(x, t)} \epsilon_D(x, t)I_u(x, t) = \frac{\epsilon_{D'}(x, t)}{H_u(x, t)} + 2I_u(x, t) \frac{\epsilon_D(x, t)}{H_u(x, t)}.$$

with \tilde{R}_u function in (3.23). In particular, the above equalities show that the last factor in (4.3) is nonnegative, being nonnegative the term on the first line of (4.3) itself. Therefore, we may use the elementary integral estimate

$$\int_{B_{r_1}(x) \setminus B_{r_0}(x)} f(z) dz \leq \frac{C}{r_0} \int_{r_0}^{2r_1} \int \phi' \left(\frac{|z - x|}{t} \right) f(z) dz dt \quad \text{for all } 0 < r_0 \leq r_1, \tag{4.4}$$

which holds true for any measurable function $f \geq 0$, in order to deduce

$$\begin{aligned} & \int_{B_{r_1}(x) \setminus B_{r_0}(x)} (\nabla u(z) \cdot (z - x) - I_u(x, r_0)u(z))^2 \frac{1}{|z - x|} dz \\ & \stackrel{(4.4)}{\leq} \frac{C}{r_0} \int_{r_0}^{2r_1} \int \phi' \left(\frac{|z - x|}{t} \right) (\nabla u(z) \cdot (z - x) - I_u(x, r_0)u(z))^2 \frac{1}{|z - x|} dz dt \end{aligned}$$

$$\begin{aligned}
 &\leq -\frac{C}{r_0} \int_{r_0}^{2r_1} \int \phi' \left(\frac{|z-x|}{t} \right) (\nabla u(z) \cdot (z-x) - I_u(x,t)u(z))^2 \frac{1}{|z-x|} dz dt \\
 &\quad - \frac{C}{r_0} \int_{r_0}^{2r_1} \int \phi' \left(\frac{|z-x|}{t} \right) (I_u(x,t) - I_u(x,r_0))^2 u^2(z) \frac{1}{|z-x|} dz dt \\
 &\stackrel{(4.3)}{\leq} \frac{C}{r_0} \int_{r_0}^{2r_1} \frac{t}{2} H_u(x,t) (I'_u(x,t) - \bar{R}_u(x,t)) dt \\
 &\quad + \frac{C}{r_0} ((I_u(x,2r_1) - I_u(x,r_0))^2 + (2r_1)^{2\alpha}) \int_{r_0}^{2r_1} H_u(x,t) dt, \tag{4.5}
 \end{aligned}$$

where in the last estimate we have used Corollary 3.11 because $x \in B'_{\frac{\theta}{32}r_1} \subset B'_{r_0/2}$.

From Lemma 3.4 and Lemma 3.8, we get that $H_u(x,t) \leq CH_u(x,2r_1)$ for all $t \in [r_0, 2r_1]$. Furthermore, we can use (3.24) in Remark 3.10 to estimate $|\bar{R}_u(x,t)|$ for all $x \in B_{\frac{\theta}{32}r_1}(x_0)$ and $t \in [r_0, 2r_1]$. Thus, by taking into account that $I'_u(x,t) - \bar{R}_u(x,t) \geq 0$ (cf. (4.3)), from (4.5), we get

$$\begin{aligned}
 &\int_{B_{r_1}(x) \setminus B_{r_0}(x)} (\nabla u(z) \cdot (z-x) - I_u(x,r_0)u(z))^2 \frac{1}{|z-x|} dz \\
 &\leq CH_u(x,2r_1) \int_{r_0}^{2r_1} (I'_u(x,t) - \bar{R}_u(x,t)) dt \\
 &\quad + CH_u(x,2r_1) ((I_u(x,2r_1) - I_u(x,r_0))^2 + (2r_1)^\alpha - r_0^\alpha) \\
 &\leq CH_u(x,2r_1) (I_u(x,2r_1) - I_u(x,r_0)) + CH_u(x,2r_1) ((2r_1)^\alpha - r_0^\alpha) \\
 &\leq CH_u(x,2r_1) \Delta_{r_0}^{2r_1}(x),
 \end{aligned}$$

where we used that $r_0 \in (\frac{\theta}{16}r_1, r_1)$ and $I_u(x,t) \leq C$ for all $t \in (r_0, 2r_1)$ and $x \in B'_{\frac{\theta}{32}r_1}$ by Lemma 3.4. ■

We are now ready to prove a spatial oscillation estimate on the frequency function in terms of the radial oscillation computed between suitable radii in all points belonging to a neighborhood of a point x_0 with $\alpha(x_0) = 0$.

Proposition 4.2. *For every $\mathfrak{m}_0 > 0$, there exist constants $\varrho, C > 0$ depending on \mathfrak{m}_0 and $[\mathbb{A}]_{0,\alpha}$ with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), then for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $\mathfrak{m}(x_0) \leq \mathfrak{m}_0$, and $\alpha(x_0) = 0$, for all $R, \rho > 0$ with $R > \frac{32}{\theta}$ and $R\rho < \varrho$, we have*

$$|I_u(x_1, R\rho) - I_u(x_2, R\rho)| \leq C ((\Delta_{R\rho/4}^{4R\rho}(x_1))^{1/2} + (\Delta_{R\rho/4}^{R\rho}(x_2))^{1/2} + (R\rho)^\alpha),$$

for every $x_1, x_2 \in B'_\rho(x_0)$ (where θ is defined in (4.2)).

Proof. Without loss of generality, we show the conclusion for $x_0 = \underline{0}$. Moreover, we set $t = R\rho$. Note that, under the assumption $R > \frac{32}{\theta}$, if $t < \varrho$ is sufficiently small, then $I_u(x,t) \leq C$ with the constant C depending on \mathfrak{m}_0 by Lemma 3.4.

The proof is based on estimating the tangential derivative of the frequency function $x \mapsto I_u(x, t)$ for $x \in B'_\rho$ by taking advantage of the H^2_{loc} regularity of u . Thus, we start off noticing that the functions $x \mapsto H_u(x, t)$ and $x \mapsto D_u(x, t)$ are differentiable and, for every $e \in \mathbb{R}^{n+1}$ with $e \cdot e_{n+1} = 0$, we have that

$$\partial_e H_u(x, t) = -2 \int \phi' \left(\frac{|z-x|}{t} \right) u(z) \partial_e u(z) \frac{1}{|z-x|} dz, \tag{4.6}$$

and setting $\mathbb{B}(z) := \mathbb{A}(z) - \mathbb{A}(\underline{0}) (= \mathbb{A}(z) - \text{Id})$,

$$\begin{aligned} \partial_e D_u(x, t) &= 2 \int \phi \left(\frac{|z-x|}{t} \right) \nabla u(z) \cdot \nabla(\partial_e u)(z) dz \\ &= 2 \int \phi \left(\frac{|z-x|}{t} \right) (\mathbb{A} - \mathbb{B})(z) \nabla u(z) \cdot \nabla(\partial_e u)(z) dz \\ &= -2t^{-1} \int \phi' \left(\frac{|z-x|}{t} \right) \partial_e u(z) \mathbb{A}(z) \nabla u(z) \cdot \frac{z-x}{|z-x|} dz \\ &\quad - 2 \int \phi \left(\frac{|z-x|}{t} \right) \mathbb{B}(z) \nabla u(z) \cdot \nabla(\partial_e u)(z) dz \\ &= -2t^{-1} \int \phi' \left(\frac{|z-x|}{t} \right) \partial_e u(z) \nabla u(z) \cdot \frac{z-x}{|z-x|} dz + \epsilon_{\partial_e D}(x, t), \end{aligned} \tag{4.7}$$

where

$$\begin{aligned} \epsilon_{\partial_e D}(x, t) &:= -2t^{-1} \int \phi' \left(\frac{|z-x|}{t} \right) \mathbb{B}(z) \partial_e u(z) \nabla u(z) \cdot \frac{z-x}{|z-x|} dz \\ &\quad - 2 \int \phi \left(\frac{|z-x|}{t} \right) \mathbb{B}(z) \nabla u(z) \cdot \nabla(\partial_e u)(z) dz. \end{aligned}$$

The third equality in (4.7) follows from the divergence theorem applied to the vector field $V(z) := \phi \left(\frac{|z-x|}{t} \right) \partial_e u(z) \mathbb{A}(z) \nabla u(z)$; note that $V \in C^\infty(B_t(x) \setminus B'_1, \mathbb{R}^{n+1})$, V has compact support, and the divergence of V does not concentrate on B'_1 . Recalling the H^2_{loc} -estimates in Theorem 2.1 and the doubling estimates in Corollary 3.6, we have

$$\begin{aligned} |\epsilon_{\partial_e D}(x, t)| &\leq C |e| t^{-1} (|x| + t)^\alpha D_u(x, 2t) + C (|x| + t)^\alpha D_u^{1/2}(x, t) \|u\|_{H^2(B_t(x))} \\ &\leq C |e| t^{-1} (|x| + t)^\alpha (D_u(x, 2t) + D_u^{1/2}(x, t) D_u^{1/2}(x, 2t)) \\ &\leq C |e| t^{-1} (|x| + t)^\alpha D_u(x, t). \end{aligned} \tag{4.8}$$

We choose $e := x_2 - x_1$ and set

$$\begin{aligned} \mathcal{E}_i(z) &:= \nabla u(z) \cdot (z - x_i) - I_u(x_i, t) u(z) \quad \text{for } i = 1, 2, \\ \Delta I &:= I_u(x_1, t) - I_u(x_2, t) \quad \text{and} \quad \Delta \mathcal{E}(z) := \mathcal{E}_1(z) - \mathcal{E}_2(z). \end{aligned}$$

Then, we have that $\partial_e u(z) = \Delta I u(z) + \Delta \mathcal{E}(z)$. Thus, from (4.6), we infer that

$$\partial_e H_u(x, t) = 2\Delta I \cdot H_u(x, t) - 2 \int \phi' \left(\frac{|z-x|}{t} \right) \Delta \mathcal{E}(z) \frac{u(z)}{|z-x|} dz,$$

while from (4.7) and (3.7), we conclude

$$\begin{aligned} \partial_e D_u(x, t) &= 2\Delta I \cdot (D_u(x, t) + \epsilon_D(x, t)) + \epsilon_{\partial_e D}(x, t) \\ &\quad - 2t^{-1} \int \phi' \left(\frac{|z-x|}{t} \right) \Delta \mathcal{E}(z) \nabla u(z) \cdot \frac{z-x}{|z-x|} dz. \end{aligned}$$

In particular, by a direct computation, we deduce that

$$\begin{aligned} &\partial_e I(x, t) \\ &= \frac{t}{H_u^2(x, t)} (H_u(x, t) \partial_e D_u(x, t) - D_u(x, t) \partial_e H_u(x, t)) \\ &= \frac{2}{H_u(x, t)} \int -\phi' \left(\frac{|z-x|}{t} \right) \Delta \mathcal{E}(z) (\nabla u(z) \cdot (z-x) - I_u(x, t) u(z)) \frac{1}{|z-x|} dz \\ &\quad + \frac{t}{H_u(x, t)} \cdot (2\Delta I \cdot \epsilon_D(x, t) + \epsilon_{\partial_e D}(x, t)). \end{aligned} \quad (4.9)$$

We estimate (4.9) (recall that $t = R\rho$ and $x \in B'_\rho$). First, notice that thanks to (3.9) we may conclude that

$$\frac{t}{H_u(x, t)} |2\Delta I \cdot \epsilon_D(x, t)| \leq C |\Delta I| t^\alpha (I_u(x, t) + I_u^{1/2}(x, t)) \leq C t^\alpha,$$

for some $C > 0$. Furthermore, by (4.8), we get that

$$\frac{t}{H_u(x, t)} |\epsilon_{\partial_e D}(x, t)| \leq C t^\alpha I_u(x, t) \leq C t^\alpha,$$

where we used that $|e| \leq 2\rho \leq t$. Note that since $x \in B'_\rho$, by elliptic regularity of u (cf. Theorem 2.1), we infer that

$$\begin{aligned} &\sup_{z \in B_{t+\rho}^+(x)} |\nabla u(z) \cdot (z-x) - I_u(x, t) u(z)| \\ &\leq t \sup_{z \in B_{t+\rho}^+(x)} |\nabla u(z)| + I_u(x, t) \|u\|_{C^0(B_{t+\rho})} \\ &\leq C t^{-\frac{n+1}{2}} \|u\|_{L^2(B_{2t+2\rho})} \leq C t^{-\frac{n}{2}} H_u^{1/2}(2t+2\rho), \end{aligned}$$

where we use (3.16) in Lemma 3.8. Hence, we have that

$$\begin{aligned} \partial_e I_u(x, t) &\leq C t^{-\frac{n}{2}} \frac{H_u^{1/2}(2t+2\rho)}{H_u(x, t)} \\ &\quad \cdot \int -\phi' \left(\frac{|z-x|}{t} \right) (|\mathcal{E}_1(z)| + |\mathcal{E}_2(z)|) \frac{1}{|z-x|} dz + C t^\alpha. \end{aligned} \quad (4.10)$$

In order to estimate the integral term in (4.10), we notice that

$$B_t(x) \setminus B_{t/2}(x) \subset B_{t+2\rho}(x_i) \setminus B_{t/2-2\rho}(x_i) \quad \forall x \in B'_{t/2}, \text{ for } i = 1, 2;$$

therefore,

$$\begin{aligned} \int_{B_t(x) \setminus B_{\frac{t}{2}}(x)} |\mathfrak{E}_i(z)| \frac{1}{|z-x|} dz &\leq \frac{2(t+2\rho)}{t} \int_{B_{t+2\rho}(x_i) \setminus B_{\frac{t}{2}-2\rho}(x_i)} |\mathfrak{E}_i(z)| \frac{1}{|z-x_i|} dz \\ &\leq C t^{\frac{n}{2}} \left(\int_{B_{t+2\rho}(x_i) \setminus B_{\frac{t}{2}-2\rho}(x_i)} \mathfrak{E}_i^2(z) \frac{1}{|z-x_i|} dz \right)^{1/2}, \end{aligned} \tag{4.11}$$

where we choose $R > 8$ and we use the direct computation

$$\int_{B_{t+2\rho}(x_i) \setminus B_{\frac{t}{2}-2\rho}(x_i)} \frac{1}{|z-x_i|} dz \leq C t^n,$$

with $C > 0$ a dimensional constant. If $R > \frac{32}{\theta}$, then we are in the position to apply Lemma 4.1 (with $r_0 = t/2 - 2\rho$ and $r_1 = t + 2\rho$) to get

$$\int_{B_{t+2\rho}(x_i) \setminus B_{t/2-2\rho}(x_i)} \mathfrak{E}_i^2(z) \frac{1}{|z-x_i|} dz \leq C_{4.1}(A) H_u(x_i, 2t+4\rho) \Delta_{\frac{t}{2}-2\rho}^{2(t+2\rho)}(x_i). \tag{4.12}$$

Using (4.10)–(4.12), we claim that for all $x \in B'_\rho$

$$\partial_e I_u(x, t) \leq C (\Delta_{t/4}^{4t}(x_1))^{1/2} + C (\Delta_{t/4}^{4t}(x_2))^{1/2} + C t^\alpha, \tag{4.13}$$

from which the conclusion follows by integrating (4.13) along the segment $\{x_1 + re : r \in [0, 1]\}$. Indeed, $4t \geq 2(t + 2\rho)$ and $\frac{t}{4} < \frac{t}{2} - 2\rho$, and the monotonicity of Corollary 3.11 in the set of radii under consideration. Moreover,

$$H_u^{1/2}(2t+2\rho) H_u^{-1}(x, t) H_u^{1/2}(x_i, 2t+4\rho) \leq H_u^{1/2}(2t+2\rho) H_u^{-1}(t) H_u^{1/2}(2t+4\rho) \leq C,$$

thanks to the estimates in Lemma 3.4 and Corollary 3.6 because $x_i \in B'_\rho$ and $t = R\rho \in (2\rho, \varrho)$. ■

4.2. Estimate of the mean flatness via the frequency function

We introduce the mean flatness.

Definition 4.3. Given a Radon measure μ in \mathbb{R}^{n+1} , for every $x_0 \in \mathbb{R}^n$ and for every $r > 0$, set

$$\beta_\mu(x, r) := \inf_{\mathcal{L}} \left(r^{-n-1} \int_{B_r(x)} \text{dist}^2(y, \mathcal{L}) d\mu(y) \right)^{1/2},$$

where the infimum is taken among all affine $(n - 1)$ -dimensional planes $\mathcal{L} \subset \mathbb{R}^{n+1}$, and $\text{dist}(y, \mathcal{L}) := \inf_{x \in \mathcal{L}} |y - x|$.

As shown in [16, 18], the mean flatness β_μ of an arbitrary measure μ supported on $\Gamma(u)$ is controlled in terms of the integration of suitable radial oscillations of the frequency with respect to μ .

Proposition 4.4. *For every $m_0 > 0$ and $R > \frac{64}{\theta}$ (where θ is defined in (4.2)), there exist constants $\varrho, C > 0$ depending on R, m_0 , and $[\mathbb{A}]_{0,\alpha}$ with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0 \in \Gamma(u) \cap B'_{1/2}$ satisfying (H4) with $m(x_0) \leq m_0$, and $\alpha(x_0) = 0$, then for every $r > 0$ with $Rr < \varrho$, for every finite Borel measure μ with $\text{spt } \mu \subseteq \Gamma(u)$, and for every $p \in \Gamma(u) \cap B'_r(x_0)$,*

$$\beta_\mu^2(p, r) \leq \frac{C}{r^{n-1}} \left(\int_{B_r(p)} \Delta_{(R-5)r/2}^{(2R+4)r}(x) \, d\mu(x) + (Rr)^\alpha \mu(B_r(p)) \right).$$

Proof. The proof is a simple adaptation of the ones in [16, Proposition 4.2] and [18, Proposition 5.1]. The condition $R > \frac{64}{\theta}$ is used in order to apply Lemma 4.1. We leave the details to the readers. ■

5. Intrinsic frequency

In this section, we introduce an elementary change of variables in order to make a generic free boundary point x_0 satisfy $\alpha(x_0) = 0$ for a different, related thin obstacle problem (cf. [20–22] for the thin obstacle, and [12] in the case of the classical obstacle problem). In such a way, we define an intrinsic frequency function for which the conclusions of Proposition 4.4 hold even without the matrix $\mathbb{A}(x_0)$ being the identity at free boundary points.

Given a solution u of (2.2) and $x_0 \in \Gamma(u) \cap B'_1$, consider the function

$$u_{\mathbb{A}(x_0)} : \Phi_{x_0}^{-1}(B_1) \rightarrow \mathbb{R}$$

defined by

$$u_{\mathbb{A}(x_0)}(x) := u(\Phi_{x_0}(x)),$$

where $\Phi_{x_0} : \mathbb{R}^{n+1} \rightarrow \mathbb{R}^{n+1}$ is the affine map $\Phi_{x_0}(x) := x_0 + \mathbb{A}^{1/2}(x_0)(x - x_0)$. In particular, changing variables by means of Φ_{x_0} leads to

$$\int_{B_1} \langle \mathbb{A}(x) \nabla u(x), \nabla u(x) \rangle \, dx = \det(\mathbb{A}^{1/2}(x_0)) \mathcal{E}_{\mathbb{A}(x_0)}(u_{\mathbb{A}(x_0)}, \Phi_{x_0}^{-1}(B_1)),$$

where

$$\mathcal{E}_{\mathbb{A}(x_0)}(v, U) := \int_U \langle \mathbb{C}_{x_0}(x) \nabla v(x), \nabla v(x) \rangle \, dx, \tag{5.1}$$

for every open set $U \subseteq \Phi_{x_0}^{-1}(B_1)$, and $\mathbb{C}_{x_0}(x) := \mathbb{A}^{-1/2}(x_0) \mathbb{A}(\Phi_{x_0}(x)) \mathbb{A}^{-1/2}(x_0)$. Note that $\mathbb{C}_{x_0}(x_0) = \text{Id}$. Therefore, $u_{\mathbb{A}(x_0)}$ turns out to be the solution to the thin obstacle problem for the energy in (5.1) among all functions in $v \in g(\Phi_{x_0}) + H_0^1(\Phi_{x_0}^{-1}(B_1))$ that are even across the corresponding hyperplane $\Phi_{x_0}^{-1}(\{x_{n+1} = 0\}) = \{x_{n+1} = 0\}$ (thanks to hypothesis (H3)), and such that $v|_{\Phi_{x_0}^{-1}(B'_1)} \geq 0$. Moreover, there is a bijection of the free boundaries: $\Gamma(u_{\mathbb{A}(x_0)}) = \Phi_{x_0}^{-1}(\Gamma(u))$.

Let u be a solution to (2.2), and let $x_0 \in B'_1$ and $r > 0$ be such that

$$\Phi_{x_0}(B_r(x_0) \cap \{x_{n+1} = 0\}) \subset B'_1.$$

Being $u_{\mathbb{A}(x_0)}$ solution to the thin obstacle problem corresponding to the matrix field $\mathbb{C}_{x_0}(\cdot)$ in $B_r(x_0)$, we consider the related frequency function

$$I_{u_{\mathbb{A}(x_0)}}(x, s) = \frac{sD_{u_{\mathbb{A}(x_0)}}(x, s)}{H_{u_{\mathbb{A}(x_0)}}(x, s)}, \quad x \in \{x_{n+1} = 0\} \cap B_r(x_0), \quad s < r - |x - x_0|. \quad (5.2)$$

In passing, note that if x_0 satisfies assumption $\alpha(x_0) = 0$, then u coincides with $u_{\mathbb{A}(x_0)}$, and correspondingly, $I_{u_{\mathbb{A}(x_0)}}$ coincides with I_u at all points in B'_1 and admissible radii. For later purposes, it is convenient to point out explicit formulas for the Dirichlet energy

$$D_{u_{\mathbb{A}(x_0)}}(x, s) = \int \phi\left(\frac{|\Phi_{x_0}^{-1}(y) - x|}{s}\right) \frac{\langle \mathbb{A}(x_0)\nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_0)} dy,$$

and for the “boundary” L^2 norm of u

$$H_{u_{\mathbb{A}(x_0)}}(x, s) = - \int \phi'\left(\frac{|\Phi_{x_0}^{-1}(y) - x|}{s}\right) \frac{u^2(y)}{|\Phi_{x_0}^{-1}(y) - x| \det \mathbb{A}^{1/2}(x_0)} dy. \quad (5.3)$$

We call $I_{u_{\mathbb{A}(x_0)}}(x_0, r)$ the intrinsic function at a free boundary point $x_0 \in \Gamma(u)$ and set

$$N_u(x_0, r) := I_{u_{\mathbb{A}(x_0)}}(x_0, r).$$

Having fixed a point x_0 of the free boundary with finite frequency, we compare the intrinsic frequency function and the (standard) Dirichlet based one at points of the free boundary with finite frequency close to x_0 . Thus, for every $\mathfrak{m}_0 > 0$, we set

$$\Gamma^{\mathfrak{m}_0}(u) := \{x \in \Gamma(u) \cap B'_{1/2} : \sup_{r \in (0, 1/2)} N_u(x, r) \leq \mathfrak{m}_0\}.$$

Proposition 5.1. *For every $\mathfrak{m}_0 > 0$, there exist constants $\varrho, C > 0$ depending on the ellipticity constant λ , \mathfrak{m}_0 , and $[\mathbb{A}]_{0,\alpha}$ with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every $x_0, x_1 \in \Gamma^{\mathfrak{m}_0}(u) \cap B'_{1/2}$, if $r \in (0, \varrho)$ and $|x_0 - x_1| < C^{-1}r$, then*

$$|I_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r) - N_u(x_1, r)| \leq C|x_0 - x_1|^{\alpha/2} N_u(x_1, r). \quad (5.4)$$

Proof. We start off noticing that by (H1)

$$|\Phi_{x_0}^{-1}(x_1) - x_0| = |\Phi_{x_0}^{-1}(x_1) - \Phi_{x_0}^{-1}(x_0)| \leq \lambda^{-1/2}|x_1 - x_0|,$$

where the square root of the ellipticity constant λ bounds from below the norm of $\mathbb{A}^{-1/2}(x_0)$. Therefore, $I_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r)$ is well defined provided that ϱ, C^{-1} are sufficiently small (cf. Lemma 3.4).

To prove the inequality in (5.4), it is convenient to recall formulas (5.2)–(5.3):

$$D_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r) = \int \phi \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y-x_1)|}{r} \right) \frac{\langle \mathbb{A}(x_0) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_0)} dy$$

and

$$H_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r) = - \int \phi' \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y-x_1)|}{r} \right) \frac{u^2(y)}{|\mathbb{A}^{-1/2}(x_0)(y-x_1)|} \frac{1}{\det \mathbb{A}^{1/2}(x_0)} dy.$$

To estimate the difference between the Dirichlet energies, we introduce the sets

$$\begin{aligned} U_r(x) &:= ((x + \mathbb{A}^{1/2}(x)B_r) \cup (x_1 + \mathbb{A}^{1/2}(x_0)B_r)) \\ &\quad \setminus ((x + \mathbb{A}^{1/2}(x)B_{r/2}) \cap (x_1 + \mathbb{A}^{1/2}(x_0)B_{r/2})) \end{aligned}$$

for all $x \in B'_{1/2}$. Then, we argue as follows:

$$\begin{aligned} &D_{u_{\mathbb{A}(x_1)}}(x_1, r) - D_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r) \\ &= \int \phi \left(\frac{|\mathbb{A}^{-1/2}(x_1)(y-x_1)|}{r} \right) \frac{\langle \mathbb{A}(x_1) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_1)} dy \\ &\quad - \int \phi \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y-x_1)|}{r} \right) \frac{\langle \mathbb{A}(x_0) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_0)} dy \\ &= \int \left(\phi \left(\frac{|\mathbb{A}^{-1/2}(x_1)(y-x_1)|}{r} \right) - \phi \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y-x_1)|}{r} \right) \right) \frac{\langle \mathbb{A}(x_1) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_1)} dy \\ &\quad + \int \phi \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y-x_1)|}{r} \right) \left(\frac{\langle \mathbb{A}(x_1) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_1)} - \frac{\langle \mathbb{A}(x_0) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_0)} \right) dy \\ &=: D^{(1)}(r) + D^{(2)}(r). \end{aligned}$$

Since $|y-x_1| \leq C(\Lambda)r$ for all $y \in U_r(x_1)$ and $\Phi_{x_1}^{-1}(U_r(x_1)) \subseteq B_{C(\lambda, \Lambda)r}(x_1)$, we deduce that

$$\begin{aligned} |D^{(1)}(r)| &\leq C[\phi]_{0,1} |\mathbb{A}^{-1/2}(x_0) - \mathbb{A}^{-1/2}(x_1)| \int_{U_r(x_1)} \frac{\langle \mathbb{A}(x_1) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_1)} dy \\ &\leq C |\mathbb{A}^{1/2}(x_0) - \mathbb{A}^{1/2}(x_1)| \int_{\Phi_{x_1}^{-1}(U_r(x_1))} |\nabla u_{\mathbb{A}(x_1)}(y)|^2 dy \\ &\stackrel{(H1)}{\leq} C |x_1 - x_0|^{\alpha/2} D_{u_{\mathbb{A}(x_1)}}(x_1, Cr), \end{aligned}$$

and, analogously,

$$\begin{aligned} |D^{(2)}(r)| &\leq \left| \text{Id} - \frac{\det \mathbb{A}^{1/2}(x_1)}{\det \mathbb{A}^{1/2}(x_0)} \mathbb{A}^{-1}(x_1) \mathbb{A}(x_0) \right| \int_{U_r(x_1)} \frac{\langle \mathbb{A}(x_1) \nabla u(y), \nabla u(y) \rangle}{\det \mathbb{A}^{1/2}(x_1)} dy \\ &\stackrel{(H1)}{\leq} C |x_1 - x_0|^{\alpha/2} D_{u_{\mathbb{A}(x_1)}}(x_1, Cr), \end{aligned}$$

for some constant $C = C(n, \lambda, \Lambda, [\phi]_{0,1}, [\mathbb{A}]_{0,\alpha}) > 0$. For ρ sufficiently small, we apply iteratively Corollary 3.6 to $u_{\mathbb{A}(x_1)}$ to conclude

$$|Du_{\mathbb{A}(x_1)}(x_1, r) - Du_{\mathbb{A}(x_0)}(\Phi_{x_0}^{-1}(x_1), r)| \leq C|x_1 - x_0|^{\alpha/2} Du_{\mathbb{A}(x_1)}(x_1, r). \tag{5.5}$$

To estimate the difference of the H -terms, we define $[0, \infty) \ni t \mapsto \psi(t) := \phi'(t)/t$ (recall that $\phi'(t) = 0$ for $t \in [0, 1/2] \cup [1, \infty)$) and notice that ψ is Lipschitz continuous on $[0, \infty)$, having assumed $\phi \in C^{1,1}$. We then argue as follows:

$$\begin{aligned} & H_{u_{\mathbb{A}(x_1)}}(x_1, r) - H_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r) \\ &= \int \psi \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y - x_1)|}{r} \right) \frac{u^2(y)}{r \det \mathbb{A}^{1/2}(x_0)} dy \\ &\quad - \int \psi \left(\frac{|\mathbb{A}^{-1/2}(x_1)(y - x_1)|}{r} \right) \frac{u^2(y)}{r \det \mathbb{A}^{1/2}(x_1)} dy \\ &= \int \left(\psi \left(\frac{|\mathbb{A}^{-1/2}(x_0)(y - x_1)|}{r} \right) - \psi \left(\frac{|\mathbb{A}^{-1/2}(x_1)(y - x_1)|}{r} \right) \right) \frac{u^2(y)}{r \det \mathbb{A}^{1/2}(x_0)} dy \\ &\quad + \int \psi \left(\frac{|\mathbb{A}^{-1/2}(x_1)(y - x_1)|}{r} \right) \left(\frac{1}{\det \mathbb{A}^{1/2}(x_0)} - \frac{1}{\det \mathbb{A}^{1/2}(x_1)} \right) \frac{u^2(y)}{r} dy \\ &=: H^{(1)}(r) + H^{(2)}(r). \end{aligned}$$

Therefore, we get straightforwardly that

$$|H^{(2)}(r)| \leq \left| \frac{\det \mathbb{A}^{1/2}(x_1)}{\det \mathbb{A}^{1/2}(x_0)} - 1 \right| H_{u_{\mathbb{A}(x_1)}}(x_1, r) \leq C|x_1 - x_0|^{\alpha/2} H_{u_{\mathbb{A}(x_1)}}(x_1, r),$$

with $C(n, \lambda, \Lambda, [\mathbb{A}]_{0,\alpha}) > 0$. To estimate $H^{(1)}$, we introduce the set

$$V_r(x_1) := (x_1 + \mathbb{A}^{1/2}(x_0)(B_r \setminus B_{r/2})) \cup (x_1 + \mathbb{A}^{1/2}(x_1)(B_r \setminus B_{r/2}))$$

and get

$$\begin{aligned} |H^{(1)}(r)| &\leq C|\mathbb{A}^{-1/2}(x_1) - \mathbb{A}^{-1/2}(x_0)| \int_{V_r(x_1)} \frac{u^2(y)}{r} dy \\ &\leq C|x_1 - x_0|^{\alpha/2} \int_{V_r(x_1)} \frac{u^2(y)}{r} dy \\ &\leq C|x_1 - x_0|^{\alpha/2} \int_{\Phi_{x_1}^{-1}(V_r(x_1))} \frac{u_{\mathbb{A}(x_1)}^2(z)}{r} dz, \end{aligned}$$

where $C(n, \lambda, \Lambda, [\phi]_{1,1}, [\mathbb{A}]_{0,\alpha}) > 0$. We have that

$$\Phi_{x_1}^{-1}(V_r(x_1)) \subseteq B_{(\lambda^{-1}\Lambda)^{1/2}r}(x_1) \setminus B_{\frac{1}{2}(\lambda\Lambda^{-1})^{1/2}r}(x_1),$$

and thus, we may estimate the r.h.s. above as follows:

$$|H^{(1)}(r)| \leq C|x_1 - x_0|^{\alpha/2} \int_{B_{(\lambda^{-1}\Lambda)^{1/2}r}(x_1) \setminus B_{\frac{1}{2}(\lambda\Lambda^{-1})^{1/2}r}(x_1)} \frac{u_{\mathbb{A}(x_1)}^2(z)}{r} dz.$$

With $u_{\mathbb{A}(x_1)}$ being a solution to a thin obstacle problem with $\alpha(x_1) = 0$, for ϱ sufficiently small, Lemma 3.8 yields

$$\int_{B_{(\lambda^{-1}\Lambda)^{1/2}r}(x_1) \setminus B_{\frac{1}{2}(\lambda\Lambda^{-1})^{1/2}r}(x_1)} u_{\mathbb{A}(x_1)}^2(z) \, dz \leq Cr H_{u_{\mathbb{A}(x_1)}}(x_1, (\lambda^{-1}\Lambda)^{1/2}r)$$

with $C = C([\mathbb{A}]_{0,\alpha}, \mathfrak{m}(x_1)) > 0$. In turn, the doubling properties of $H_{u_{\mathbb{A}(x_1)}}(x_1, \cdot)$ together with the quasi-monotonicity in (3.14) imply

$$|H^{(1)}(r)| \leq C|x_1 - x_0|^{\alpha/2} H_{u_{\mathbb{A}(x_1)}}(x_1, r).$$

Thus, we conclude that

$$|H_{u_{\mathbb{A}(x_1)}}(x_1, r) - H_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r)| \leq C|x_1 - x_0|^{\alpha/2} H_{u_{\mathbb{A}(x_1)}}(x_1, r), \tag{5.6}$$

and from estimates (5.5) and (5.6), we conclude that (always under the hypothesis that ϱ is sufficiently small)

$$|I_{u_{\mathbb{A}(x_0)}}(\Phi_{x_0}^{-1}(x_1), r) - I_{u_{\mathbb{A}(x_1)}}(x_1, r)| \leq \frac{2C|x_1 - x_0|^{\alpha/2}}{1 - C|x_1 - x_0|^{\alpha/2}} I_{u_{\mathbb{A}(x_1)}}(x_1, r). \quad \blacksquare$$

In view of the previous estimate on the intrinsic frequency function, we can rephrase the bound on the mean flatness in terms of the intrinsic frequency itself dispensing with the assumption $\alpha(x_0) = 0$ for the base point.

For points $x_0 \in \Gamma(u)$ with $\sup_{r \in (0, 1/2)} N_u(x_0, r) < \infty$, we set

$$\Xi_\rho^r(x_0) := N_u(x_0, r) + Cr^\alpha - (N_u(x_0, \rho) + C\rho^\alpha), \quad 0 < \rho < r,$$

for $C > 0$ the constant in (4.1) and r sufficiently small. Finally, we note that the seminorm $[(\det \mathbb{A}^{1/2}(x_0))\mathbb{C}_{x_0}]_{0,\alpha}$ is uniformly bounded; therefore, the constants θ in (4.2) for this new matrices are uniformly bounded with respect to x_0 in view of (H1) and (H2). We denote by $\theta_0 > 0$ its infimum.

Proposition 5.2. *For every $\mathfrak{m}_0 > 0$ and $R > 64/\theta_0$, there exist constants $\varrho, C > 0$ depending on R, \mathfrak{m}_0 , and $[\mathbb{A}]_{0,\alpha}$, with this property. If u is a solution to (2.1) under assumptions (H1)–(H3), for every finite Borel measure μ with $\text{spt } \mu \subseteq \Gamma^{\mathfrak{m}_0}(u)$, for every $x_0 \in \Gamma^{\mathfrak{m}_0}(u)$, then for every $r > 0$ with $Rr < \varrho$ and $p \in \Gamma^{\mathfrak{m}_0}(u) \cap B_r'(x_0)$*

$$\beta_\mu^2(p, r) \leq \frac{C}{r^{n-1}} \left(\int_{B_{R_2 r}(p)} \Xi_{R_1 r}^{R_2 r}(x) \, d\mu(x) + (R_2 r)^{\alpha/2} \mu(B_{R_2 r}(p)) \right),$$

for every $R_2 > \max\{2R^2, 2R + 4\}$ and $R_1 < \frac{1}{2}(R - 5)r$.

Proof. Set $\mu_{\mathbb{A}(x_0)} := (\Phi_{x_0}^{-1})_\# \mu$; then, $\text{spt}(\mu_{\mathbb{A}(x_0)}) \subseteq \Gamma(u_{\mathbb{A}(x_0)})$. Note that $\Phi_{x_0}^{-1}(B_r(p)) = x_0 + \mathbb{A}^{-1/2}(x_0)B_r(p - x_0) \subseteq B_{(1+2\lambda^{-1/2})r}(p)$. Thus, from the very definition of the mean

flatness β_μ , we infer that for all $p \in B'_r(x_0)$

$$\begin{aligned} \beta_\mu^2(p, r) &= \inf_{\mathcal{L}} r^{-n-1} \int_{B_r(p)} \text{dist}^2(y, \mathcal{L}) \, d(\Phi_{x_0})_{\#} \mu_{\mathbb{A}(x_0)}(y) \\ &= \inf_{\mathcal{L}} r^{-n-1} \int_{\Phi_{x_0}^{-1}(B_r(p))} \text{dist}^2(\Phi_{x_0}(y), \mathcal{L}) \, d\mu_{\mathbb{A}(x_0)}(y) \\ &= \inf_{\mathcal{L}} r^{-n-1} \int_{\Phi_{x_0}^{-1}(B_r(p))} \text{dist}^2(\Phi_{x_0}(y), \Phi_{x_0}(\mathcal{L})) \, d\mu_{\mathbb{A}(x_0)}(y) \\ &\leq \Lambda \inf_{\mathcal{L}} r^{-n-1} \int_{\Phi_{x_0}^{-1}(B_r(p))} \text{dist}^2(y, \mathcal{L}) \, d\mu_{\mathbb{A}(x_0)}(y) \\ &\leq CR^{n+1} \Lambda \beta_{\mu_{\mathbb{A}(x_0)}}^2(p, Rr), \end{aligned}$$

if $R \geq 1 + 2\lambda^{-1/2}$. Since $u_{\mathbb{A}(x_0)}$ satisfies the hypotheses of Proposition 4.4 in x_0 , recalling that $\mu_{\mathbb{A}(x_0)} = (\Phi_{x_0}^{-1})_{\#} \mu$ we deduce that

$$\begin{aligned} \beta_\mu^2(p, r) &\leq CR^{n+1} \Lambda \beta_{\mu_{\mathbb{A}(x_0)}}^2(p, Rr) \\ &\leq \frac{C}{r^{n-1}} \left(\int_{B_{Rr}(p)} (\Delta u_{\mathbb{A}(x_0)})_{\frac{1}{2}(R-5)r}^{(2R+4)r}(x) \, d\mu_{\mathbb{A}(x_0)}(x) + (Rr)^\alpha \mu_{\mathbb{A}(x_0)}(B_{Rr}(p)) \right) \\ &= \frac{C}{r^{n-1}} \int_{\Phi_{x_0}(B_{Rr}(p))} (\Delta u_{\mathbb{A}(x_0)})_{\frac{1}{2}(R-5)r}^{(2R+4)r}(\Phi_{x_0}^{-1}(x)) \, d\mu(x) \\ &\quad + \frac{C}{r^{n-1}} (Rr)^\alpha \mu(\Phi_{x_0}(B_{Rr}(p))), \end{aligned}$$

where we denote by $\Delta u_{\mathbb{A}(x_0)}$ the quantity defined in (4.1) by means of $I_{u_{\mathbb{A}(x_0)}}$, $R \geq (1 + 2\lambda^{-1/2}) \vee \frac{64}{\theta_0}$, and r is sufficiently small (cf. Proposition 4.4).

Eventually, Proposition 5.1 provides the conclusion as $\text{spt } \mu \subseteq \Gamma^{\text{mo}}(u)$, i.e.,

$$\beta_\mu^2(p, r) \leq \frac{C}{r^{n-1}} \left(\int_{B_{R_2 r}(p)} \Xi_{R_1 r}^{R_2 r}(x) \, d\mu(x) + (Rr)^{\alpha/2} \mu(B_{R_2 r}(p)) \right),$$

for $R_2 > \max\{2R^2, 2R + 4\}$ and $R_1 < \frac{1}{2}(R - 5)r$, and r is sufficiently small (cf. Proposition 5.1). ■

6. The measure and the structure of the free boundary

We recall the definition of homogeneous and almost homogeneous solutions to the (standard) thin obstacle problem:

$$\mathcal{H} := \left\{ w \in H_{\text{loc}}^1(\mathbb{R}^{n+1}) \setminus \{0\} : w(x) = |x|^\lambda w\left(\frac{x}{|x|}\right), \lambda \geq 3/2, \right. \\ \left. w|_{B_1} \text{ solves (2.1) with } \mathbb{A} \equiv \text{Id} \right\}.$$

Given a solution u to (2.1) with \mathbb{A} satisfying (H1)–(H3), we set

$$\Gamma^{\text{finite}}(u) := \{x \in \Gamma(u) : \limsup_{r \rightarrow 0^+} N_u(x, r) < +\infty\}.$$

Note that, for every $\mathfrak{m}_0 > 0$, we have that $\Gamma^{\mathfrak{m}_0}(u) \subseteq \Gamma^{\text{finite}}(u)$. For any point $x_0 \in \Gamma^{\text{finite}}(u)$, we set

$$J_u(x_0, t) := e^{Ct^\alpha} N_u(x_0, t),$$

for all $t > 0$ such that $J_u(x_0, t)$ is monotone, namely, for all $t \in (0, \varrho)$ with $\varrho > 0$ a constant depending on $[\mathbb{A}]_{0,\alpha}$ and \mathfrak{m}_0 as in the statement of Proposition 3.9.

Definition 6.1. Let $\eta > 0$, and let $u : B_1 \rightarrow \mathbb{R}$ be a solution to thin obstacle problem (2.1). Assume that $x_0 \in \Gamma^{\text{finite}}(u) \cap B'_{1/2}$ and $r \in (0, 1/2)$ is such that $J_u(x_0, r)$ is defined. Then, u is called η -almost homogeneous of (2.1) in $B_r(x_0)$ if

$$J_u(x_0, r/2) - J_u(x_0, r/4) \leq \eta.$$

The following lemma justifies this terminology.

Lemma 6.2. For every $\varepsilon > 0$ and $\mathfrak{m}_0 > 0$, there exist $\eta, \varrho > 0$ with the following property: if u is an η -almost homogeneous solution in $B_r(x_0)$ with $r \leq \varrho$ and $x_0 \in \Gamma^{\mathfrak{m}_0}(u) \cap B'_{1/2}$, then

$$\inf_{w \in \mathcal{H}} \|(u_{\mathbb{A}(x_0)})_{x_0, r} - w\|_{H^1(B_1)} \leq \varepsilon.$$

Proof. The proof follows by a contradiction argument similar to [16, Lemma 5.5]. Assume that for $\varepsilon > 0$ we could find sequences r_l of numbers and u_l of $\frac{1}{7}$ -almost homogeneous solutions in $B_{r_l}(x_l)$ such that

$$\inf_l \inf_{w \in \mathcal{H}} \|((u_l)_{\mathbb{A}(x_l)})_{x_l, r_l} - w\|_{H^1(B_1)} \geq \varepsilon, \tag{6.1}$$

with $x_l \in \Gamma^{\text{finite}}(u_l) \cap B'_{1/2}$ and $\mathfrak{m}(x_l) \leq \mathfrak{m}_0$. By Proposition 3.1, there exists a subsequence (not relabeled) of $v_l = ((u_l)_{\mathbb{A}(x_l)})_{x_l, r_l}$ converging to a solution v_∞ to the thin obstacle problem in B_1 for the standard Dirichlet energy. Moreover, we can assume that the points x_l converge to $x_\infty \in \overline{B}'_{1/2}$. From Proposition 3.3, we infer that

$$H_{v_\infty}(1/2) = \lim_l H_{v_l}(1/2) \geq C \lim_l H_{v_l}(1) > 0$$

so that v_∞ is not zero. On the other hand, we have that

$$I_{v_\infty}(1/2) - I_{v_\infty}(1/4) = \lim_l (I_{v_l}(1/2) - I_{v_l}(1/4)) = \lim_l (J_u(x_l, r_l/2) - J_u(x_l, r_l/4)) = 0.$$

This implies that v_∞ is a solution with constant frequency and thus is homogeneous (see, for instance, [16, Proposition 2.7]), contradicting (6.1). ■

A rigidity property of the type shown in [16, Proposition 5.6] holds in the case of non-smooth coefficients as well. We call spine $S(w)$ of a function $w \in \mathcal{H}$ the maximal subspace of invariance of u ,

$$S(w) := \{y \in \mathbb{R}^n \times \{0\} : w(x + y) = w(x) \quad \forall x \in \mathbb{R}^{n+1}\}.$$

We recall that the maximal dimension of the spine of a function w in \mathcal{H} is at most $n - 1$ (cf. [16, Section 5.2]), and we set \mathcal{H}^{top} for the set of homogeneous solutions w with $\dim S(w) = n - 1$; whereas $\mathcal{H}^{\text{low}} := \mathcal{H} \setminus \mathcal{H}^{\text{top}}$.

Proposition 6.3. *For every $\tau > 0$ and $m_0 > 0$, there exists $\eta, \varrho > 0$ with this property. If u is an η -almost homogeneous solution in $B_r(x_0)$, $r \leq \varrho$ and $x_0 \in \Gamma^{m_0}(u) \cap B'_{r/2}$ with $m(x_0) \leq m_0$, then the following dichotomy holds:*

- (i) *either for every point $x \in \Gamma^{m_0}(u) \cap B'_{r/2}(x_0)$, we have*

$$|J_u(x, r/2) - J_u(x_0, r/2)| \leq \tau,$$

- (ii) *or there exists a linear subspace $V \subset \mathbb{R}^n \times \{0\}$ of dimension $n - 2$ such that*

$$\begin{cases} y \in \Gamma^{m_0}(u) \cap B'_{r/2}(x_0), \\ J_u(y, r/8) - J_u(y, r/16) \leq \eta \end{cases} \implies \text{dist}(y, x_0 + V) \leq \tau r.$$

Proof. The proof proceeds by contradiction and follows the ideas in [16, Proposition 5.6]. Let $\tau > 0$ be a given constant and assume that there exist r_l and a sequence $(u_l)_{l \in \mathbb{N}}$ of $1/l$ -almost homogeneous solutions in B_{r_l} (this clearly holds up to horizontal translations) such that

- (i) *there exists $x_l \in \Gamma^{m_0}(u_l) \cap B'_{r_l/2}$ for which*

$$|J_u(x_l, r_l/2) - J_u(0, r_l/2)| > \tau, \tag{6.2}$$

- (ii) *for every linear subspace $V \in \mathbb{R}^n \times \{0\}$ of dimension $n - 2$, there exists $y_l \in \Gamma^{m_0}(u_l) \cap B'_{r_l/2}(x_0)$ (a priori depending on V) such that*

$$J_u(y_l, r_l/8) - J_u(y_l, r_l/16) \leq 1/l \quad \text{and} \quad \text{dist}(y_l, V) > \tau r_l. \tag{6.3}$$

We consider the rescaled functions $v_l := (u_l)_{0, r_l}$. By the compactness result in Proposition 3.1, v_l converge, up to a subsequence, to a nonzero solution to the thin obstacle problem with constant coefficients v_∞ . In particular, $v_\infty \in \mathcal{H}$ thanks to Lemma 6.2.

If $v_\infty \in \mathcal{H}^{\text{top}}$, then (6.2) is contradicted. Indeed, up to choosing a further subsequence, we can assume that $z_l := r_l^{-1}x_l \rightarrow z_\infty \in \bar{B}_{1/2}$; moreover, z_∞ is a critical point for v_∞ , because both $v_l(z_l) = |\nabla v_l(z_l)| = 0$ and the convergence is C^1 , and by a simple change of variables, we have that

$$|I_{v_\infty}(z_\infty, 1/2) - I_{v_\infty}(0, 1/2)| = \lim_{l \rightarrow \infty} |J_u(x_l, r_l/2) - J_u(0, r_l/2)| \geq \tau,$$

which is a contradiction to the constancy of the frequency at critical points of homogeneous solutions $v_\infty \in \mathcal{H}^{\text{top}}$ (see [16, Lemma 5.3]).

On the other hand, if $v_\infty \in \mathcal{H}^{\text{low}}$, we show a contradiction to the second condition in (6.3) with V any $(n - 2)$ -dimensional subspace containing $S(v_\infty)$. Indeed, let y_l be as in (6.3) for such a choice of V . By compactness, up to passing to a subsequence (not relabeled), $z_l := r_l^{-1}y_l \rightarrow z_\infty$ for some $z_\infty \in \bar{B}_{1/2}$ with $\text{dist}(z_\infty, V) \geq \tau > 0$. Arguing as before, we obtain

$$|I_{v_\infty}(z_\infty, 1/8) - I_{v_\infty}(z_\infty, 1/16)| = \lim_{l \rightarrow \infty} |J_u(y_l, r_l/8) - J_u(y_l, r_l/16)| \leq \lim_{l \rightarrow \infty} 1/l = 0,$$

where the last inequality is given by the $1/l$ -almost homogeneity of the functions u_l (cf. the first condition in (6.3)). Using [16, Proposition 2.7, Lemma 5.2], it follows that $z_\infty \in S(v_\infty)$, from which we infer a contradiction as $z_\infty \in S(v_\infty) \subseteq V$ and $\text{dist}(z_\infty, V) \geq \tau$. ■

Proof of Theorem 1.1

The proof is now a simple consequence of the results established in the previous sections. Indeed, we can follow verbatim [16, Section 6] (see also [18, Section 5.3]). Recall that

$$\Gamma^{\text{finite}}(u) := \{x \in B'_1 : \limsup_{r \rightarrow 0^+} N_u(x, r) < +\infty\}.$$

By a simple rescaling argument, if $x_0 \in B'_1$ and $r < \text{dist}(x_0, \partial B_1)$, then the function $u_r(y) := u(x_0 + ry)$ solves a thin obstacle problem (1.1) with \mathbb{A} satisfying (H1)–(H3) and

$$\Gamma^{\text{finite}}(u_r) \cap B'_{\frac{1}{2}} = \bigcup_{m_0 \geq \frac{3}{2}} \Gamma^{m_0}(u_r).$$

Therefore, it is enough to show that $\Gamma^{m_0}(u) \cap B'_{1/2}$ is rectifiable. To this aim, we fix $\rho_0 > 0$ such that the conclusions of all propositions in the previous sections hold for points $x \in \Gamma^{m_0}(u) \cap B'_{1/2}$ and radii $\rho \leq \rho_0$. We can then follow the proof of [16, Section 6] applied to the intrinsic frequency N_u starting at ρ_0 : indeed, the proof uses only the lower bound of the frequency (cf. Corollary 3.6), the estimate of the spatial oscillation of the frequency in terms of the mean flatness (cf. Proposition 5.2) and the rigidity of Proposition 6.3, together with the Reifenberg-type rectifiability criteria provided in the work by Naber and Valtorta [31].

Finally, we note that the proof of the rectifiability also gives the local finiteness of the measure of each $\Gamma^{m_0}(u)$, which we will use for the proof of Theorem 1.2 in the next section.

7. Finiteness of the frequency for $\mathbb{A} \in W^{1,\infty}$

In this section, we prove the finiteness of the intrinsic frequency N_u at all free boundary points for a solution u of (2.2), assuming that the matrix field \mathbb{A} satisfies (H1) with $p = \infty$,

(H2), and (H3) (see also [22]). Given this for granted, Theorem 1.2 is then an immediate consequence of Theorem 1.1.

We first establish several auxiliary results under the simplifying assumptions that the base point is the origin and that $\mathbb{A}(\underline{0}) = \text{Id}$ in the spirit of [12, Section 3.2]. Consider then the function

$$\mu : B_1 \rightarrow [0, \infty)$$

defined by

$$\mu(x) := \langle \mathbb{A}(x)v(x), v(x) \rangle \quad \text{if } x \neq 0 \text{ and } \mu(\underline{0}) = 1,$$

where $v(x) = \frac{x}{|x|}$. Recalling that \mathbb{A} is Lipschitz-continuous, we infer that $\mu \in C^{0,1}(B_1)$, and

$$\lambda \leq \mu(x) \leq \Lambda, \quad \text{for every } x \in B_1, \tag{7.1}$$

where λ, Λ are the ellipticity constants in (H2) (for a proof, see [12, Lemma 3.10]).

Here, for the sake of simplicity, we follow the computations in [12] which use Almgren original frequency function (cf. [2]) tailored for Lipschitz coefficients. Let us define the functions

$$\mathcal{E}_u(r) := \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle dx \quad \text{and} \quad \mathcal{H}_u(r) := \int_{\partial B_r} \mu u^2 d\mathcal{H}^n,$$

and the energy-driven frequency function

$$\mathcal{J}_u(r) := \frac{r \mathcal{E}_u(r)}{\mathcal{H}_u(r)}.$$

It is useful for the sequel to observe that

$$\mathcal{E}_u(r) = \int_{\partial B_r} u \langle \mathbb{A} \nabla u, \nu \rangle d\mathcal{H}^n. \tag{7.2}$$

This equality follows by computing the divergence of the vector field $u\mathbb{A}\nabla u$, by taking into account (2.2), and by exploiting the Signorini’s ambiguous conditions together with (H3).

In order to establish the monotonicity of \mathcal{J}_u , we start with the following lemma.

Lemma 7.1. *Let \mathbb{A} satisfy (H1) with $p = \infty$, (H2), and $\mathbb{A}(\underline{0}) = \text{Id}$, and let μ be as above. Then, there exists a constant $C > 0$ depending on n and on $[\mathbb{A}]_{0,1}$ such that, for every $r \in (0, 1)$ and $x \in B_r$, we have that*

$$|\text{Tr } \mathbb{A}(x) - (n + 1)\mu(x)| \leq Cr.$$

Proof. Fixing a point $x \in B_r$, let $\{\lambda_i\}_{i=1}^{n+1}$ be the eigenvalues of the matrix $\mathbb{A}(x)$ and $\{e_i\}_{i=1}^{n+1}$ be the corresponding orthonormal base of eigenvectors. Set $y_i := r e_i$; then,

$$\begin{aligned} & |\operatorname{Tr} \mathbb{A}(x) - (n+1)\mu(x)| \\ &= \left| \sum_{i=1}^{n+1} (\lambda_i - \mu(x)) \right| \\ &= \left| \sum_{i=1}^{n+1} (\langle \mathbb{A}(x)e_i, e_i \rangle - \langle \mathbb{A}(x)v(x), v(x) \rangle) \right| \\ &\leq \sum_{i=1}^{n+1} (|\langle \mathbb{A}(x)e_i, e_i \rangle - \langle \mathbb{A}(y_i)e_i, e_i \rangle| + |\langle \mathbb{A}(y_i)e_i, e_i \rangle - \langle \mathbb{A}(x)v(x), v(x) \rangle|) \\ &= \sum_{i=1}^{n+1} (|\langle (\mathbb{A}(x) - \mathbb{A}(y_i))e_i, e_i \rangle| + |\mu(y_i) - \mu(x)|) \\ &\leq C \sum_{i=1}^{n+1} (|e_i|^2 + 1)|x - y_i| \leq Cr, \end{aligned}$$

where we used the Lipschitz continuity of \mathbb{A} and μ , and that $x, y_i \in \bar{B}_r$. ■

Remark 7.2. From Lemma 7.1, we deduce that

$$\operatorname{Tr} \mathbb{A}(x) - (n+1)\mu(x) \geq -Cr,$$

in turn implying that for every $x \in B_1$

$$\mu^{-1}(x) \operatorname{Tr} \mathbb{A}(x) \geq -Cr\mu^{-1}(x) + (n+1). \tag{7.3}$$

First, we compute the derivative of \mathcal{E}_u .

Proposition 7.3. *Let u be a solution to (2.1) under assumptions (H1)–(H3) with*

$$p = \infty \quad \text{and} \quad \mathbb{A}(\underline{0}) = \operatorname{Id},$$

and let μ be as above. Then, there exists a constant $C > 0$ depending on n, λ, Λ , and $[\mathbb{A}]_{0,1}$ such that for \mathcal{L}^1 a.e. $r \in (0, 1)$

$$\mathcal{E}'_u(r) = 2 \int_{\partial B_r} \mu^{-1} \langle \mathbb{A}v, \nabla u \rangle^2 d\mathcal{H}^n + E_r,$$

with

$$E_r \geq -C \mathcal{E}_u(r) + \frac{n-1}{r} \mathcal{E}_u(r).$$

Proof. By the coarea formula and [12, Lemma 3.4] applied to the Lipschitz vector field $\mathbf{F}(x) := \frac{\mathbb{A}(x)x}{r\mu(x)}$, we have

$$\begin{aligned} \mathcal{E}'_u(r) &= \int_{\partial B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle d\mathcal{H}^n \\ &= 2 \int_{\partial B_r} \mu^{-1} \langle \mathbb{A} \nu, \nabla u \rangle^2 d\mathcal{H}^n + \frac{1}{r} \int_{B_r} \mu^{-1} \nabla \mathbb{A} : \mathbb{A} x \otimes \nabla u \otimes \nabla u \, dx \\ &\quad + \frac{1}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \operatorname{div}(\mu^{-1} \mathbb{A} x) \, dx - \frac{2}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla^T(\mu^{-1} \mathbb{A} x) \nabla u \rangle \, dx \\ &=: 2 \int_{\partial B_r} \mu^{-1} \langle \mathbb{A} \nu, \nabla u \rangle^2 d\mathcal{H}^n + R_1 + R_2 + R_3. \end{aligned} \tag{7.4}$$

We now estimate the R_i 's. We start with R_1 . By using the Lipschitz continuity of \mathbb{A} , (7.1), and (H2), we get

$$\begin{aligned} |R_1| &\leq \frac{1}{r} \int_{B_r} \sum_{i,j,k,l} |\mu^{-1} \partial_i a_{j,l} a_{i,k} x_k \partial_j u \partial_l u| \, dx \\ &\leq \lambda^{-1} \int_{B_r} \sum_{i,j,k,l} |\partial_i a_{j,l}| |a_{i,k} \partial_j u \partial_l u| \, dx \\ &\leq C \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \, dx = C \mathcal{E}_u(r). \end{aligned} \tag{7.5}$$

By computing explicitly the divergence, R_2 rewritten as

$$\begin{aligned} R_2 &= \frac{1}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \sum_{i,j}^{n+1} \partial_i(\mu^{-1} a_{ij} x_j) \, dx \\ &= \frac{1}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \sum_{i,j}^{n+1} \partial_i(\mu^{-1} a_{ij}) x_j \, dx + \frac{1}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \mu^{-1} \operatorname{Tr} \mathbb{A} \, dx \\ &\geq -C \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \, dx + \frac{1}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \mu^{-1} \operatorname{Tr} \mathbb{A} \, dx \\ &\geq -C \mathcal{E}_u(r) + \frac{n+1}{r} \mathcal{E}_u(r), \end{aligned} \tag{7.6}$$

where we used the Lipschitz continuity of $\mu^{-1} \mathbb{A}$, (7.1), and (7.3). Analogously, for R_3 , we have

$$\begin{aligned} R_3 &= -\frac{2}{r} \int_{B_r} \langle \mathbb{A} \nabla u, (\nabla^T(\mu^{-1} \mathbb{A})x) \nabla u \rangle \, dx - \frac{2}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \mu^{-1} \mathbb{A} \nabla u \rangle \, dx \\ &\geq -C \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \, dx - \frac{2}{r} \int_{B_r} \langle \mathbb{A} \nabla u, (\mu^{-1} \mathbb{A} - \operatorname{Id}) \nabla u \rangle - \frac{2}{r} \int_{B_r} \langle \mathbb{A} \nabla u, \nabla u \rangle \, dx \\ &\geq -C \mathcal{E}_u(r) - \frac{2}{r} \mathcal{E}_u(r), \end{aligned} \tag{7.7}$$

where we used the Lipschitz continuity of $\mu^{-1}\mathbb{A}$, (H2), (7.1), and $\mu^{-1}(\underline{0})\mathbb{A}(\underline{0}) = \text{Id}$. Collecting (7.4)–(7.7), we conclude. ■

We now focus on the derivative of \mathcal{H}_u .

Proposition 7.4. *Let u be a solution to (2.1) under assumptions (H1)–(H3) with $p = \infty$, and $\mathbb{A}(\underline{0}) = \text{Id}$, and let μ be as above. Then, there exists a constant $C > 0$ depending on n, λ, Λ , and $[\mathbb{A}]_{0,1}$ such that for \mathcal{L}^1 a.e. $r \in (0, 1)$*

$$\mathcal{H}'_u(r) = \frac{n}{r}\mathcal{H}_u(r) + 2 \int_{\partial B_r} u \langle \mathbb{A} \nabla u, \nu \rangle \, d\mathcal{H}^n + H_r,$$

with

$$|H_r| \leq C\mathcal{H}_u(r).$$

Proof. First, note that by the definition of μ, ν , the divergence theorem implies

$$\mathcal{H}_u(r) = \frac{1}{r} \int_{B_r} \text{div}(u^2 \mathbb{A}x) \, dx.$$

Thus, the coarea formula and Lemma 7.1 yield for \mathcal{L}^1 a.e. $r \in (0, 1)$

$$\begin{aligned} \mathcal{H}'_u(r) &= -\frac{1}{r}\mathcal{H}_u(r) + \frac{1}{r} \int_{\partial B_r} \text{div}(u^2 \mathbb{A}x) \, d\mathcal{H}^n \\ &= -\frac{1}{r}\mathcal{H}_u(r) + \frac{2}{r} \int_{\partial B_r} u \langle \mathbb{A}x, \nabla u \rangle \, d\mathcal{H}^n + \frac{1}{r} \int_{\partial B_r} u^2 \left(\sum_{i,j}^{n+1} \partial_i a_{i,j} x_j + \text{Tr } \mathbb{A} \right) \, d\mathcal{H}^n \\ &= \frac{n}{r}\mathcal{H}_u + 2 \int_{\partial B_r} u \langle \mathbb{A} \nu, \nabla u \rangle \, d\mathcal{H}^n + \frac{1}{r} \int_{\partial B_r} u^2 \sum_{i,j}^{n+1} \partial_i a_{i,j} x_j \, d\mathcal{H}^n \\ &\quad + \frac{1}{r} \int_{\partial B_r} u^2 (\text{Tr } \mathbb{A} - (n+1)\mu) \, d\mathcal{H}^n. \end{aligned}$$

We now estimate the last two summands. Thanks to the Lipschitz continuity of \mathbb{A} and (7.1), we have

$$\left| \frac{1}{r} \int_{\partial B_r} u^2 \sum_{i,j}^{n+1} \partial_i a_{i,j} x_j \, d\mathcal{H}^n \right| \leq C \int_{\partial B_r} u^2 \, d\mathcal{H}^n \leq C\mathcal{H}_u(r).$$

Moreover, using Lemma 7.1, we have that

$$\left| \frac{1}{r} \int_{\partial B_r} u^2 (\text{Tr } \mathbb{A} - (n+1)\mu) \, d\mathcal{H}^n \right| \leq C \int_{\partial B_r} u^2 \, d\mathcal{H}^n \leq C\mathcal{H}_u(r).$$

The conclusion then follows at once. ■

We now prove the quasi-monotonicity of \mathcal{J}_u .

Proposition 7.5. *Let \mathbb{A} satisfy (H1) with $p = \infty$, (H2), (H3), and $\mathbb{A}(\underline{0}) = \text{Id}$, and let μ be as above. Then, there exists a constant $C > 0$ depending on n, λ, Λ , and $[\mathbb{A}]_{0,1}$ such that the function*

$$(0, 1] \ni r \mapsto e^{Cr} \mathcal{J}_u(r)$$

is non-decreasing, where we recall that $\mathcal{J}_u(r) = \frac{r \mathcal{E}_u(r)}{\mathcal{H}_u(r)}$.

Proof. Propositions 7.3 and 7.4, formula (7.2), and the Cauchy–Schwarz inequality give for \mathcal{L}^1 a.e. $r \in (0, 1)$

$$\begin{aligned} \mathcal{J}'_u(r) &= \frac{d}{dr} \left(\frac{r \mathcal{E}_u(r)}{\mathcal{H}_u(r)} \right) = \frac{\mathcal{J}_u(r)}{r} + r \frac{\mathcal{E}'_u(r) \mathcal{H}_u(r) - \mathcal{E}_u(r) \mathcal{H}'_u(r)}{\mathcal{H}_u^2(r)} \\ &= \frac{\mathcal{J}_u(r)}{r} \\ &\quad + \frac{r}{\mathcal{H}_u^2(r)} \left(2\mathcal{H}_u(r) \int_{\partial B_r} \mu^{-1} \langle \mathbb{A}v, \nabla u \rangle^2 d\mathcal{H}^n - 2 \left(\int_{\partial B_r} u \langle \mathbb{A} \nabla u, v \rangle d\mathcal{H}^n \right)^2 \right) \\ &\quad + \frac{r}{\mathcal{H}_u^2(r)} \left(E_r \mathcal{H}_u(r) - \frac{n}{r} \mathcal{H}_u(r) \mathcal{E}_u(r) - H_r \mathcal{E}_u(r) \right) \\ &\geq \frac{\mathcal{J}_u(r)}{r} + \frac{r}{\mathcal{H}_u^2(r)} \left(E_r \mathcal{H}_u(r) - \frac{n}{r} \mathcal{H}_u(r) \mathcal{E}_u(r) - H_r \mathcal{E}_u(r) \right) \\ &\geq \frac{\mathcal{J}_u(r)}{r} + \frac{r}{\mathcal{H}_u^2(r)} \left(-C \mathcal{E}_u(r) \mathcal{H}_u(r) - \frac{1}{r} \mathcal{H}_u(r) \mathcal{E}_u(r) \right) = -C \mathcal{J}_u(r). \end{aligned}$$

The conclusion then follows at once. ■

The quasi-monotonicity of \mathcal{J}_u is exploited in what follows to show the finiteness of the intrinsic frequency N_u . To this aim, we will also need the following auxiliary result.

Lemma 7.6. *Let u be a solution to (2.1) under assumptions (H1)–(H3) with $p = \infty$, and $\mathbb{A}(\underline{0}) = \text{Id}$. Then, there exists $\beta > 0$ such that*

$$\frac{\mathcal{H}_u(t)}{t^\beta} e^{Ct} \leq \frac{\mathcal{H}_u(r)}{r^\beta} e^{Cr} \quad \forall 0 < r < t < 1.$$

Proof. From Proposition 7.4 and (7.2), we have for \mathcal{L}^1 a.e. $r \in (0, 1)$

$$\mathcal{H}'_u(r) \leq \frac{n}{r} \mathcal{H}_u(r) + 2\mathcal{E}_u(r) + C \mathcal{H}_u(r)$$

so that

$$\frac{\mathcal{H}'_u(r)}{\mathcal{H}_u(r)} \leq \frac{n}{r} + 2 \frac{\mathcal{J}_u(r)}{r} + C \leq \frac{n + C \mathcal{J}_u(1)}{r} + C,$$

from which we obtain

$$\frac{d}{dr} \ln \left(\frac{\mathcal{H}_u(r)}{r^\beta} \right) \leq C,$$

where $\beta = n + C \mathcal{J}_u(1)$. The conclusion then follows by a simple integration. ■

We can finally prove the finiteness of $N_u(x_0, 0^+)$ for every point x_0 in $\Gamma(u)$.

Proposition 7.7. *Let u be a solution to (2.1) under assumptions (H1)–(H3) with $p = \infty$. Then, $\Gamma^{\text{finite}}(u) = \Gamma(u)$, i.e., $N_u(x_0, 0^+) < \infty$ for every $x_0 \in \Gamma(u)$.*

Proof. Without loss of generality, we verify the finiteness of the frequency in $x_0 = \underline{0} \in \Gamma(u)$. Moreover, by the arguments in Section 5, it is enough to consider the case $\mathbb{A}(\underline{0}) = \text{Id}$: indeed, the intrinsic frequency function is defined after change of coordinates Φ_{x_0} which sets the matrix $\mathbb{A}(0)$ to be the identity. In the sequel, we use the convention adopted throughout the paper to drop the base point being equal to the origin.

We begin by estimating $H_u(r)$ and $D_u(r)$ in terms of $\mathcal{H}_u(r)$ and $\mathcal{E}_u(r)$, respectively. Let us begin with $H_u(r)$: from Lemma 7.6, we get

$$\begin{aligned} H_u(r) &= \int -\phi' \left(\frac{|x|}{r} \right) \frac{u^2}{|x|} dx = \int_{r/2}^r \int_{\partial B_s} -\phi' \left(\frac{|x|}{r} \right) \frac{u^2}{s} d\mathcal{H}^n ds \\ &\geq \int_{r/2}^r -\phi' \left(\frac{s}{r} \right) \frac{1}{\Lambda s} \int_{\partial B_s} u^2 \mu d\mathcal{H}^n ds = \int_{r/2}^r -\phi' \left(\frac{s}{r} \right) \frac{1}{\Lambda s} \mathcal{H}_u(s) ds \\ &\geq \int_{r/2}^r -\phi' \left(\frac{|x|}{r} \right) \frac{1}{\Lambda s} \mathcal{H}_u(r) e^{C(r-s)} \frac{s^\beta}{r^\beta} ds \\ &\geq C \mathcal{H}_u(r) \int_{r/2}^r -\phi' \left(\frac{|x|}{r} \right) \frac{1}{s} ds \geq C \mathcal{H}_u(r). \end{aligned} \tag{7.8}$$

Instead, for \mathcal{E}_u and D_u , we have

$$D_u(r) = \int_{B_r} \phi \left(\frac{|x|}{r} \right) |\nabla u|^2 dx \leq \int_{B_r} |\nabla u|^2 dx \leq \lambda^{-1} \mathcal{E}_u(r). \tag{7.9}$$

Thus, from (7.8), (7.9), and the definition of the frequency $I_u(r)$, we conclude by taking into account Proposition 7.5:

$$N_u(r) = I_u(r) = \frac{r D_u(r)}{H_u(r)} \leq C \frac{r \mathcal{E}_u(r)}{\mathcal{H}_u(r)} = C \mathcal{J}_u(r) \leq C \mathcal{J}_u(1) < \infty.$$

We then conclude that all points of the free boundary have finite frequency, i.e., $\Gamma(u) = \Gamma^{\text{finite}}(u)$. ■

We are then in the position to prove Theorem 1.2.

Proof of Theorem 1.2. The rectifiability of the free boundary is a consequence of Theorem 1.1 and the previous Proposition 7.7 which establishes the fact that all free boundary points belong to $\Gamma^{\text{finite}}(u)$.

In order to deduce the local finiteness of the Minkowski content of $\Gamma(u)$, we observe that the intrinsic frequency is locally bounded in $B'_{\frac{1}{2}} \cap \Gamma(u)$: i.e., there exists $m_0 > 0$ such that

$$B'_{\frac{1}{2}} \cap \Gamma(u) \subset \Gamma^{m_0}(u).$$

Indeed, we have that $N_u(x, r) \leq C\mathcal{J}_u(x, r) \leq C\mathcal{J}_u(x, 1/2)$ for every $r \in (0, \frac{1}{2}]$; taking into account the continuity of $\bar{B}'_{1/2} \cap \Gamma(u) \ni x \rightarrow \mathcal{J}_u(x, 1/2)$, we infer that $N_u(x, r)$ is bounded in $\bar{B}'_{1/2} \cap \Gamma(u)$ for every $r \in (0, 1/2]$.

By simple covering and scaling arguments, the conclusion of Theorem 1.2 is shown for every compact $K \subset\subset B'_1$. ■

7.1. Free boundary of nonlinear thin obstacle problems

The results proven above can be applied to the case of nonlinear thin obstacle problems studied in [1], i.e., to the class of problems

$$\min_{u \in \mathcal{A}} \int_{B_1^+} f(\nabla u) \, dx, \tag{7.10}$$

where the energy density $f : \mathbb{R}^{n+1} \rightarrow \mathbb{R}$ is convex and is of the form

$$f(p) = h(|p|) \tag{7.11}$$

for every $p \in \mathbb{R}^{n+1}$, and the matrix $D_p^2 f(p)$ is uniformly coercive on compact subsets, i.e., fulfills the following local ellipticity condition: for every $M > 0$, there exists $\lambda = \lambda(M) > 0$ such that

$$\langle D_p^2 f(p)\xi, \xi \rangle \geq \lambda|\xi|^2 \tag{7.12}$$

for every $|p| \leq M$ and $\xi \in \mathbb{R}^{n+1}$.

As shown in [1], if the function $h \in C^2(\mathbb{R})$ satisfies

$$h(0) = h'(0) = 0, \quad h''(t) = 1 + O(t) \quad \text{for } t \rightarrow 0^+,$$

then the solutions to the variational problem in (7.10) are $C_{\text{loc}}^{1,1/2}(B_1^+ \cup B'_1)$. Here, we show that, if in addition

$$h''(t) = 1 + O(t^2) \quad \text{for } t \rightarrow 0^+, \tag{7.13}$$

and, for the sake of simplicity, $h \in C^\infty$, then we may apply Theorem 1.2 to infer all the results on the free boundary regularity in that statement.

Proposition 7.8. *Let $u \in W_{\text{loc}}^{1,\infty}(B_1^+)$ be a solution to (7.10) in \mathcal{A} with $h \in C^\infty$ under the assumptions (7.11), (7.12), and (7.13); then, $u \in C_{\text{loc}}^{1,1/2}(B_1^+ \cup B'_1)$, and the free boundary $\Gamma(u)$ is $(n - 1)$ -rectifiable and its Minkowski content is locally finite; i.e., for every $K \subset\subset B'_1$, there exists a constant $C(K) > 0$ such that*

$$\mathcal{L}^{n+1}(\mathcal{T}_r(\Gamma(u) \cap K)) \leq C(K)r^2$$

for every $r \in (0, 1)$.

Moreover, there exists a set $\Sigma(u) \subset \Gamma(u)$ with Hausdorff dimension at most $n - 2$ such that for every $x \in \Gamma(u) \setminus \Sigma(u)$

$$N_u(x, 0^+) \in \{2m, 2m - 1/2, 2m + 1\}_{m \in \mathbb{N} \setminus \{0\}}.$$

Proof. The solution to (7.10) is $C_{\text{loc}}^{1,1/2}(B_1^+ \cup B'_1)$ by [1] and by standard elliptic regularity $u \in C^\infty(B_1^+)$ (thanks to the simplifying assumption $h \in C^\infty$). Moreover, u can be characterized as the weak solution to the system

$$\begin{cases} \operatorname{div}(\nabla_p f(\nabla u)) = 0 & \text{in } B_1^+, \\ u \partial_{n+1} f(\nabla u) = 0 & \text{on } B'_1, \\ -\partial_{n+1} f(\nabla u) \geq 0 & \text{on } B'_1, \\ u \geq 0 & \text{on } B'_1, \\ u = g & \text{on } (\partial B_1)^+. \end{cases} \tag{7.14}$$

In particular, we deduce from the first equation in (7.14) that for every $\phi \in C^1(B_1^+)$ with support non-intersecting $(\partial B_1)^+$

$$\int_{B_1^+} \langle \nabla_p f(\nabla u), \nabla \phi(x) \rangle dx = 0. \tag{7.15}$$

We can write assumption (7.13) in the form $h''(t) = 1 + \omega(t)$ with $|\omega(t)| \leq Ct^2$ for t sufficiently small. Integrating, we infer that

$$h'(t) = t(1 + \tilde{\omega}(t)), \quad \tilde{\omega}(t) = \frac{1}{t} \int_0^t \omega(s) ds,$$

and $\tilde{\omega} \in C^1$ with $\tilde{\omega}(0) = \tilde{\omega}'(0) = 0$ and

$$\tilde{\omega}'(t) = -\frac{1}{t^2} \int_0^t \omega(s) ds + \frac{1}{t} \omega(t) \implies |\tilde{\omega}'(t)| \leq Ct,$$

for t sufficiently small. The first variation (7.15) reads then for every $\phi \in C^1(B_1^+)$ with support non-intersecting $(\partial B_1)^+$

$$\int_{B_1^+} \langle (1 + \tilde{\omega}(|\nabla u(x)|)) \nabla u(x), \nabla \phi(x) \rangle dx = 0,$$

which is the Euler–Lagrange equation of the linear thin obstacle problem driven by the quadratic energy

$$\int_{B_1^+} \theta(x) |\nabla u(x)|^2 dx, \quad \theta(x) := 1 + \tilde{\omega}(|\nabla u(x)|).$$

Note that $1 \leq \theta(x) \leq 1 + C|\nabla u(x)|$; therefore, θ is locally bounded on $B_1^+ \cup B'_1$. Thus, if we prove that the function θ is locally Lipschitz continuous on $B_1^+ \cup B'_1$, we can apply Theorem 1.2 and conclude all the results about the structure of $\Gamma(u)$.

To this aim, we notice that for nontrivial solutions u in B_1^+ we have

$$\nabla \theta(x) = \tilde{\omega}'(|\nabla u(x)|) D^2 u(x) \frac{\nabla u(x)}{|\nabla u(x)|} \quad \text{if } |\nabla u(x)| \neq 0.$$

Extend for simplicity u by even reflection to the whole B_1 (without renaming the function u), and let $d : B_1 \rightarrow [0, \infty)$ be the distance from the free boundary $\Gamma(u)$. By (7.14), the function u satisfies the nonlinear elliptic equation

$$\operatorname{div}(\nabla_p f(\nabla u)) = 0 \quad \text{on } B_1 \setminus \{(x', 0) : u(x', 0) = 0\},$$

and thus, the following classical elliptic estimates hold locally in $B_1 \setminus \{(x', 0) : u(x', 0) = 0\}$:

$$\begin{aligned} |\nabla u(x)| &\leq C d(x)^{-1} \|u\|_{L^\infty(B_{d(x)}(x))} \leq C d(x)^{1/2}, \\ |D^2 u(x)| &\leq C d(x)^{-2} \|u\|_{L^\infty(B_{d(x)}(x))} \leq C d(x)^{-1/2}. \end{aligned}$$

Recalling that by assumption

$$\tilde{\omega}'(|\nabla u(x)|) \leq C |\nabla u(x)|,$$

we then conclude for points outside the contact set, i.e., $B_1 \setminus \{(x', 0) : u(x', 0) = 0\}$, that

$$|\nabla \theta(x)| \leq \frac{|\tilde{\omega}'(|\nabla u(x)|)|}{|\nabla u(x)|} |D^2 u(x)| |\nabla u(x)| \leq C d(x)^{-1/2} d(x)^{1/2} \leq C. \quad (7.16)$$

Moreover, if $x \in \Gamma(u)$, then $|\nabla u(x)| = 0$ so that for every $y \in B_1^+ \cup B_1'$

$$\begin{aligned} |\theta(x) - \theta(y)| &= |\tilde{\omega}(|\nabla u(y)|)| \leq \int_0^1 |\tilde{\omega}'(t|\nabla u(y))| |\nabla u(y)| dt \\ &\leq C |\nabla u(y)|^2 \leq C |x - y|, \end{aligned}$$

using the optimal regularity of u . Finally, if x belongs to the relative interior of $\{(x', 0) : u(x', 0) = 0\}$ in B_1' , we use the odd reflection across the hyperplane $\{x_{n+1} = 0\}$ as in [1, Theorem 4.1] to infer that (7.16) holds as well.

In conclusion, θ is locally Lipschitz continuous on $B_1^+ \cup B_1'$. ■

A. Order of contact

We introduce the definition of lower and upper order of contact with zero at a point.

Definition A.1. Let $v \in H^1(\Omega)$, $x_0 \in \Omega \subset \mathbb{R}^{n+1}$, the lower and upper orders of contact with 0 of v at x_0 are defined, respectively as

$$\underline{\vartheta}(x_0) := \sup \left\{ \vartheta \in \mathbb{R} : \limsup_{\rho \rightarrow 0^+} \frac{H_v(x_0, \rho)}{\rho^{n+2\vartheta}} < \infty \right\}, \quad (A.1)$$

$$\bar{\vartheta}(x_0) := \inf \left\{ \vartheta \in \mathbb{R} : \liminf_{\rho \rightarrow 0^+} \frac{H_v(x_0, \rho)}{\rho^{n+2\vartheta}} > 0 \right\}. \quad (A.2)$$

Few elementary properties of $\underline{\vartheta}(x_0)$ and $\bar{\vartheta}(x_0)$ are resumed in the ensuing list: for all $x_0 \in \Omega$, we have

(1) $-\infty \leq \underline{\vartheta}(x_0) \leq \bar{\vartheta}(x_0) \leq \infty$,

(2) the lower order of contact can be characterized equivalently as

$$\underline{\vartheta}(x_0) = \sup \left\{ \vartheta \in \mathbb{R} : \lim_{\rho \rightarrow 0^+} \frac{H_u(x_0, \rho)}{\rho^{n+2\vartheta}} = 0 \right\},$$

(3) the upper order of contact can be characterized equivalently as

$$\bar{\vartheta}(x_0) = \inf \left\{ \vartheta \in \mathbb{R} : \lim_{\rho \rightarrow 0^+} \frac{H_u(x_0, \rho)}{\rho^{n+2\vartheta}} = \infty \right\}.$$

Additionally, we compare the latter notions with those used by Koch, Rüländ, and Shi [28].

Proposition A.2. *Let $u \in \mathcal{A}$ be a solution to (1.1) under the hypotheses (H1) and (H2), and $x_0 \in \Gamma(u)$ with $\mathbb{A}(x_0) = \text{Id}$. Then, on setting $A_\rho(x_0) := B_\rho(x_0) \setminus B_{\rho/2}(x_0)$, we have that*

$$\underline{\vartheta}(x_0) = \liminf_{\rho \rightarrow 0^+} \frac{\ln(\int_{A_\rho(x_0)} u^2 dx)^{1/2}}{\ln \rho} =: \underline{\kappa}(x_0), \tag{A.3}$$

$$\bar{\vartheta}(x_0) = \limsup_{\rho \rightarrow 0^+} \frac{\ln(\int_{A_\rho(x_0)} u^2 dx)^{1/2}}{\ln \rho} =: \bar{\kappa}(x_0). \tag{A.4}$$

Proof. We will only prove the equality in (A.3), the other in (A.4) being completely analogous. We first note that by Lemma 3.8

$$\frac{1}{2} \|\phi'\|_\infty^{-1} \rho H_u(x_0, \rho) \leq \int_{A_\rho(x_0)} |u(x)|^2 dx \stackrel{(3.16)}{\leq} C \rho H_u(x_0, \rho), \tag{A.5}$$

for points on the free boundary with $\mathbb{A}(x_0) = \text{Id}$.

Assume $\underline{\kappa}(x_0) \in \mathbb{R}$; then, for every $\varepsilon > 0$, there are $\rho_\varepsilon \in (0, 1)$ and $\rho_j \downarrow 0$ such that for all $\rho \in (0, \rho_\varepsilon)$

$$\int_{A_\rho(x_0)} u^2 dx \leq \rho^{2(\underline{\kappa}(x_0) - \varepsilon)},$$

and for all $j \in \mathbb{N}$

$$\int_{A_{\rho_j}(x_0)} u^2 dx \geq \rho_j^{2(\underline{\kappa}(x_0) + \varepsilon)}.$$

From the former inequality and (A.5), we infer that $\underline{\kappa}(x_0) - \varepsilon \leq \underline{\vartheta}(x_0)$, and thus, $\underline{\kappa}(x_0) \leq \underline{\vartheta}(x_0)$. Instead, from the latter inequality and (A.5), we deduce that $\underline{\kappa}(x_0) + 2\varepsilon > \underline{\vartheta}(x_0)$; thus, $\underline{\kappa}(x_0) \geq \underline{\vartheta}(x_0)$. Therefore, $\underline{\kappa}(x_0) = \underline{\vartheta}(x_0)$.

If $\underline{\kappa}(x_0) = -\infty$, then there exists $\rho_j \downarrow 0$ such that for all $i \in \mathbb{N}$ there exists $j_i \in \mathbb{N}$ such that for all $j \geq j_i$

$$\int_{A_{\rho_j}(x_0)} u^2 dx \geq \rho_j^{-2i},$$

and thus, $-i + \frac{1}{2} > \underline{\vartheta}(x_0)$, in turn implying $\underline{\vartheta}(x_0) = -\infty$.

If $\underline{\kappa}(x_0) = \infty$, then for every $i \in \mathbb{N}$ there is $\rho_i \in (0, 1)$ such that for all $\rho \in (0, \rho_i)$

$$\int_{A_\rho(x_0)} u^2 \, dx \leq \rho^{2i},$$

from which we conclude that $\underline{\vartheta}(x_0) \geq i$, and thus, $\underline{\vartheta}(x_0) = \infty$.

In conclusion, $\underline{\vartheta}(x_0) = \underline{\kappa}(x_0)$ in all possible instances. ■

For solutions to the thin obstacle problem, the points with finite frequency are points with finite order of contact.

Lemma A.3. *Let u be a solution to the thin obstacle problem (2.2) in B_1 . Then, for every $x_0 \in \Gamma(u)$,*

$$\limsup_{r \rightarrow 0^+} I_u(x_0, r) \geq \bar{\vartheta}(x_0). \tag{A.6}$$

Moreover, if $\limsup_{r \rightarrow 0^+} I_u(x_0, r) < \infty$, then the limsup is actually a limit and

$$\lim_{r \rightarrow 0^+} I_u(x_0, r) = \underline{\vartheta}(x_0) = \bar{\vartheta}(x_0) \in [3/2, \infty). \tag{A.7}$$

Proof. Without loss of generality, we take $x_0 = \underline{0}$, and set $\bar{I}_u(0^+) := \limsup_{r \rightarrow 0^+} I_u(\underline{0}, r)$.

We start off proving (A.6). Without loss of generality, we assume $\bar{I}_u(0^+) < \infty$, the inequality being trivial otherwise. Hence, the doubling of both $H_u(\underline{0}, \cdot)$ and $D_u(\underline{0}, \cdot)$ holds thanks to Proposition 3.3. Then, we use the equality in (3.17), namely,

$$H'_u(r) = \frac{n}{r} H_u(r) + 2G_u(r),$$

and (3.9) with $x = x_0 = \underline{0}$ and $\kappa = 0$, to infer that

$$\begin{aligned} \left| H'_u(r) - \frac{n}{r} H_u(r) - 2D_u(r) \right| &\leq C r^\alpha \left(D_u(r) + \frac{1}{r^{1/2}} H_u^{1/2}(r) D_u^{1/2}(r) \right) \\ &\leq C r^\alpha D_u(r) (1 + I_u^{-1/2}(r)), \end{aligned}$$

in turn implying

$$\begin{aligned} \left| \frac{d}{dr} \ln \left(\frac{H_u(r)}{r^n} \right) - \frac{2}{r} I_u(r) \right| &\leq C r^{\alpha-1} I_u(r) (1 + I_u^{-1/2}(r)) \\ &= C r^{\alpha-1} (I_u(r) + I_u^{1/2}(r)). \end{aligned} \tag{A.8}$$

Then, for every $\varepsilon > 0$, there is $r_\varepsilon > 0$ such that $I_u(r) \leq \bar{I}_u(0^+) + \varepsilon$ for every $r \in (0, r_\varepsilon)$.

We use (A.8) to deduce for such radii that

$$\frac{d}{dr} \ln \left(\frac{H_u(r)}{r^n} \right) \leq \frac{2}{r} (\bar{I}_u(0^+) + \varepsilon) + C r^{\alpha-1}.$$

Hence, by direct integration, we get that for all $0 < r < s < r_\varepsilon$

$$0 < \frac{H_u(s)}{s^{n+2(\bar{I}_u(0^+)+\varepsilon)}} e^{-\frac{C}{\alpha} s^\alpha} \leq \frac{H_u(r)}{r^{n+2(\bar{I}_u(0^+)+\varepsilon)}} e^{-\frac{C}{\alpha} r^\alpha}.$$

From this and the very definition of $\bar{\vartheta}(0)$ in (A.2), we have $\bar{\vartheta}(0) \leq \bar{I}_u(0^+) + \varepsilon$ for every $\varepsilon > 0$, which implies (A.6).

In order to prove (A.7), we combine the results in (A.6) with those in Proposition 3.9 (cf. (3.20)) to infer that the lim sup of the frequency is actually a limit so that the latter is rewritten as

$$\lim_{r \rightarrow 0^+} I_u(r) \geq \bar{\vartheta}(0). \tag{A.9}$$

Therefore, arguing as above, by the inequality in (3.26) of Corollary 3.11 and (A.8), we get that for every $r \in (0, r_\varepsilon)$

$$\frac{d}{dr} \ln\left(\frac{H_u(r)}{r^n}\right) \geq \frac{2}{r}(\bar{I}_u(0^+) - \varepsilon) - Cr^{\alpha-1},$$

from which we conclude by integration that for all $0 < r < s < r_\varepsilon$

$$0 < \frac{H_u(r)}{r^{n+2(\bar{I}_u(0^+) - \varepsilon)}} e^{\frac{c}{\alpha} r^\alpha} \leq \frac{H_u(s)}{s^{n+2(\bar{I}_u(0^+) - \varepsilon)}} e^{\frac{c}{\alpha} s^\alpha}.$$

Hence, we deduce that $\bar{I}_u(0^+) - \varepsilon \leq \bar{\vartheta}(0)$ for every $\varepsilon > 0$; (A.7) then follows at once from the last inequality, (A.9), and Corollary 3.11. ■

As a consequence of a Carleman-type estimate in [28], it is established there that for the solutions to the variable coefficients thin obstacle problem:

- (a) $\underline{\vartheta}(x_0) = \bar{\vartheta}(x_0)$ for every $x_0 \in B'_1$;
- (b) if $\underline{\vartheta}(x_0) < \infty$, then doubling for $H_u(x_0, \cdot)$ holds provided $\mathbb{A}(x_0) = \text{Id}$.

Items (a) and (b) right above yield the doubling of $H_{u_{\mathbb{A}(x_0)}}(x_0, \cdot)$, in turn implying that for $D_{u_{\mathbb{A}(x_0)}}(x_0, \cdot)$ thanks to an elementary Cacciopoli inequality. The latter and the proof of Proposition 3.9 imply the quasi-monotonicity of $N_u(x_0, \cdot) = I_{u_{\mathbb{A}(x_0)}}(x_0, \cdot)$ and thus the finiteness of N_u . On the other hand, item (a) of Lemma A.3 shows that points with finite frequency have finite order of contact. Therefore, we infer the following corollary.

Corollary A.4. *Let $u \in \mathcal{A}$ be a solution to (1.1) under the hypotheses (H1) and (H2). Then, the subset of points of the free boundary with finite order of contact is well defined:*

$$\begin{aligned} \Gamma^{\text{finite}}(u) &= \{x_0 \in \Gamma(u) : \limsup_{r \rightarrow 0^+} N_u(x_0, r) < \infty\} \\ &= \{x_0 \in \Gamma(u) : \underline{\kappa}(x_0) = \bar{\kappa}(x_0) < \infty\}. \end{aligned}$$

In particular, the points with finite order of contact do not depend on the choice of the cut-off function ϕ in the definition of the frequency function.

Funding. G. Andreucci and M. Focardi are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). M. Focardi has been supported by the European Union – Next Generation EU, Mission 4 Component 1 CUP 2022J4FYNJ, PRIN2022

project “Variational methods for stationary and evolution problems with singularities and interfaces”. E. Spadaro has been partially supported by the ERC-STG grant 759229 HiCoS and by PRIN 2022PJ9EFL “Geometric Measure Theory: Structure of Singular Measures, Regularity Theory and Applications in the Calculus of Variations” funded by the European Union – Next Generation EU, Mission 4 Component 2 – CUP: E53D23005860006.

References

- [1] A. Abbatiello, G. Andreucci, and E. Spadaro, On the nonlinear thin obstacle problem. 2024, arXiv:2403.19487v1, to appear in *Rev. Mat. Iberoam*.
- [2] F. J. Almgren, Jr., *Almgren’s big regularity paper*. World Sci. Monogr. Ser. Math. 1, World Scientific, River Edge, NJ, 2000 Zbl 0985.49001 MR 1777737
- [3] G. Andreucci and M. Focardi, [The classical obstacle problem with Hölder continuous coefficients](#). *Nonlinear Anal.* **226** (2023), article no. 113139 Zbl 1501.35222 MR 4484854
- [4] I. Athanasopoulos and L. A. Caffarelli, [Optimal regularity of lower dimensional obstacle problems](#). *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **310** (2004), no. Kraev. Zadachi Mat. Fiz. i Smezh. Vopr. Teor. Funkts. 35 [34], 49–66 Zbl 1108.35038 MR 2120184
- [5] I. Athanasopoulos, L. A. Caffarelli, and S. Salsa, [The structure of the free boundary for lower dimensional obstacle problems](#). *Amer. J. Math.* **130** (2008), no. 2, 485–498 Zbl 1185.35339 MR 2405165
- [6] L. A. Caffarelli, [Further regularity for the Signorini problem](#). *Comm. Partial Differential Equations* **4** (1979), no. 9, 1067–1075 Zbl 0427.35019 MR 0542512
- [7] M. Colombo, L. Spolaor, and B. Velichkov, [Direct epiperimetric inequalities for the thin obstacle problem and applications](#). *Comm. Pure Appl. Math.* **73** (2020), no. 2, 384–420 Zbl 1433.49010 MR 4054360
- [8] D. De Silva and O. Savin, [\$C^\infty\$ regularity of certain thin free boundaries](#). *Indiana Univ. Math. J.* **64** (2015), no. 5, 1575–1608 Zbl 1331.35397 MR 3418452
- [9] L. Di Fazio and E. Spadaro, [Regularity of solutions to nonlinear thin and boundary obstacle problems](#). *Adv. Math.* **401** (2022), article no. 108263 Zbl 1494.35198 MR 4387851
- [10] X. Fernández-Real and X. Ros-Oton, [Free boundary regularity for almost every solution to the Signorini problem](#). *Arch. Ration. Mech. Anal.* **240** (2021), no. 1, 419–466 Zbl 07331727 MR 4228865
- [11] N. Filonov, [Second-order elliptic equation of divergence form having a compactly supported solution](#). *J. Math. Sci. (New York)* **106** (2001), no. 3, 3078–3086 Zbl 0991.35020 MR 1906035
- [12] M. Focardi, M. S. Gelli, and E. Spadaro, [Monotonicity formulas for obstacle problems with Lipschitz coefficients](#). *Calc. Var. Partial Differential Equations* **54** (2015), no. 2, 1547–1573 Zbl 1328.35332 MR 3396423
- [13] M. Focardi, F. Geraci, and E. Spadaro, [Quasi-monotonicity formulas for classical obstacle problems with Sobolev coefficients and applications](#). *J. Optim. Theory Appl.* **184** (2020), no. 1, 125–138 Zbl 1431.35251 MR 4047132
- [14] M. Focardi and E. Spadaro, [An epiperimetric inequality for the thin obstacle problem](#). *Adv. Differential Equations* **21** (2016), no. 1-2, 153–200 Zbl 1336.35370 MR 3449333

- [15] M. Focardi and E. Spadaro, [Correction to: On the measure and the structure of the free boundary of the lower dimensional obstacle problem](#). *Arch. Ration. Mech. Anal.* **230** (2018), no. 2, 783–784 Zbl 1466.35376 MR 3842059
- [16] M. Focardi and E. Spadaro, [On the measure and the structure of the free boundary of the lower dimensional obstacle problem](#). *Arch. Ration. Mech. Anal.* **230** (2018), no. 1, 125–184 Zbl 1405.35257 MR 3840912
- [17] M. Focardi and E. Spadaro, [How a minimal surface leaves a thin obstacle](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **37** (2020), no. 4, 1017–1046 Zbl 1461.35250 MR 4104833
- [18] M. Focardi and E. Spadaro, [The local structure of the free boundary in the fractional obstacle problem](#). *Adv. Calc. Var.* **15** (2022), no. 3, 323–349 Zbl 1501.35463 MR 4451895
- [19] N. Garofalo and A. Petrosyan, [Some new monotonicity formulas and the singular set in the lower dimensional obstacle problem](#). *Invent. Math.* **177** (2009), no. 2, 415–461 Zbl 1175.35154 MR 2511747
- [20] N. Garofalo, A. Petrosyan, and M. Smit Vega Garcia, [An epiperimetric inequality approach to the regularity of the free boundary in the Signorini problem with variable coefficients](#). *J. Math. Pures Appl. (9)* **105** (2016), no. 6, 745–787 Zbl 1341.35036 MR 3491531
- [21] N. Garofalo, A. Petrosyan, and M. Smit Vega Garcia, [The singular free boundary in the Signorini problem for variable coefficients](#). *Indiana Univ. Math. J.* **67** (2018), no. 5, 1893–1934 Zbl 1409.35239 MR 3875246
- [22] N. Garofalo and M. Smit Vega Garcia, [New monotonicity formulas and the optimal regularity in the Signorini problem with variable coefficients](#). *Adv. Math.* **262** (2014), 682–750 Zbl 1295.35155 MR 3228440
- [23] S. Jeon and A. Petrosyan, [Almost minimizers for the thin obstacle problem](#). *Calc. Var. Partial Differential Equations* **60** (2021), no. 4, article no. 124 Zbl 1468.49041 MR 4277328
- [24] S. Jeon, A. Petrosyan, and M. Smit Vega Garcia, [Almost minimizers for the thin obstacle problem with variable coefficients](#). *Interfaces Free Bound.* **26** (2024), no. 3, 321–380 Zbl 07902357 MR 4762086
- [25] D. Kinderlehrer, [The smoothness of the solution of the boundary obstacle problem](#). *J. Math. Pures Appl. (9)* **60** (1981), no. 2, 193–212 Zbl 0459.35092 MR 0620584
- [26] H. Koch, A. Petrosyan, and W. Shi, [Higher regularity of the free boundary in the elliptic Signorini problem](#). *Nonlinear Anal.* **126** (2015), 3–44 Zbl 1329.35362 MR 3388870
- [27] H. Koch, A. Rüländ, and W. Shi, [The variable coefficient thin obstacle problem: Carleman inequalities](#). *Adv. Math.* **301** (2016), 820–866 Zbl 1346.35240 MR 3539391
- [28] H. Koch, A. Rüländ, and W. Shi, [The variable coefficient thin obstacle problem: Optimal regularity and regularity of the regular free boundary](#). *Ann. Inst. H. Poincaré C Anal. Non Linéaire* **34** (2017), no. 4, 845–897 Zbl 1435.35421 MR 3661863
- [29] N. Mandache, [On a counterexample concerning unique continuation for elliptic equations in divergence form](#). *Math. Phys. Anal. Geom.* **1** (1998), no. 3, 273–292 Zbl 0920.35034 MR 1686671
- [30] K. Miller, [Nonunique continuation for uniformly parabolic and elliptic equations in self-adjoint divergence form with Hölder continuous coefficients](#). *Arch. Rational Mech. Anal.* **54** (1974), 105–117 Zbl 0289.35046 MR 0342822
- [31] A. Naber and D. Valtorta, [Rectifiable-Reifenberg and the regularity of stationary and minimizing harmonic maps](#). *Ann. of Math. (2)* **185** (2017), no. 1, 131–227 Zbl 1393.58009 MR 3583353
- [32] A. Naber and D. Valtorta, [The singular structure and regularity of stationary varifolds](#). *J. Eur. Math. Soc. (JEMS)* **22** (2020), no. 10, 3305–3382 Zbl 1460.35143 MR 4153109

- [33] A. Pliš, On non-uniqueness in Cauchy problem for an elliptic second order differential equation. *Bull. Acad. Polon. Sci. Sér. Sci. Math. Astronom. Phys.* **11** (1963), 95–100
Zbl [0107.07901](#) MR [0153959](#)
- [34] A. Rüländ and W. Shi, [Optimal regularity for the thin obstacle problem with \$C^{0,\alpha}\$ coefficients](#). *Calc. Var. Partial Differential Equations* **56** (2017), no. 5, article no. 129 Zbl [1378.35340](#)
MR [3689152](#)
- [35] L. Simon and N. Wickramasekera, [A frequency function and singular set bounds for branched minimal immersions](#). *Comm. Pure Appl. Math.* **69** (2016), no. 7, 1213–1258 Zbl [1341.53097](#)
MR [3503021](#)
- [36] N. N. Ural'tseva, Hölder continuity of gradients of solutions of parabolic equations with boundary conditions of Signorini type. *Dokl. Akad. Nauk SSSR* **280** (1985), no. 3, 563–565
Zbl [0629.35047](#) MR [0775926](#)
- [37] N. N. Ural'tseva, Estimation on the boundary of the domain of derivatives of solutions of variational inequalities. In *Linear and nonlinear boundary value problems. Spectral theory (Russian)*, pp. 92–105. Translated in *J. Soviet Math.* **45** (1989), no. 3, 1181–1191, Probl. Mat. Anal. 10, Leningrad University, Leningrad, 1986 Zbl [0703.35076](#) MR [860572](#)
- [38] N. N. Ural'tseva, [Regularity of solutions of variational inequalities](#) (in Russian). *Uspekhi Mat. Nauk* **42** (1987), no. 6(258), 151–174. [English translation: Russ. Math. Surv.](#) **42** (1987), no. 6, 191–219 Zbl [0696.49022](#) MR [0933999](#)

Received 22 July 2024.

Giovanna Andreucci

Dipartimento di Matematica, Sapienza Università di Roma, Piazzale Aldo Moro 5, 00185 Rome, Italy; giovanna.andreucci@uniroma1.it

Matteo Focardi

DIMAI, Università degli Studi di Firenze, Viale Morgagni 67/A, 50134 Firenze, Italy; matteo.focardi@unifi.it

Emanuele Spadaro

Dipartimento di Matematica, Sapienza Università di Roma, Piazzale Aldo Moro 5, 00185 Rome, Italy; spadaro@mat.uniroma1.it