

Yetter–Drinfeld post-Hopf algebras and Yetter–Drinfeld relative Rota–Baxter operators

Andrea Sciandra

Abstract. Recently, Li, Sheng, and Tang introduced post-Hopf algebras and relative Rota–Baxter operators (on cocommutative Hopf algebras), providing an adjunction between the respective categories under the assumption that the structures involved are cocommutative. We introduce Yetter–Drinfeld post-Hopf algebras, which become usual post-Hopf algebras in the cocommutative setting. In analogy with the correspondence between cocommutative post-Hopf algebras and cocommutative Hopf braces, the category of Yetter–Drinfeld post-Hopf algebras is isomorphic to the category of Yetter–Drinfeld braces introduced by the author in a joint work with D. Ferri. This allows to explore the connection with matched pairs of actions and provide examples of Yetter–Drinfeld post-Hopf algebras. Moreover, we prove that the category of Yetter–Drinfeld post-Hopf algebras is equivalent to a subcategory of bijective Yetter–Drinfeld relative Rota–Baxter operators. The latter structures coincide with the inverse maps of Yetter–Drinfeld 1-cocycles introduced by the author and D. Ferri and generalise bijective relative Rota–Baxter operators on cocommutative Hopf algebras. Hence, the previous equivalence passes to cocommutative post-Hopf algebras and bijective relative Rota–Baxter operators. Once the surjectivity of the Yetter–Drinfeld relative Rota–Baxter operators is removed, the equivalence is replaced by an adjunction and one can recover the result of Li, Sheng, and Tang in the cocommutative case.

1. Introduction

Post-Lie algebras have been introduced in [30] and have found several interesting applications [7, 10, 25], just to name a few; while Rota–Baxter operators on Lie algebras first appeared in [19] and are now studied in different fields, such as quantum field theory [9, 18]. Post-Lie algebras and Rota–Baxter operators are strictly related: given a post-Lie algebra, the identity morphism is a Rota–Baxter operator on the subadjacent Lie algebra and, vice versa, one can induce a post-Lie algebra structure on the domain of a Rota–Baxter operator. In [22], Li, Sheng, and Tang introduced post-Hopf algebras and relative Rota–Baxter operators on cocommutative Hopf algebras, the latter generalising Rota–Baxter operators given in [13] (where the action involved is the adjoint one). The space of primitive elements of a post-Hopf algebra results to be a post-Lie algebra and a relative

Mathematics Subject Classification 2020: 16T05 (primary); 17B38 (secondary).

Keywords: Yetter–Drinfeld post-Hopf algebras, Yetter–Drinfeld relative Rota–Baxter operators, Yetter–Drinfeld braces, matched pairs, 1-cocycles, Hopf braces.

Rota–Baxter operator on a cocommutative Hopf algebra restricts to a Rota–Baxter operator between the Lie algebras of primitive elements. Moreover, in [22], the aforementioned connection between post-Lie algebras and Rota–Baxter operators is extended to cocommutative post-Hopf algebras and relative Rota–Baxter operators on cocommutative Hopf algebras, providing an adjunction between the respective categories. In [21] (which is the arXiv version of [22]), a correspondence between cocommutative post-Hopf algebras and cocommutative Hopf braces is exhibited, then further investigated [31] in terms of relative Rota–Baxter operators. Hopf braces were introduced by Angiono, Galindo and Vendramin in [2] as the Hopf-theoretic version of skew braces, introduced in [14]. Moreover, cocommutative Hopf braces are shown to be equivalent to matched pairs of cocommutative Hopf algebras. In [12], the author and D. Ferri proved that, removing the cocommutativity hypothesis, matched pairs result to be equivalent to the more general Yetter–Drinfeld braces. The latter structures coincide with Hopf braces if the cocommutativity is assumed.

The purpose of this paper is to generalise post-Hopf algebras and relative Rota–Baxter operators in a non-cocommutative setting, keeping in mind the connection with Yetter–Drinfeld braces.

More precisely, the content of the paper is the following. In Section 2, we recall the notions and results that motivate this work and are frequently used throughout the paper. In Section 3, we introduce Yetter–Drinfeld post-Hopf algebras, which coincide with post-Hopf algebras in the cocommutative case, and we prove that the category of Yetter–Drinfeld post-Hopf algebras is isomorphic to the category of Yetter–Drinfeld braces (and thus to the category of matched pairs of actions). Explicitly, a Yetter–Drinfeld post-Hopf algebra is the datum $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ where $(H, \cdot, 1)$ is an algebra, (H, Δ, ε) is a coalgebra, S is the convolution inverse of the morphism Id_H and \rightharpoonup is a morphism of coalgebras satisfying some technical hypotheses. Then, one can build the subadjacent Hopf algebra $H_{\rightharpoonup} := (H, \bullet_{\rightharpoonup}, 1, \Delta, \varepsilon, S_{\rightharpoonup})$ so that $(H, \cdot, 1, \Delta, \varepsilon, S)$ results to be an object in $\text{Hopf}(\overset{H}{\underset{H_{\rightharpoonup}}{\mathcal{YD}}})$ and $(H, \cdot, \bullet_{\rightharpoonup}, 1, \Delta, \varepsilon, S, S_{\rightharpoonup})$ becomes a Yetter–Drinfeld brace. A Yetter–Drinfeld post-Hopf algebra induces a post-Lie algebra structure on its space of primitive elements. At the end of the section, we provide examples of Yetter–Drinfeld post-Hopf algebras coming from Yetter–Drinfeld braces. Finally, in Section 4, we introduce Yetter–Drinfeld relative Rota–Baxter operators; these generalise relative Rota–Baxter operators on cocommutative Hopf algebras and, under the assumption of bijectivity, they coincide with the inverse maps of Yetter–Drinfeld 1-cocycles introduced in [12]. As in the cocommutative case, we can restrict Yetter–Drinfeld relative Rota–Baxter operators to primitive elements and group-like elements, obtaining Rota–Baxter operators on Lie algebras and groups, respectively. We show that a subcategory of bijective Yetter–Drinfeld relative Rota–Baxter operators is equivalent to the category of Yetter–Drinfeld post-Hopf algebras: given a Yetter–Drinfeld post-Hopf algebra (H, \rightharpoonup) the identity morphism $\text{Id}_H : H \rightarrow H_{\rightharpoonup}$ is a Yetter–Drinfeld relative Rota–Baxter operator and, vice versa, one can induce a structure of Yetter–Drinfeld post-Hopf algebra on the domain (or codomain) of a bijective Yetter–Drinfeld relative Rota–Baxter operator. This equivalence passes to cocommutative post-Hopf algebras and bijective relative Rota–Baxter operators on cocommutative

Hopf algebras. Once the surjectivity of the Yetter–Drinfeld relative Rota–Baxter operators is removed, the equivalence is replaced by an adjunction and one can recover, in the cocommutative case, the result given in [22]. Then, the aforementioned correspondence between post-Lie algebras and Rota–Baxter operators, generalised in [22] for cocommutative Hopf algebras, is now given in full generality.

Notations and conventions. We denote by \mathbb{k} an arbitrary field and all vector spaces will be \mathbb{k} -vector spaces. By a linear map we mean a \mathbb{k} -linear map and the unadorned tensor product \otimes is the one of \mathbb{k} -vector spaces. Algebras over \mathbb{k} will be associative and unital and coalgebras over \mathbb{k} will be coassociative and counital. We use symbols like \bullet and \cdot for the multiplication of an algebra; equivalently, the multiplication will be denoted by m, m_\bullet, m_\cdot . The unit of an algebra A will be denoted by 1 or $u : \mathbb{k} \rightarrow A$. The comultiplication and the counit of a coalgebra C will be denoted by Δ and ε , respectively. We will use Sweedler’s notation for calculations involving the coproduct, i.e., we write $\Delta(c) = c_1 \otimes c_2$ for all $c \in C$, omitting the summation. The antipode of a Hopf algebra will be denoted by S or T .

2. Preliminaries

In this section, we recall the notions and results that we are going to use throughout the paper. For more details about Hopf algebras and category theory, we refer the reader to [24, 27, 29] and [6], respectively.

Post-Lie algebras and relative Rota–Baxter operators. The notion of post-Lie algebra was introduced in [30] and it has found several interesting applications, such as [7, 10, 25]. Recall that a *post-Lie algebra* $(\mathfrak{g}, [\cdot, \cdot], \rightharpoonup)$ is the datum of a Lie algebra $(\mathfrak{g}, [\cdot, \cdot])$ and a linear map $\rightharpoonup : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ such that, for all $x, y, z \in \mathfrak{g}$, the following equalities hold:

$$x \rightharpoonup [y, z] = [x \rightharpoonup y, z] + [y, x \rightharpoonup z], \tag{2.1}$$

$$([x, y] + (x \rightharpoonup y) - (y \rightharpoonup x)) \rightharpoonup z = (x \rightharpoonup (y \rightharpoonup z)) - (y \rightharpoonup (x \rightharpoonup z)). \tag{2.2}$$

If $(\mathfrak{g}, [\cdot, \cdot])$ is abelian, $(\mathfrak{g}, [\cdot, \cdot], \rightharpoonup)$ is called a *pre-Lie algebra* [8]. More precisely, a *pre-Lie algebra* $(\mathfrak{g}, \rightharpoonup)$ is just a vector space \mathfrak{g} equipped with a linear map $\rightharpoonup : \mathfrak{g} \otimes \mathfrak{g} \rightarrow \mathfrak{g}$ such that

$$((x \rightharpoonup y) - (y \rightharpoonup x)) \rightharpoonup z = (x \rightharpoonup (y \rightharpoonup z)) - (y \rightharpoonup (x \rightharpoonup z)).$$

To any post-Lie algebra $(\mathfrak{g}, [\cdot, \cdot], \rightharpoonup)$ one can associate a Lie algebra $\mathfrak{g}_{\rightharpoonup} := (\mathfrak{g}, [\cdot, \cdot]_{\rightharpoonup})$, called the *subadjacent Lie algebra*, where the Lie bracket $[\cdot, \cdot]_{\rightharpoonup}$ is defined by

$$[x, y]_{\rightharpoonup} := (x \rightharpoonup y) - (y \rightharpoonup x) + [x, y],$$

so that (2.1)–(2.2) equivalently mean that $\mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g}), x \mapsto (x \rightharpoonup (-))$ is an action of the Lie algebra $(\mathfrak{g}, [\cdot, \cdot]_{\rightharpoonup})$ on the Lie algebra $(\mathfrak{g}, [\cdot, \cdot])$. Recently, many studies on the universal enveloping algebra of a pre-Lie algebra and a post-Lie algebra were conducted, see, e.g., [11, 26].

We also mention that the corresponding notion of *post-group* was introduced in [4]. It consists of a group (G, \cdot) equipped with a map $\dashv : G \times G \rightarrow G$ such that, for each $a \in G$, the map $a \dashv (-) : G \rightarrow G$ is an automorphism of the group (G, \cdot) , that is,

$$a \dashv (b \cdot c) = (a \dashv b) \cdot (a \dashv c).$$

Moreover, the following deformed associativity holds:

$$a \dashv (b \dashv c) = (a \cdot (a \dashv b)) \dashv c.$$

As shown in [1], the notion of post-group formalises properties of group-like elements in the completion of the enveloping algebra of a post-Lie algebra, see also [11]. The category of post-groups is proven to be isomorphic to the category of skew braces [4, Theorem 3.25].

We also recall that, given an action $\varphi : \mathfrak{g} \rightarrow \text{Der}(\mathfrak{h})$ of a Lie algebra $(\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{g}})$ on a Lie algebra $(\mathfrak{h}, [\cdot, \cdot]_{\mathfrak{h}})$, a linear map $R : \mathfrak{h} \rightarrow \mathfrak{g}$ is called a *relative Rota–Baxter operator of weight 1* on \mathfrak{h} with respect to (\mathfrak{h}, φ) if the following condition holds true for all $x, y \in \mathfrak{h}$:

$$[R(x), R(y)]_{\mathfrak{g}} = R(\varphi(R(x))y - \varphi(R(y))x + [x, y]_{\mathfrak{h}}).$$

It was shown in [3] that, given a relative Rota–Baxter operator $R : \mathfrak{h} \rightarrow \mathfrak{g}$, the Lie algebra \mathfrak{h} inherits a post-Lie algebra structure defined by

$$x \dashv y := \varphi(R(x))y \quad \text{for all } x, y \in \mathfrak{h},$$

and, given a post-Lie algebra, the identity map is a relative Rota–Baxter operator on the subadjacent Lie algebra. Similarly, given an action $\phi : G \rightarrow \text{Aut}(H)$ of a group G on a group H , a map $R : H \rightarrow G$ is called a *relative Rota–Baxter operator of weight 1* on G with respect to (H, ϕ) if

$$R(h) \cdot_G R(k) = R(h \cdot_H \phi(R(h))k) \quad \text{for all } h, k \in H.$$

This notion was introduced in [17] and its connection with skew braces was studied in [5].

Yetter–Drinfeld braces and matched pairs. The notion of *skew brace* was introduced in [14], generalising that of *brace* given in [28]. The Hopf-theoretic version of a skew brace, namely, a *Hopf brace*, was introduced in [2]. Hopf braces are equivalent to bijective 1-cocycles, see [2, Theorem 1.12]. Moreover, in the cocommutative case, Hopf braces are also equivalent to matched pairs of Hopf algebras in the sense of [24, Definition 7.2.1] which satisfy an additional relation (precisely (2.8) below), see [2, Theorem 3.3]. In [12], we have introduced Yetter–Drinfeld braces, which coincide with Hopf braces in the cocommutative case, and in [12, Theorem 3.22], we proved that the category of Yetter–Drinfeld braces is isomorphic to the category of matched pairs of actions on a Hopf algebra.

Let us recall the notions of Yetter–Drinfeld brace and matched pair of actions on a Hopf algebra. In order to do this, we first recall the definition of Yetter–Drinfeld module.

Definition 2.1. Let $(H, \bullet, 1, \Delta, \varepsilon, T)$ be a Hopf algebra. A (left-left) Yetter–Drinfeld module on H is the datum of a left H -module V with an action $\rightharpoonup: H \otimes V \rightarrow V$, which is also a left H -comodule with a coaction $\rho: V \rightarrow H \otimes V$, satisfying the following compatibility condition:

$$\rho(h \rightharpoonup v) = h_1 \bullet v_{-1} \bullet T(h_3) \otimes (h_2 \rightharpoonup v_0).$$

A morphism of Yetter–Drinfeld modules is a morphism of both left H -modules and left H -comodules. We denote by ${}^H_H\mathcal{YD}$ the category of Yetter–Drinfeld modules over H .

More generally, one can define Yetter–Drinfeld modules over a bialgebra but we will work with Hopf algebras in the following.

It is known that, see, e.g., [27], given any monoidal category $(\mathcal{M}, \otimes, \mathbf{1})$, one can consider the categories of algebras $\text{Mon}(\mathcal{M})$ and coalgebras $\text{Comon}(\mathcal{M})$ in \mathcal{M} . Once the category is braided (or even just prebraided) one can also define the categories of bialgebras $\text{Bimon}(\mathcal{M})$ and Hopf algebras $\text{Hopf}(\mathcal{M})$ in \mathcal{M} .

The category ${}^H_H\mathcal{YD}$ of Yetter–Drinfeld modules over a Hopf algebra H is monoidal and it is equipped with a prebraiding defined, for all $X, Y \in {}^H_H\mathcal{YD}$, as follows:

$$c_{X,Y}: X \otimes Y \rightarrow Y \otimes X, \quad x \otimes y \mapsto (x_{-1} \rightharpoonup y) \otimes x_0,$$

and this is bijective if the antipode of H is bijective, see, e.g., [27]. Thus, one can define Hopf algebras $\text{Hopf}({}^H_H\mathcal{YD})$ in the category ${}^H_H\mathcal{YD}$.

Definition 2.2 ([12, Definition 3.3]). A Yetter–Drinfeld brace (or YD-brace) $(H, \cdot, \bullet, 1, \Delta, \varepsilon, S, T)$ is the datum of a Hopf algebra $H^\bullet = (H, \bullet, 1, \Delta, \varepsilon, T)$, a second operation \cdot on H , and a linear map $S: H \rightarrow H$ such that

- (1) $(H, \cdot, 1, \Delta, \varepsilon, S)$ is in $\text{Hopf}({}^{H^\bullet}_H\mathcal{YD})$ with the action \rightharpoonup defined by $a \rightharpoonup b := S(a_1) \cdot (a_2 \bullet b)$, and the coaction given by Ad_L ;
- (2) if we define $a \leftarrow b := T(a_1 \rightharpoonup b_1) \bullet a_2 \bullet b_2$, the two maps $\rightharpoonup, \leftarrow$ satisfy (2.9);
- (3) the two operations \bullet and \cdot satisfy the Hopf brace compatibility:

$$a \bullet (b \cdot c) = (a_1 \bullet b) \cdot S(a_2) \cdot (a_3 \bullet c). \tag{2.3}$$

Given two Yetter–Drinfeld braces H and K , a morphism of Yetter–Drinfeld braces is a map $f: H \rightarrow K$ that is a morphism in $\text{Hopf}(\text{Vec}_{\mathbb{k}})$ between the respective Hopf algebras in $\text{Vec}_{\mathbb{k}}$, and satisfies

$$f(a \cdot_H b) = f(a) \cdot_K f(b).$$

In particular, this implies $S_K f = f S_H$, $f \rightharpoonup_H = \rightharpoonup_K (f \otimes f)$ and $(\text{Ad}_L)_K f = (f \otimes f)(\text{Ad}_L)_H$. We denote the category of Yetter–Drinfeld braces by $\mathcal{YD}\text{Br}(\text{Vec}_{\mathbb{k}})$.

Definition 2.3 (cf. [12, Definition 2.1]). Let H be a bialgebra. A matched pair of actions $(H, \rightharpoonup, \leftarrow)$ on H is the datum of a left action \rightharpoonup and a right action \leftarrow of H on itself, such that H is a left H -module coalgebra and a right H -module coalgebra with the respective

actions, and the following conditions hold for all $a, b, c \in H$:

$$a \rightharpoonup 1 = \varepsilon(a)1, \tag{2.4}$$

$$1 \leftharpoonup a = \varepsilon(a)1, \tag{2.5}$$

$$a \rightharpoonup (b \cdot c) = (a_1 \rightharpoonup b_1) \cdot ((a_2 \leftharpoonup b_2) \rightharpoonup c), \tag{2.6}$$

$$(a \cdot b) \leftharpoonup c = (a \leftharpoonup (b_1 \rightharpoonup c_1)) \cdot (b_2 \leftharpoonup c_2), \tag{2.7}$$

$$a \cdot b = (a_1 \rightharpoonup b_1) \cdot (a_2 \leftharpoonup b_2). \tag{2.8}$$

A morphism of matched pairs of actions between $(H, \rightharpoonup_H, \leftharpoonup_H)$ and $(K, \rightharpoonup_K, \leftharpoonup_K)$ is a morphism of bialgebras $H \rightarrow K$ that intertwines the two left actions and the two right actions, respectively, in the following sense:

$$f(a \rightharpoonup_H b) = f(a) \rightharpoonup_K f(b), \quad f(a \leftharpoonup_H b) = f(a) \leftharpoonup_K f(b).$$

The category of matched pairs of actions (in $\text{Vec}_{\mathbb{k}}$) is denoted by $\text{MP}(\text{Vec}_{\mathbb{k}})$.

Given a matched pairs of actions $(H, \rightharpoonup, \leftharpoonup)$ on a Hopf algebra H , the following equality is always satisfied:

$$(a_1 \rightharpoonup b_1) \otimes (a_2 \leftharpoonup b_2) = (a_2 \rightharpoonup b_2) \otimes (a_1 \leftharpoonup b_1), \tag{2.9}$$

see [12, Corollary 3.6]. Thus, matched pairs of actions on a Hopf algebra form a subclass of matched pairs of Hopf algebras in the sense of [24, Definition 7.2.1].

We also recall that in [12, Definition 4.1] we introduced the category of Yetter–Drinfeld 1-cocycles and in [12, Theorem 4.2] we proved that a subcategory of it is equivalent to the category of Yetter–Drinfeld braces. We will give more details about Yetter–Drinfeld 1-cocycles in the last section.

3. Yetter–Drinfeld post-Hopf algebras

The notion of *post-Hopf algebra* is introduced in [22, Definition 2.1] and in [22, Theorem 2.7], it is shown that the subspace of primitive elements of a post-Hopf algebra is a post-Lie algebra. Moreover, in [21, Theorem 2.13] (where [21] is the arXiv version of [22]) it was also proven that, in the cocommutative case, one can obtain a Hopf brace from a post-Hopf algebra and vice versa. Here, we introduce a generalisation of the notion of post-Hopf algebra, which we call *Yetter–Drinfeld post-Hopf algebra*, that coincides with a post-Hopf algebra in the cocommutative case. The category of Yetter–Drinfeld post-Hopf algebras will turn out to be isomorphic to the category of Yetter–Drinfeld braces given in [12].

Definition 3.1. A *Yetter–Drinfeld post-Hopf algebra* $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$, also denoted as a pair (H, \rightharpoonup) , is the datum of an algebra $(H, \cdot, 1)$, a coalgebra (H, Δ, ε) , a morphism

$S : H \rightarrow H$ such that $x_1 \cdot S(x_2) = S(x_1) \cdot x_2 = \varepsilon(x)1_H$ for all $x \in H$ and a morphism of coalgebras $\rightharpoonup : H \otimes H \rightarrow H$ satisfying the following equalities, for any $x, y, z \in H$:

$$x \rightharpoonup (y \cdot z) = (x_1 \rightharpoonup y) \cdot (x_2 \rightharpoonup z), \tag{3.1}$$

$$x \rightharpoonup (y \rightharpoonup z) = (x_1 \cdot (x_2 \rightharpoonup y)) \rightharpoonup z. \tag{3.2}$$

Moreover, the morphism $\alpha_{\rightharpoonup} : H \rightarrow \text{End}(H)$ defined by

$$(\alpha_{\rightharpoonup}x)(y) := x \rightharpoonup y \quad \text{for all } x, y \in H,$$

is convolution invertible in $\text{Hom}(H, \text{End}(H))$, i.e., there exists a unique morphism $\beta_{\rightharpoonup} : H \rightarrow \text{End}(H)$ such that

$$(\alpha_{\rightharpoonup}x_1) \circ (\beta_{\rightharpoonup}x_2) = (\beta_{\rightharpoonup}x_1) \circ (\alpha_{\rightharpoonup}x_2) = \varepsilon(x)\text{Id}_H \quad \text{for all } x \in H. \tag{3.3}$$

Moreover, ε is a morphism of algebras, i.e., $\varepsilon(a \cdot b) = \varepsilon(a)\varepsilon(b)$ and $\varepsilon(1_H) = 1_{\mathbb{k}}$, $\Delta(1_H) = 1_H \otimes 1_H$ and the following compatibility condition between Δ and \cdot holds true:

$$\Delta(x \cdot y) = (x_1 \cdot (\alpha_{\rightharpoonup}x_2(\beta_{\rightharpoonup}x_4(y_1)))) \otimes (x_3 \cdot y_2). \tag{3.4}$$

Finally, by defining

$$x \bullet_{\rightharpoonup} y := x_1 \cdot (x_2 \rightharpoonup y), \tag{3.5}$$

$$S_{\rightharpoonup}(x) := \beta_{\rightharpoonup}x_1(S(x_2)) \tag{3.6}$$

and

$$x \leftarrow y := S_{\rightharpoonup}(x_1 \rightharpoonup y_1) \bullet_{\rightharpoonup} x_2 \bullet_{\rightharpoonup} y_2, \tag{3.7}$$

we require that S_{\rightharpoonup} is anti-comultiplicative, i.e., it satisfies $\Delta S_{\rightharpoonup}(x) = S_{\rightharpoonup}(x_2) \otimes S_{\rightharpoonup}(x_1)$, and the pair $(\rightharpoonup, \leftarrow)$ satisfies (2.9).

A morphism of Yetter–Drinfeld post-Hopf algebras from (H, \rightharpoonup) to (H', \rightharpoonup') is a morphism of algebras and coalgebras $g : H \rightarrow H'$ satisfying

$$g(x \rightharpoonup y) = g(x) \rightharpoonup' g(y) \quad \text{for all } x, y \in H. \tag{3.8}$$

We denote the category of Yetter–Drinfeld post-Hopf algebras and morphisms of Yetter–Drinfeld post-Hopf algebras by $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$.

Remark 3.2. We use notations $\bullet_{\rightharpoonup}$ and S_{\rightharpoonup} , similarly to those used in [22, Theorem 2.5]. Let us observe that one can recover \rightharpoonup as

$$x \rightharpoonup y = S(x_1) \cdot x_2 \cdot (x_3 \rightharpoonup y) \stackrel{(3.5)}{=} S(x_1) \cdot (x_2 \bullet_{\rightharpoonup} y) \tag{3.9}$$

and rewrite (3.2) as

$$x \rightharpoonup (y \rightharpoonup z) = (x \bullet_{\rightharpoonup} y) \rightharpoonup z. \tag{3.10}$$

Remark 3.3. Suppose H is cocommutative. Then, we obtain

$$\begin{aligned} \Delta(x \cdot y) &\stackrel{(3.4)}{=} (x_1 \cdot (\alpha_{\rightarrow} x_2 (\beta_{\rightarrow} x_4 (y_1)))) \otimes (x_3 \cdot y_2) \\ &= (x_1 \cdot (\alpha_{\rightarrow} x_2 (\beta_{\rightarrow} x_3 (y_1)))) \otimes (x_4 \cdot y_2) \\ &\stackrel{(3.3)}{=} (x_1 \cdot y_1) \otimes (x_2 \cdot y_2), \end{aligned}$$

hence, H becomes a standard Hopf algebra and (H, \rightarrow) is a post-Hopf algebra, see [22, Definition 2.1]. Moreover, clearly, (2.9) is automatically satisfied and, as we will observe later, S_{\rightarrow} becomes comultiplicative.

One can prove that the same properties given in [22, Lemma 2.4] hold for Yetter–Drinfeld post-Hopf algebras. We include the proof for the sake of completeness, showing that there are no problems with this more general definition.

Lemma 3.4. *Let (H, \rightarrow) be a Yetter–Drinfeld post-Hopf algebra. Then, the following properties are satisfied, for all $x, y \in H$:*

$$x \rightarrow 1 = \varepsilon(x)1, \tag{3.11}$$

$$1 \rightarrow x = x, \tag{3.12}$$

$$S(x \rightarrow y) = x \rightarrow S(y). \tag{3.13}$$

Proof. Since \rightarrow is a morphism of coalgebras, we have

$$\begin{aligned} x \rightarrow 1 &= (x_1 \rightarrow 1)\varepsilon(x_2 \rightarrow 1) \\ &= (x_1 \rightarrow 1) \cdot (x_2 \rightarrow 1) \cdot S(x_3 \rightarrow 1) \stackrel{(3.1)}{=} (x_1 \rightarrow 1) \cdot S(x_2 \rightarrow 1) \\ &= \varepsilon(x \rightarrow 1)1 = \varepsilon(x)1. \end{aligned}$$

By (3.3), we have $(\alpha_{\rightarrow} 1) \circ (\beta_{\rightarrow} 1) = (\beta_{\rightarrow} 1) \circ (\alpha_{\rightarrow} 1) = \text{Id}_H$, so $\alpha_{\rightarrow} 1$ is an automorphism of H with inverse given by $\beta_{\rightarrow} 1$. Moreover, we have

$$(\alpha_{\rightarrow} 1)^2(x) = 1 \rightarrow (1 \rightarrow x) \stackrel{(3.2)}{=} (1 \rightarrow 1) \rightarrow x \stackrel{(3.11)}{=} 1 \rightarrow x = (\alpha_{\rightarrow} 1)(x),$$

hence, $1 \rightarrow x = (\alpha_{\rightarrow} 1)(x) = x$ (and so, also $\beta_{\rightarrow} 1 = \text{Id}$). Finally, we have

$$\begin{aligned} S(x \rightarrow y) &= S(x_1 \rightarrow y_1)\varepsilon(x_2)\varepsilon(y_2) \stackrel{(3.11)}{=} S(x_1 \rightarrow y_1) \cdot (x_2 \rightarrow \varepsilon(y_2)1) \\ &= S(x_1 \rightarrow y_1) \cdot (x_2 \rightarrow y_2 \cdot S(y_3)) \stackrel{(3.1)}{=} S(x_1 \rightarrow y_1) \cdot (x_2 \rightarrow y_2) \cdot (x_3 \rightarrow S(y_3)) \\ &= \varepsilon(x_1 \rightarrow y_1)(x_2 \rightarrow S(y_2)) = x \rightarrow S(y). \quad \blacksquare \end{aligned}$$

Remark 3.5. Given a Yetter–Drinfeld post-Hopf algebra (H, \rightarrow) we know by definition that α_{\rightarrow} has convolution inverse given by β_{\rightarrow} . We prove that $\alpha_{\rightarrow} S_{\rightarrow}$ is a right convolution inverse of α_{\rightarrow} in order to obtain $\beta_{\rightarrow} = \alpha_{\rightarrow} S_{\rightarrow}$. We compute

$$\begin{aligned} \alpha_{\rightarrow} x_1 (\alpha_{\rightarrow} S_{\rightarrow} (x_2)(a)) &= x_1 \rightarrow (S_{\rightarrow} (x_2) \rightarrow a) \stackrel{(3.2)}{=} (x_1 \cdot (x_2 \rightarrow S_{\rightarrow} (x_3))) \rightarrow a \\ &\stackrel{(3.6)}{=} (x_1 \cdot (\alpha_{\rightarrow} x_2 (\beta_{\rightarrow} x_3 (S(x_4)))))) \rightarrow a \stackrel{(3.3)}{=} (x_1 \cdot S(x_2)) \rightarrow a \\ &\stackrel{(3.12)}{=} \varepsilon(x)a, \end{aligned}$$

thus, we get

$$\alpha_{\rightarrow} S_{\rightarrow} = \beta_{\rightarrow}. \tag{3.14}$$

Remark 3.6. Notice that, from (3.10) and (3.12), we have that H is an object in $H_{\rightarrow} \mathfrak{M}$, the category of left H_{\rightarrow} -modules. Moreover, from (3.1) and (3.11), we have that $(H, \cdot, 1)$ is in $\text{Mon}(H_{\rightarrow} \mathfrak{M})$. Recall also that, by definition of Yetter–Drinfeld post-Hopf algebra, \rightarrow is a morphism of coalgebras, i.e., (H, Δ, ε) is in $\text{Comon}(H_{\rightarrow} \mathfrak{M})$.

We can prove the following result.

Lemma 3.7. *The following equations are satisfied, for all $x \in H$:*

$$\Delta \circ (\alpha_{\rightarrow} x) = (\alpha_{\rightarrow} x_1 \otimes \alpha_{\rightarrow} x_2) \circ \Delta, \tag{3.15}$$

$$\Delta \circ (\beta_{\rightarrow} x) = (\beta_{\rightarrow} x_2 \otimes \beta_{\rightarrow} x_1) \circ \Delta, \tag{3.16}$$

$$(\alpha_{\rightarrow} x) \circ m. = m. \circ (\alpha_{\rightarrow} x_1 \otimes \alpha_{\rightarrow} x_2), \tag{3.17}$$

$$(\beta_{\rightarrow} x) \circ m. = m. \circ (\beta_{\rightarrow} x_2 \otimes \beta_{\rightarrow} x_1). \tag{3.18}$$

Proof. First, we compute

$$\begin{aligned} \Delta((\alpha_{\rightarrow} x)(y)) &= \Delta(x \rightarrow y) = (x_1 \rightarrow y_1) \otimes (x_2 \rightarrow y_2) = (\alpha_{\rightarrow} x_1)(y_1) \otimes (\alpha_{\rightarrow} x_2)(y_2) \\ &= (\alpha_{\rightarrow} x_1 \otimes \alpha_{\rightarrow} x_2) \Delta(y), \end{aligned}$$

hence, (3.15) holds true. Thus, we have

$$\begin{aligned} (\alpha_{\rightarrow} x_1 \otimes \alpha_{\rightarrow} x_3)(\beta_{\rightarrow} x_2 \otimes \beta_{\rightarrow} x_4) \Delta(y) &\stackrel{(3.3)}{=} \varepsilon(x_1) \varepsilon(x_2) \Delta(y) = \Delta(\varepsilon(x) y) \\ &\stackrel{(3.3)}{=} \Delta(\alpha_{\rightarrow} x_1(\beta_{\rightarrow} x_2(y))) \\ &\stackrel{(3.15)}{=} (\alpha_{\rightarrow} x_1 \otimes \alpha_{\rightarrow} x_2) \Delta(\beta_{\rightarrow} x_3(y)), \end{aligned} \tag{3.19}$$

hence,

$$\begin{aligned} (\text{Id} \otimes \alpha_{\rightarrow} x_1) \Delta(\beta_{\rightarrow} x_2(y)) &\stackrel{(3.3)}{=} (\beta_{\rightarrow} x_1 \alpha_{\rightarrow} x_2 \otimes \alpha_{\rightarrow} x_3) \Delta(\beta_{\rightarrow} x_4(y)) \\ &\stackrel{(3.19)}{=} (\beta_{\rightarrow} x_1 \alpha_{\rightarrow} x_2 \otimes \alpha_{\rightarrow} x_4) (\beta_{\rightarrow} x_3 \otimes \beta_{\rightarrow} x_5) \Delta(y) \\ &\stackrel{(3.3)}{=} (\text{Id} \otimes \alpha_{\rightarrow} x_2) (\beta_{\rightarrow} x_1 \otimes \beta_{\rightarrow} x_3) \Delta(y), \end{aligned} \tag{3.20}$$

and then, we obtain

$$\begin{aligned} \Delta((\beta_{\rightarrow} x)(y)) &\stackrel{(3.3)}{=} (\text{Id} \otimes \beta_{\rightarrow} x_1 \alpha_{\rightarrow} x_2) \Delta(\beta_{\rightarrow} x_3(y)) \\ &\stackrel{(3.20)}{=} (\text{Id} \otimes \beta_{\rightarrow} x_1 \alpha_{\rightarrow} x_3) (\beta_{\rightarrow} x_2 \otimes \beta_{\rightarrow} x_4) \Delta(y) \\ &\stackrel{(3.3)}{=} (\beta_{\rightarrow} x_2 \otimes \beta_{\rightarrow} x_1) \Delta(y). \end{aligned}$$

Hence, (3.16) is satisfied. Moreover, we have

$$(\alpha_{\rightarrow} x) m.(a \otimes b) = x \rightarrow (a \cdot b) \stackrel{(3.1)}{=} (x_1 \rightarrow a) \cdot (x_2 \rightarrow b) = m.(\alpha_{\rightarrow} x_1 \otimes \alpha_{\rightarrow} x_2)(a \otimes b),$$

so (3.17) holds true. The proof of (3.18) is the dual of the proof of (3.16). ■

Remark 3.8. Notice that

$$\begin{aligned} \Delta(S_{\rightarrow}(x)) &\stackrel{(3.6)}{=} \Delta\beta_{\rightarrow x_1}(S(x_2)) \\ &\stackrel{(3.16)}{=} (\beta_{\rightarrow x_2} \otimes \beta_{\rightarrow x_1})\Delta(S(x_3)) = \beta_{\rightarrow x_2}(S(x_4)) \otimes \beta_{\rightarrow x_1}(S(x_3)), \end{aligned}$$

hence, the fact that S_{\rightarrow} is anti-comultiplicative reads in terms on β_{\rightarrow} as follows:

$$\beta_{\rightarrow x_2}(S(x_4)) \otimes \beta_{\rightarrow x_1}(S(x_3)) = \beta_{\rightarrow x_3}(S(x_4)) \otimes \beta_{\rightarrow x_1}(S(x_2)).$$

Remark 3.9. Notice that

$$\beta_{\rightarrow x}(1) = \beta_{\rightarrow x_1}(\varepsilon(x_2)1) \stackrel{(3.11)}{=} \beta_{\rightarrow x_1}(\alpha_{\rightarrow x_2}(1)) \stackrel{(3.3)}{=} \varepsilon(x)1,$$

and then,

$$\beta_{\rightarrow x_2}(S(x_3)) \cdot \beta_{\rightarrow x_1}(x_4) \stackrel{(3.18)}{=} \beta_{\rightarrow x_1}(S(x_2) \cdot x_3) = \beta_{\rightarrow x}(1) = \varepsilon(x)1 \quad (3.21)$$

for all $x \in H$. Observe also that if H is cocommutative, we have $\Delta \circ (\beta_{\rightarrow x}) = (\beta_{\rightarrow x_1} \otimes \beta_{\rightarrow x_2})\Delta$ and $(\beta_{\rightarrow x}) \circ m. = m. \circ (\beta_{\rightarrow x_1} \otimes \beta_{\rightarrow x_2})$ for all $x \in H$, and moreover, S_{\rightarrow} is comultiplicative.

The following result generalises [22, Theorem 2.5], removing the hypothesis of cocommutativity.

Proposition 3.10. *Let (H, \rightarrow) be a Yetter–Drinfeld post-Hopf algebra and define \bullet_{\rightarrow} and S_{\rightarrow} as in (3.5) and (3.6), respectively. Then, $H_{\rightarrow} := (H, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S_{\rightarrow})$ is a Hopf algebra. We call H_{\rightarrow} the subadjacent Hopf algebra in analogy with [22].*

Proof. By definition of Yetter–Drinfeld post-Hopf algebra, we already know that $\Delta(1_H) = 1_H \otimes 1_H$ and $\varepsilon(1_H) = 1_{\mathbb{k}}$. We show that Δ and ε are multiplicative with respect to \bullet_{\rightarrow} :

$$\begin{aligned} \Delta(x \bullet_{\rightarrow} y) &\stackrel{(3.5)}{=} \Delta(x_1 \cdot (x_2 \rightarrow y)) \stackrel{(3.4)}{=} (x_1 \cdot (\alpha_{\rightarrow x_2}(\beta_{\rightarrow x_4}(x_5 \rightarrow y_1)))) \otimes (x_3 \cdot (x_6 \rightarrow y_2)) \\ &= x_1 \cdot (\alpha_{\rightarrow x_2}(\beta_{\rightarrow x_4}(\alpha_{\rightarrow x_5}(y_1)))) \otimes (x_3 \cdot (x_6 \rightarrow y_2)) \\ &\stackrel{(3.3)}{=} (x_1 \cdot (x_2 \rightarrow y_1)) \otimes (x_3 \cdot (x_4 \rightarrow y_2)) \\ &= (x_1 \bullet_{\rightarrow} y_1) \otimes (x_2 \bullet_{\rightarrow} y_2) \end{aligned}$$

and

$$\varepsilon(x \bullet_{\rightarrow} y) \stackrel{(3.5)}{=} \varepsilon(x_1 \cdot (x_2 \rightarrow y)) = \varepsilon(x_1)\varepsilon(x_2 \rightarrow y) = \varepsilon(x_1)\varepsilon(x_2)\varepsilon(y) = \varepsilon(x)\varepsilon(y).$$

We show that $(H, \bullet_{\rightarrow}, 1)$ is an algebra, so that $(H, \bullet_{\rightarrow}, 1, \Delta, \varepsilon)$ is a bialgebra. We have

$$1 \bullet_{\rightarrow} x = 1 \cdot (1 \rightarrow x) \stackrel{(3.12)}{=} 1 \cdot x = x, \quad x \bullet_{\rightarrow} 1 = x_1 \cdot (x_2 \rightarrow 1) \stackrel{(3.11)}{=} x_1 \cdot \varepsilon(x_2)1 = x,$$

and also,

$$\begin{aligned}
 x \bullet_{\rightarrow} (y \bullet_{\rightarrow} z) &= x_1 \cdot (x_2 \rightarrow (y_1 \cdot (y_2 \rightarrow z))) \\
 &\stackrel{(3.1)}{=} x_1 \cdot (x_2 \rightarrow y_1) \cdot (x_3 \rightarrow (y_2 \rightarrow z)) \\
 &\stackrel{(3.2)}{=} (x_1 \cdot (x_2 \rightarrow y_1)) \cdot ((x_3 \cdot (x_4 \rightarrow y_2)) \rightarrow z) \\
 &= (x_1 \bullet_{\rightarrow} y_1) \cdot ((x_2 \bullet_{\rightarrow} y_2) \rightarrow z) \\
 &= (x \bullet_{\rightarrow} y) \bullet_{\rightarrow} z,
 \end{aligned}$$

where the last equality follows from the fact that Δ is multiplicative with respect to \bullet_{\rightarrow} . Finally, we prove that S_{\rightarrow} is an antipode. We compute

$$\begin{aligned}
 x_1 \bullet_{\rightarrow} S_{\rightarrow}(x_2) &\stackrel{(3.5)}{=} x_1 \cdot (x_2 \rightarrow S_{\rightarrow}(x_3)) \\
 &\stackrel{(3.6)}{=} x_1 \cdot (\alpha_{\rightarrow} x_2 (\beta_{\rightarrow} x_3 (S(x_4)))) = x_1 \cdot S(x_2) = \varepsilon(x)1.
 \end{aligned}$$

Moreover, since S_{\rightarrow} is anti-comultiplicative, we obtain

$$\begin{aligned}
 S_{\rightarrow}(x_1) \bullet_{\rightarrow} x_2 &\stackrel{(3.5)}{=} S_{\rightarrow}(x_1)_1 \cdot (S_{\rightarrow}(x_1)_2 \rightarrow x_2) \\
 &= S_{\rightarrow}(x_2) \cdot (S_{\rightarrow}(x_1) \rightarrow x_3) \\
 &= \beta_{\rightarrow} x_2 (S(x_3)) \cdot (\alpha_{\rightarrow} S_{\rightarrow} x_1(x_4)) \\
 &\stackrel{(3.14)}{=} \beta_{\rightarrow} x_2 (S(x_3)) \cdot \beta_{\rightarrow} x_1(x_4) \\
 &\stackrel{(3.21)}{=} \varepsilon(x)1,
 \end{aligned}$$

hence, S_{\rightarrow} is an antipode and $(H, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S_{\rightarrow})$ is a Hopf algebra. ■

Remark 3.11. Let us observe that since S_{\rightarrow} is an antipode, one can recover \cdot as

$$a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_2) \rightarrow b) = a_1 \cdot (a_2 \rightarrow (S_{\rightarrow}(a_3) \rightarrow b)) = a_1 \cdot (a_2 \bullet_{\rightarrow} S_{\rightarrow}(a_3) \rightarrow b) = a \cdot b. \tag{3.22}$$

The relation between post-Hopf algebras and post-Lie algebras given in [22, Theorem 2.7] still remains true for Yetter–Drinfeld post-Hopf algebras as it is written in the following result.

Proposition 3.12. *Let (H, \rightarrow) be a Yetter–Drinfeld post-Hopf algebra. Then, its subspace of primitive elements $P(H)$ is a post-Lie algebra.*

The proof is exactly the same of [22, Theorem 2.7]. Indeed, this only involves the equations (3.1), (3.2), (3.11), and (3.12) and the fact that \rightarrow is a morphism of coalgebras. This shows that the notion of Yetter–Drinfeld post-Hopf algebra is the natural one in the non-cocommutative setting.

Remark 3.13. A kind of converse of the previous result holds. Indeed, it is known that, given a post-Lie algebra $(\mathfrak{g}, [\cdot, \cdot], \rightharpoonup)$, the map \rightharpoonup can be extended to its universal enveloping algebra and induces a subadjacent Hopf algebra structure isomorphic to the universal enveloping algebra of the subadjacent Lie algebra $\mathfrak{g}_{\rightharpoonup}$, see [11, Proposition 3.1 and Theorem 3.4]. In this case, since universal enveloping algebras are cocommutative, the definition of Yetter–Drinfeld post-Hopf algebra coincides with that of post-Hopf algebra and one can refer to [22, Theorem 2.8]. Observe also that one can easily obtain examples of cocommutative post-Hopf algebras which do not come from Lie algebras considering group algebras with action coming from Rota–Baxter operators on the base group, as it is done in [22, Example 2.11].

Remark 3.14. In [22, Definition 2.3], a post-Hopf algebra (H, \rightharpoonup) is called a *pre-Hopf algebra* if H is commutative. In this case, $P(H)$ results to be a pre-Lie algebra since $[x, y] = 0$ for all $x, y \in P(H)$. We will introduce the notion of *Yetter–Drinfeld pre-Hopf algebra* later and it will coincide with the notion of pre-Hopf algebra in the cocommutative case.

3.1. Yetter–Drinfeld post-Hopf algebras and Yetter–Drinfeld braces

We are ready to prove that the category of Yetter–Drinfeld post-Hopf algebras is isomorphic to the category of Yetter–Drinfeld braces. First, we show some preliminary results.

Lemma 3.15. *Let (H, \rightharpoonup) be a Yetter–Drinfeld post-Hopf algebra and define \leftarrow as in (3.7). Then, we have $\varepsilon(a \leftarrow b) = \varepsilon(a)\varepsilon(b)$ and $(a_1 \rightharpoonup b_1) \bullet_{\rightharpoonup} (a_2 \leftarrow b_2) = a \bullet_{\rightharpoonup} b$.*

Proof. We compute

$$\begin{aligned} \varepsilon(a \leftarrow b) &= \varepsilon(S_{\rightharpoonup}(a_1 \rightharpoonup b_1) \bullet_{\rightharpoonup} a_2 \bullet_{\rightharpoonup} b_2) = \varepsilon(S_{\rightharpoonup}(a_1 \rightharpoonup b_1))\varepsilon(a_2)\varepsilon(b_2) \\ &= \varepsilon(a \rightharpoonup b) = \varepsilon(a)\varepsilon(b), \end{aligned}$$

and also,

$$\begin{aligned} (a_1 \rightharpoonup b_1) \bullet_{\rightharpoonup} (a_2 \leftarrow b_2) &= (a_1 \rightharpoonup b_1) \bullet_{\rightharpoonup} S_{\rightharpoonup}(a_2 \rightharpoonup b_2) \bullet_{\rightharpoonup} a_3 \bullet_{\rightharpoonup} b_3 \\ &= \varepsilon(a_1 \rightharpoonup b_1)a_2 \bullet_{\rightharpoonup} b_2 = a \bullet_{\rightharpoonup} b. \quad \blacksquare \end{aligned}$$

Remark 3.16. By [12, Lemma 3.5], we have that (2.9) is equivalent to \leftarrow being a morphism of coalgebras, i.e., $\Delta(a \leftarrow b) = (a_1 \leftarrow b_1) \otimes (a_2 \leftarrow b_2)$ and also to

$$(a_1 \rightharpoonup b_1) \bullet_{\rightharpoonup} (a_3 \leftarrow b_3) \otimes (a_2 \rightharpoonup b_2) = (a_1 \bullet_{\rightharpoonup} b_1) \otimes (a_2 \rightharpoonup b_2). \tag{3.23}$$

Let us also observe that, from the definition of \leftarrow and the fact that S_{\rightharpoonup} is the antipode of H_{\rightharpoonup} , we obtain

$$S_{\rightharpoonup}(a \rightharpoonup b) = (a_1 \leftarrow b_1) \bullet_{\rightharpoonup} S_{\rightharpoonup}(b_2) \bullet_{\rightharpoonup} S_{\rightharpoonup}(a_2). \tag{3.24}$$

Now, we show that a Yetter–Drinfeld post-Hopf algebra is automatically a Hopf monoid in the category of (left-left) Yetter–Drinfeld modules over H_{\rightarrow} .

Theorem 3.17. *Given a Yetter–Drinfeld post-Hopf algebra (H, \rightarrow) one has $(H, \rightarrow, \text{Ad}_L)$ is in ${}^{H_{\rightarrow}}\mathcal{YD}$. Moreover, $(H, \cdot, 1, \Delta, \varepsilon, S)$ is in $\text{Hopf}({}^{H_{\rightarrow}}\mathcal{YD})$.*

Proof. Given the left adjoint coaction $\text{Ad}_L : H \rightarrow H_{\rightarrow} \otimes H, a \mapsto a_1 \bullet_{\rightarrow} S_{\rightarrow}(a_3) \otimes a_2$ and the action $\rightarrow : H_{\rightarrow} \otimes H \rightarrow H$, we verify the compatibility condition

$$\begin{aligned} \text{Ad}_L(a \rightarrow b) &= (a_1 \rightarrow b_1) \bullet_{\rightarrow} S_{\rightarrow}(a_3 \rightarrow b_3) \otimes (a_2 \rightarrow b_2) \\ &\stackrel{(3.24)}{=} (a_1 \rightarrow b_1) \bullet_{\rightarrow} (a_3 \leftarrow b_3) \bullet_{\rightarrow} S_{\rightarrow}(b_4) \bullet_{\rightarrow} S_{\rightarrow}(a_4) \otimes (a_2 \rightarrow b_2) \\ &\stackrel{(3.23)}{=} a_1 \bullet_{\rightarrow} b_1 \bullet_{\rightarrow} S_{\rightarrow}(b_3) \bullet_{\rightarrow} S_{\rightarrow}(a_3) \otimes (a_2 \rightarrow b_2). \end{aligned}$$

Hence, we obtain the braiding operator $\sigma_{H,H}^{\mathcal{YD}} : H \otimes H \rightarrow H \otimes H$ given by

$$\sigma_{H,H}^{\mathcal{YD}} : a \otimes b \mapsto (a_1 \bullet_{\rightarrow} S_{\rightarrow}(a_3) \rightarrow b) \otimes a_2.$$

Let us observe that

$$\begin{aligned} (a_1 \bullet_{\rightarrow} S_{\rightarrow}(a_3) \rightarrow b) \otimes a_2 &= (a_1 \rightarrow (S_{\rightarrow}(a_3) \rightarrow b)) \otimes a_2 \\ &= \alpha_{\rightarrow} a_1 (\alpha_{\rightarrow} S_{\rightarrow}(a_3)(b)) \otimes a_2 \\ &\stackrel{(3.14)}{=} \alpha_{\rightarrow} a_1 (\beta_{\rightarrow} a_3(b)) \otimes a_2. \end{aligned}$$

We compute

$$\begin{aligned} (m \cdot \otimes m.) (\text{Id}_H \otimes \sigma_{H,H}^{\mathcal{YD}} \otimes \text{Id}_H) (\Delta \otimes \Delta) (a \otimes b) &= (a_1 \cdot (\alpha_{\rightarrow} a_2 (\beta_{\rightarrow} a_4(b_1)))) \otimes (a_3 \cdot b_2) \\ &\stackrel{(3.4)}{=} \Delta m.(a \otimes b), \end{aligned}$$

thus, since $\varepsilon(a \cdot b) = \varepsilon(a)\varepsilon(b)$, $\varepsilon(1_H) = 1_{\mathbb{k}}$ and $\Delta(1_H) = 1_H \otimes 1_H$ hold by definition of Yetter–Drinfeld post-Hopf algebra, the compatibility condition for a bialgebra in ${}^{H_{\rightarrow}}\mathcal{YD}$ is satisfied. By Remark 3.6, we already know that $(H, \cdot, 1)$ is in $\text{Mon}(H_{\rightarrow}\mathfrak{M})$ and (H, Δ, ε) is in $\text{Comon}(H_{\rightarrow}\mathfrak{M})$. Moreover, clearly, u, Δ and ε are morphisms in ${}^{H_{\rightarrow}}\mathfrak{M}$ with respect to the adjoint coaction Ad_L . Hence, we show that also $m.$ is left colinear in order to obtain that $(H, \cdot, 1, \Delta, \varepsilon)$ is in $\text{Bimon}({}^{H_{\rightarrow}}\mathcal{YD})$. Let us first observe that

$$\begin{aligned} (\text{Id} \otimes \Delta) \Delta(a \cdot b) &= (a_1 \cdot (\alpha_{\rightarrow} a_2 (\beta_{\rightarrow} a_4(b_1)))) \otimes \Delta(a_3 \cdot b_2) \\ &= (a_1 \cdot (\alpha_{\rightarrow} a_2 (\beta_{\rightarrow} a_7(b_1)))) \otimes (a_3 \cdot (\alpha_{\rightarrow} a_4 (\beta_{\rightarrow} a_6(b_2)))) \otimes (a_5 \cdot b_3). \end{aligned}$$

From (3.23), we obtain that

$$\begin{aligned} (S_{\rightarrow}(a_3) \rightarrow b_1) \bullet_{\rightarrow} (S_{\rightarrow}(a_1) \leftarrow b_3) \otimes (S_{\rightarrow}(a_2) \rightarrow b_2) \\ = (S_{\rightarrow}(a_2) \bullet_{\rightarrow} b_1) \otimes (S_{\rightarrow}(a_1) \rightarrow b_2), \end{aligned}$$

which is equivalent to

$$\begin{aligned} & (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_5) \rightarrow b_1) \bullet_{\rightarrow} (S_{\rightarrow}(a_3) \leftarrow b_3) \bullet_{\rightarrow} S_{\rightarrow}(b_4)) \otimes (a_2 \bullet_{\rightarrow} (S_{\rightarrow}(a_4) \rightarrow b_2)) \\ &= (a_1 \bullet_{\rightarrow} S_{\rightarrow}(a_4) \bullet_{\rightarrow} b_1 \bullet_{\rightarrow} S_{\rightarrow}(b_3)) \otimes (a_2 \bullet_{\rightarrow} (S_{\rightarrow}(a_3) \rightarrow b_2)) \\ &\stackrel{(3.22)}{=} (a_1 \bullet_{\rightarrow} S_{\rightarrow}(a_3) \bullet_{\rightarrow} b_1 \bullet_{\rightarrow} S_{\rightarrow}(b_3)) \otimes (a_2 \cdot b_2). \end{aligned}$$

Moreover, we have

$$\begin{aligned} & (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_5) \rightarrow b_1) \bullet_{\rightarrow} (S_{\rightarrow}(a_3) \leftarrow b_3) \bullet_{\rightarrow} S_{\rightarrow}(b_4)) \otimes (a_2 \bullet_{\rightarrow} (S_{\rightarrow}(a_4) \rightarrow b_2)) \\ &\stackrel{(3.7)}{=} (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_6) \rightarrow b_1) \bullet_{\rightarrow} S_{\rightarrow}(S_{\rightarrow}(a_4) \rightarrow b_3) \bullet_{\rightarrow} S_{\rightarrow}(a_3) \bullet_{\rightarrow} b_4 \bullet_{\rightarrow} S_{\rightarrow}(b_5)) \\ &\quad \otimes (a_2 \bullet_{\rightarrow} (S_{\rightarrow}(a_5) \rightarrow b_2)) \\ &= (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_6) \rightarrow b_1) \bullet_{\rightarrow} S_{\rightarrow}(S_{\rightarrow}(a_4) \rightarrow b_3) \bullet_{\rightarrow} S_{\rightarrow}(a_3)) \\ &\quad \otimes (a_2 \bullet_{\rightarrow} (S_{\rightarrow}(a_5) \rightarrow b_2)) \\ &= (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_6) \rightarrow b_1) \bullet_{\rightarrow} S_{\rightarrow}(a_3 \bullet_{\rightarrow} (S_{\rightarrow}(a_4) \rightarrow b_3))) \\ &\quad \otimes (a_2 \bullet_{\rightarrow} (S_{\rightarrow}(a_5) \rightarrow b_2)) \\ &= (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_2) \rightarrow b))_1 \bullet_{\rightarrow} S_{\rightarrow}((a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_2) \rightarrow b))_3) \\ &\quad \otimes (a_1 \bullet_{\rightarrow} (S_{\rightarrow}(a_2) \rightarrow b))_2 \\ &\stackrel{(3.22)}{=} (a \cdot b)_1 \bullet_{\rightarrow} S_{\rightarrow}((a \cdot b)_3) \otimes (a \cdot b)_2, \end{aligned}$$

hence, we obtain

$$(a \cdot b)_1 \bullet_{\rightarrow} S_{\rightarrow}((a \cdot b)_3) \otimes (a \cdot b)_2 = (a_1 \bullet_{\rightarrow} S_{\rightarrow}(a_3) \bullet_{\rightarrow} b_1 \bullet_{\rightarrow} S_{\rightarrow}(b_3)) \otimes (a_2 \cdot b_2),$$

i.e., m is left colinear. Thus, $(H, \cdot, 1, \Delta, \varepsilon)$ is in $\text{Bimon}(\overset{H}{\leftarrow} \mathcal{YD})$. Finally, we already know that S satisfies $a_1 \cdot S(a_2) = \varepsilon(a)1_H = S(a_1) \cdot a_2$ for all $a \in H$, hence, $(H, \cdot, 1, \Delta, \varepsilon, S)$ is in $\text{Hopf}(\overset{H}{\leftarrow} \mathcal{YD})$ by [12, Proposition 3.9]. \blacksquare

Remark 3.18. If H is cocommutative, the braiding $\sigma_{H,H}^{\mathcal{YD}} : a \otimes b \mapsto \alpha_{\rightarrow} a_1(\beta_{\rightarrow} a_3(b)) \otimes a_2$ becomes the canonical flip $\tau_{H,H}$ and $(H, \cdot, 1, \Delta, \varepsilon, S)$ becomes a standard Hopf algebra.

We can prove the following result.

Proposition 3.19. *Let $(H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow)$ be a Yetter–Drinfeld post-Hopf algebra. Then, the set of group-like elements $G(H)$ is a post-group.*

Proof. We know that $\Delta(1_H) = 1_H \otimes 1_H$ and $\varepsilon(1_H) = 1_{\mathbb{k}}$, so $1_H \in G(H)$. In order to obtain that $(G(H), \cdot)$ is a group, we need to prove that $G(H)$ is closed under \cdot and S . Indeed, we already know that $x \cdot S(x) = S(x) \cdot x = 1_H$ for all $x \in G(H)$. Since $\alpha_{\rightarrow} : H \rightarrow \text{End}(H)$ is convolution invertible with convolution inverse β_{\rightarrow} , we obtain $(\alpha_{\rightarrow} x) \circ (\beta_{\rightarrow} x) = (\beta_{\rightarrow} x) \circ (\alpha_{\rightarrow} x) = \text{Id}$ for all $x \in G(H)$ and then, for all $x, y \in G(H)$, we have

$$\Delta(x \cdot y) \stackrel{(3.4)}{=} (x \cdot (\alpha_{\rightarrow} x(\beta_{\rightarrow} x(y)))) \otimes (x \cdot y) = (x \cdot y) \otimes (x \cdot y), \quad \varepsilon(x \cdot y) = \varepsilon(x)\varepsilon(y) = 1_{\mathbb{k}},$$

hence, $x \cdot y \in G(H)$. Moreover, by Theorem 3.17, we know that $(H, \cdot, 1, \Delta, \varepsilon, S)$ is in $\text{Hopf}({}_{H\leftarrow}^H\mathcal{YD})$, hence, S satisfies $(S \otimes S)\sigma_{H,H}^{\mathcal{YD}}\Delta = \Delta S$ and $\varepsilon S = \varepsilon$. Hence, given $x \in G(H)$, we obtain

$$(S \otimes S)\sigma_{H,H}^{\mathcal{YD}}\Delta(x) = (S \otimes S)\sigma_{H,H}^{\mathcal{YD}}(x \otimes x) = (S \otimes S)(\alpha_{\leftarrow}x(\beta_{\leftarrow}x(x)) \otimes x) = S(x) \otimes S(x),$$

so $\Delta(S(x)) = S(x) \otimes S(x)$ and $\varepsilon(S(x)) = \varepsilon(x) = 1_{\mathbb{k}}$ and then $S(x) \in G(H)$. Thus, $(G(H), \cdot)$ is a group. Moreover, since \rightarrow is a coalgebra morphism, for all $x, y \in G(H)$ we have

$$\Delta(x \rightarrow y) = (x_1 \rightarrow y_1) \otimes (x_2 \rightarrow y_2) = (x \rightarrow y) \otimes (x \rightarrow y), \quad \varepsilon(x \rightarrow y) = \varepsilon(x)\varepsilon(y) = 1_{\mathbb{k}},$$

hence, we obtain a map $\rightarrow: G(H) \times G(H) \rightarrow G(H)$. Since (3.1) and (3.2) are satisfied for all $x, y, z \in H$, we have

$$x \rightarrow (y \cdot z) = (x \rightarrow y) \cdot (x \rightarrow z), \quad x \rightarrow (y \rightarrow z) = (x \cdot (x \rightarrow y)) \rightarrow z$$

for all $x, y, z \in G(H)$. In particular, $\alpha_{\leftarrow}x: G(H) \rightarrow G(H)$ is an automorphism of the group $G(H)$ for all $x \in G(H)$ and then $(G(H), \cdot, \rightarrow)$ is a post-group. ■

Since a Yetter–Drinfeld post-Hopf algebra (H, \rightarrow) is an object in $\text{Mon}({}_{H\leftarrow}^H\mathcal{YD})$, we call it a *Yetter–Drinfeld pre-Hopf algebra* if H is braided commutative, i.e., $m \circ \sigma_{H,H}^{\mathcal{YD}} = m$, which on elements $a, b \in H$ reads

$$\alpha_{\leftarrow}a_1(\beta_{\leftarrow}a_3(b)) \cdot a_2 = a \cdot b.$$

If H is cocommutative, then $\sigma_{H,H}^{\mathcal{YD}} = \tau_{H,H}$ and the braided commutativity becomes the classical commutativity.

Remark 3.20. Since a Yetter–Drinfeld post-Hopf algebra (H, \rightarrow) is an object in $\text{Hopf}({}_{H\leftarrow}^H\mathcal{YD})$, one can consider the braided commutator $[a, b]_{\mathcal{YD}} := a \cdot b - m \circ \sigma_{H,H}^{\mathcal{YD}}(a \otimes b)$ studying primitive elements. But if $a \in P(H)$, recalling that $\alpha_{\leftarrow}1 = \beta_{\leftarrow}1 = \text{Id}_H$, we obtain

$$\begin{aligned} \sigma_{H,H}^{\mathcal{YD}}(a \otimes b) &= \alpha_{\leftarrow}1(\beta_{\leftarrow}a(b)) \otimes 1 + \alpha_{\leftarrow}1(\beta_{\leftarrow}1(b)) \otimes a + \alpha_{\leftarrow}a(\beta_{\leftarrow}1(b)) \otimes 1 \\ &\stackrel{(3.14)}{=} \alpha_{\leftarrow}S_{\leftarrow}a(b) \otimes 1 + b \otimes a + \alpha_{\leftarrow}a(b) \otimes 1 \\ &= -\alpha_{\leftarrow}a(b) \otimes 1 + b \otimes a + \alpha_{\leftarrow}a(b) \otimes 1 \\ &= b \otimes a, \end{aligned}$$

hence, the braided commutator coincides with the classical one.

Finally, we obtain that any Yetter–Drinfeld post-Hopf algebra provides a Yetter–Drinfeld brace.

Corollary 3.21. *Suppose that (H, \rightarrow) is a Yetter–Drinfeld post-Hopf algebra. Then, we have that $(H, \cdot, \bullet_{\leftarrow}, 1, \Delta, \varepsilon, S, S_{\leftarrow})$ is a Yetter–Drinfeld brace. This yields a functor $F: \mathcal{YDPH}(\text{Vec}_{\mathbb{k}}) \rightarrow \mathcal{YDBr}(\text{Vec}_{\mathbb{k}})$.*

Proof. We know that $(H, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S_{\rightarrow})$ is a Hopf algebra by Proposition 3.10 and $(H, \cdot, 1, \Delta, \varepsilon, S)$ is in $\text{Hopf}(\mathcal{H}_{H_{\rightarrow}}^{\bullet} \mathcal{YD})$ by Theorem 3.17, where the coaction is given by Ad_L . Moreover, by (3.9), we know that the action \rightarrow is given by $a \rightarrow b = S(a_1) \cdot (a_2 \bullet_{\rightarrow} b)$ and, if we define \leftarrow as in (3.7), the pair $(\rightarrow, \leftarrow)$ satisfies (2.9). In order to conclude that $(H, \cdot, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S, S_{\rightarrow})$ is a Yetter–Drinfeld brace, it remains to prove that \cdot and \bullet_{\rightarrow} satisfy the compatibility of Hopf braces

$$\begin{aligned} x \bullet_{\rightarrow} (y \cdot z) &= x_1 \cdot (x_2 \rightarrow (y \cdot z)) \stackrel{(3.1)}{=} x_1 \cdot ((x_2 \rightarrow y) \cdot (x_3 \rightarrow z)) \\ &= x_1 \cdot (x_2 \rightarrow y) \cdot S(x_3) \cdot x_4 \cdot (x_5 \rightarrow z) \\ &= (x_1 \bullet_{\rightarrow} y) \cdot S(x_2) \cdot (x_3 \bullet_{\rightarrow} z). \end{aligned}$$

Let $f : (H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow) \rightarrow (H', \cdot', 1', \Delta', \varepsilon', S', \rightarrow')$ be a morphism of Yetter–Drinfeld post-Hopf algebras. We already know that f is a morphism of algebras and coalgebras, so we just have to show that it is multiplicative with respect to \bullet_{\rightarrow} to obtain that it is a morphism of Yetter–Drinfeld braces $f : (H, \cdot, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S, S_{\rightarrow}) \rightarrow (H', \cdot', \bullet'_{\rightarrow'}, 1', \Delta', \varepsilon', S', S'_{\rightarrow'})$. We compute

$$\begin{aligned} f(a \bullet_{\rightarrow} b) &= f(a_1 \cdot (a_2 \rightarrow b)) = f(a_1) \cdot f(a_2 \rightarrow b) \stackrel{(3.8)}{=} f(a_1) \cdot (f(a_2) \rightarrow' f(b)) \\ &= f(a)_1 \cdot (f(a)_2 \rightarrow' f(b)) = f(a) \bullet_{\rightarrow'} f(b). \end{aligned}$$

Clearly, the assignment F respects identities and compositions. ■

We can also prove that any Yetter–Drinfeld brace provides a Yetter–Drinfeld post-Hopf algebra as follows.

Proposition 3.22. *Let us assume that $(H, \cdot, \bullet, 1, \Delta, \varepsilon, S, T)$ is a Yetter–Drinfeld brace. Then, $(H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow)$ is a Yetter–Drinfeld post-Hopf algebra with \rightarrow defined by*

$$a \rightarrow b := S(a_1) \cdot (a_2 \bullet b)$$

and $\beta_{\rightarrow} a : b \mapsto (T(a) \rightarrow b)$ for all $a, b \in H$. This yields a functor $G : \mathcal{YD}\text{Br}(\text{Vec}_{\mathbb{k}}) \rightarrow \mathcal{YD}\text{PH}(\text{Vec}_{\mathbb{k}})$.

Proof. Since $(H, \cdot, \bullet, 1, \Delta, \varepsilon, S, T)$ is a Yetter–Drinfeld brace, by definition $(H, \cdot, 1, \Delta, \varepsilon, S)$ is in $\text{Hopf}(\mathcal{H}_H^{\bullet} \mathcal{YD})$, where $H^{\bullet} := (H, \bullet, 1, \Delta, \varepsilon, T)$ is a Hopf algebra, the H^{\bullet} -action and the H^{\bullet} -coaction on H are given by $a \rightarrow b := S(a_1) \cdot (a_2 \bullet b)$ and $\text{Ad}_L : a \mapsto a_1 \bullet T(a_3) \otimes a_2$, respectively. In particular, $(H, \cdot, 1)$ is an algebra, (H, Δ, ε) is a coalgebra, S satisfies $a_1 \cdot S(a_2) = \varepsilon(a)1 = S(a_1) \cdot a_2$ and \rightarrow is a morphism of coalgebras which satisfies (3.1). From $a_1 \cdot (a_2 \rightarrow b) = a_1 \cdot S(a_2) \cdot (a_3 \bullet b) = a \bullet b$, also (3.2) is automatically true as \rightarrow is an action. Define $(\beta_{\rightarrow} a)(b) := T(a) \rightarrow b$, for all $a, b \in H$. Hence, we have

$$\beta_{\rightarrow} a_1(\alpha_{\rightarrow} a_2(b)) = T(a_1) \rightarrow (a_2 \rightarrow b) = (T(a_1) \bullet a_2) \rightarrow b = \varepsilon(a)1 \rightarrow b = \varepsilon(a)b,$$

and, similarly,

$$\alpha_{\rightarrow} a_1(\beta_{\rightarrow} a_2(b)) = \varepsilon(a)b,$$

so α_{\rightarrow} is convolution invertible in $\text{Hom}(H, \text{End}(H))$ with inverse given by β_{\rightarrow} . Moreover, we already know that ε and Δ are morphisms of algebras in $\frac{H}{H} \bullet \mathcal{YD}$, i.e., $\varepsilon(a \cdot b) = \varepsilon(a)\varepsilon(b)$, $\varepsilon(1_H) = 1_{\mathbb{k}}$, $\Delta(1_H) = 1_H \otimes 1_H$ and the following compatibility condition holds true:

$$\Delta(a \cdot b) = a_1 \cdot (a_2 \bullet T(a_4) \rightarrow b_1) \otimes (a_3 \cdot b_2). \tag{3.25}$$

Hence, we have

$$\Delta(a \cdot b) \stackrel{(3.25)}{=} a_1 \cdot (a_2 \rightarrow (T(a_4) \rightarrow b_1)) \otimes (a_3 \cdot b_2) = (a_1 \cdot (\alpha_{\rightarrow} a_2 (\beta_{\rightarrow} a_4 (b_1)))) \otimes (a_3 \cdot b_2),$$

i.e., (3.4) is satisfied. Finally, $a \bullet_{\rightarrow} b := a_1 \cdot (a_2 \rightarrow b) = a \bullet b$ and since $a_1 \rightarrow T(a_2) = S(a_1) \cdot (a_2 \bullet T(a_3)) = S(a)$, we obtain

$$S_{\rightarrow}(a) \stackrel{(3.6)}{=} \beta_{\rightarrow} a_1 (S(a_2)) = T(a_1) \rightarrow (a_2 \rightarrow T(a_3)) = (T(a_1) \bullet a_2) \rightarrow T(a_3) = T(a).$$

Hence, S_{\rightarrow} is anti-comultiplicative and $a \leftarrow b := S_{\rightarrow}(a_1 \rightarrow b_1) \bullet_{\rightarrow} a_2 \bullet_{\rightarrow} b_2 = T(a_1 \rightarrow b_1) \bullet a_2 \bullet b_2$. Thus, by definition of Yetter–Drinfeld brace, the pair $(\rightarrow, \leftarrow)$ satisfies (2.9) and (H, \rightarrow) is a Yetter–Drinfeld post-Hopf algebra.

Let $f : (H, \cdot, \bullet, 1, \Delta, \varepsilon, S, T) \rightarrow (H', \cdot', \bullet', 1', \Delta', \varepsilon', S', T')$ be a morphism of Yetter–Drinfeld braces, so $f : (H, \bullet, 1, \Delta, \varepsilon, T) \rightarrow (H', \bullet', 1', \Delta', \varepsilon', T')$ is a morphism of Hopf algebras and satisfies $f(a \cdot b) = f(a) \cdot' f(b)$ and $fS = S'f$. We prove that $f(a \rightarrow b) = f(a) \rightarrow' f(b)$, so that $f : (H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow) \rightarrow (H', \cdot', 1', \Delta', \varepsilon', S', \rightarrow')$ is a morphism of Yetter–Drinfeld post-Hopf algebras:

$$\begin{aligned} f(a \rightarrow b) &= f(S(a_1) \cdot (a_2 \bullet b)) = f(S(a_1)) \cdot' f(a_2 \bullet b) = S'(f(a_1)) \cdot' (f(a_2) \bullet' f(b)) \\ &= S'(f(a)_1) \cdot' (f(a)_2 \bullet' f(b)) = f(a) \rightarrow' f(b). \end{aligned}$$

Clearly, the assignment G respects identities and compositions. ■

Remark 3.23. The previous proof shows that the subadjacent Hopf algebra of the Yetter–Drinfeld post-Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow)$, where the latter is obtained from the Yetter–Drinfeld brace $(H, \cdot, \bullet, 1, \Delta, \varepsilon, S, T)$, is given by the Hopf algebra $(H, \bullet, 1, \Delta, \varepsilon, T)$. Then, $FG = \text{Id}$. Moreover, given a Yetter–Drinfeld post-Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S \rightarrow)$ and the Yetter–Drinfeld brace $F((H, \rightarrow)) = (H, \cdot, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S, S_{\rightarrow})$, then $GF((H, \rightarrow))$ is the Yetter–Drinfeld post-Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow')$ where

$$a \rightarrow' b := S(a_1) \cdot (a_2 \bullet_{\rightarrow} b) = S(a_1) \cdot a_2 \cdot (a_3 \rightarrow b) = a \rightarrow b.$$

Hence, $GF = \text{Id}$ and we obtain the following result.

Corollary 3.24. *The categories $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$ and $\mathcal{YDBr}(\text{Vec}_{\mathbb{k}})$ are isomorphic.*

By [12, Theorem 3.22], we know that the category $\mathcal{YDBr}(\text{Vec}_{\mathbb{k}})$ is isomorphic to the category $\text{MP}(\text{Vec}_{\mathbb{k}})$ of matched pairs of actions. Thus, we also obtain the following result.

Corollary 3.25. *The categories $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$ and $\text{MP}(\text{Vec}_{\mathbb{k}})$ are isomorphic. Explicitly, from a Yetter–Drinfeld post-Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ one obtains a matched pair of actions $(H_{\rightharpoonup}, \rightharpoonup, \leftarrow)$ on the subadjacent Hopf algebra H_{\rightharpoonup} , where \leftarrow is defined as in (3.7). Conversely, given a matched pairs of actions $(H, \rightharpoonup, \leftarrow)$ on a Hopf algebra $(H, \bullet, 1, \Delta, \varepsilon, T)$ one obtains a Yetter–Drinfeld post-Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ where $a \cdot b := a_1 \bullet (T(a_2) \rightharpoonup b)$ and $S(a) := a_1 \rightharpoonup T(a_2)$.*

The previous result generalises [22, Corollary 4.5].

To end this subsection, we make a connection with braiding operators. It is known that matched pairs of actions on cocommutative Hopf algebras, braiding operators on cocommutative Hopf algebras and cocommutative Hopf braces are equivalent [16]. Moreover, in [15, Theorem 6.34] the category of Yetter–Drinfeld braces is proven to be isomorphic to the category of braiding operators on Hopf algebras [15, Definition 5.1]. Hence, from Corollary 3.24, we obtain the following corollary.

Corollary 3.26. *The category $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$ is isomorphic to the category of braiding operators on Hopf algebras.*

Let us recall the definition of *braiding operator on a Hopf algebra H* as stated in [20, Definition 3.1]. It consists of a coalgebra morphism $r : H \otimes H \rightarrow H \otimes H$ such that

- (i) $mr = m$,
- (ii) $r(m \otimes \text{Id}) = (\text{Id} \otimes m)(r \otimes \text{Id})(\text{Id} \otimes r)$,
- (iii) $r(\text{Id} \otimes m) = (m \otimes \text{Id})(\text{Id} \otimes r)(r \otimes \text{Id})$,
- (iv) $r(u \otimes \text{Id}) = \text{Id} \otimes u$,
- (v) $r(\text{Id} \otimes u) = u \otimes \text{Id}$,

where m and u denote the multiplication and the unit of H , respectively.

In [15, Theorem 5.27], it is proved that braiding operators on Hopf algebras provide solutions of the braid equation. Thus, as it is shown in [20, Theorem 3.2], matched pairs of actions on Hopf algebras $(H, \rightharpoonup, \leftarrow)$ provide solutions to the braid equation $r(x \otimes y) := (x_1 \rightharpoonup y_1) \otimes (x_2 \leftarrow y_2)$. This result extends [2, Corollary 2.4], obtained under the cocommutativity assumption. Therefore, from Corollary 3.25, we obtain the following proposition.

Proposition 3.27. *Let $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ be a Yetter–Drinfeld post-Hopf algebra. Then,*

$$r : H \otimes H \rightarrow H \otimes H, \quad x \otimes y \mapsto (x_1 \rightharpoonup y_1) \otimes (x_2 \leftarrow y_2) \tag{3.26}$$

is a solution of the braid equation, where \leftarrow is defined as in (3.7). More explicitly, we have

$$r(x \otimes y) = (x_1 \rightharpoonup y_1) \otimes (\beta_{\rightharpoonup}(x_4 \rightharpoonup y_4)(x_5 \rightharpoonup S(y_5)) \cdot \beta_{\rightharpoonup}(x_3 \rightharpoonup y_3)(x_6) \cdot \beta_{\rightharpoonup}(x_2 \rightharpoonup y_2)(x_7 \rightharpoonup y_6)),$$

where β_{\rightharpoonup} is the convolution inverse of α_{\rightharpoonup} .

Proof. We compute

$$\begin{aligned}
 & (x_1 \rightharpoonup y_1) \otimes (x_2 \leftarrow y_2) \\
 & \stackrel{(3.7)}{=} (x_1 \rightharpoonup y_1) \otimes (S_{\leftarrow}(x_2 \rightharpoonup y_2) \bullet_{\leftarrow} x_3 \bullet_{\leftarrow} y_3) \\
 & \stackrel{(3.5)}{=} (x_1 \rightharpoonup y_1) \otimes (S_{\leftarrow}(x_3 \rightharpoonup y_3) \cdot (S_{\leftarrow}(x_2 \rightharpoonup y_2) \rightharpoonup (x_4 \cdot (x_5 \rightharpoonup y_4)))) \\
 & \stackrel{(3.1)}{=} (x_1 \rightharpoonup y_1) \otimes (S_{\leftarrow}(x_4 \rightharpoonup y_4) \cdot (S_{\leftarrow}(x_3 \rightharpoonup y_3) \rightharpoonup x_5) \\
 & \qquad \qquad \qquad \cdot (S_{\leftarrow}(x_2 \rightharpoonup y_2) \rightharpoonup (x_6 \rightharpoonup y_5))) \\
 & = (x_1 \rightharpoonup y_1) \otimes (S_{\leftarrow}(x_4 \rightharpoonup y_4) \cdot \alpha_{\leftarrow} S_{\leftarrow}(x_3 \rightharpoonup y_3)(x_5) \\
 & \qquad \qquad \qquad \cdot \alpha_{\leftarrow} S_{\leftarrow}(x_2 \rightharpoonup y_2)(x_6 \rightharpoonup y_5)) \\
 & \stackrel{(3.14)}{=} (x_1 \rightharpoonup y_1) \otimes (S_{\leftarrow}(x_4 \rightharpoonup y_4) \cdot \beta_{\leftarrow}(x_3 \rightharpoonup y_3)(x_5) \cdot \beta_{\leftarrow}(x_2 \rightharpoonup y_2)(x_6 \rightharpoonup y_5)) \\
 & \stackrel{(3.6), (3.13)}{=} (x_1 \rightharpoonup y_1) \otimes (\beta_{\leftarrow}(x_4 \rightharpoonup y_4)(x_5 \rightharpoonup S(y_5)) \cdot \beta_{\leftarrow}(x_3 \rightharpoonup y_3)(x_6) \\
 & \qquad \qquad \qquad \cdot \beta_{\leftarrow}(x_2 \rightharpoonup y_2)(x_7 \rightharpoonup y_6)),
 \end{aligned}$$

where β_{\leftarrow} is the convolution inverse of α_{\leftarrow} . ■

3.2. Examples of Yetter–Drinfeld post-Hopf algebras

By Proposition 3.22, we can use Yetter–Drinfeld braces to provide examples of Yetter–Drinfeld post-Hopf algebras. In [12, Section 6] some examples of Yetter–Drinfeld braces are given, using matched pairs of actions coming from coquasitriangular structures \mathcal{R} (we refer the reader to [24] for coquasitriangular Hopf algebras). In this case, as it is said in [12, Remark 5.6], the structure $(H, \cdot, 1, \Delta, \varepsilon, S)$ results to be the transmutation of $(H, \bullet, 1, \Delta, \varepsilon, T, \mathcal{R})$, see [24, dual of Example 9.4.10] and [23]. Hence, translating [12, Theorem 5.5] into our setting, we get the following result.

Lemma 3.28. *Let $(H, \bullet, 1, \Delta, \varepsilon, T, \mathcal{R})$ be a coquasitriangular Hopf algebra. Define \cdot and S as*

$$a \cdot b := a_1 \bullet b_2 \mathcal{R}^{-1}(T(a_2) \otimes b_1 \bullet T(b_3)), \quad S(a) := \mathcal{R}(a_1 \otimes a_5) T(a_4) \mathcal{R}(a_2 \otimes T(a_3))$$

for all $a, b \in H$, where \mathcal{R}^{-1} denotes the convolution inverse of $\mathcal{R} : H \otimes H \rightarrow \mathbb{k}$. Then, we have $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ is a Yetter–Drinfeld post-Hopf algebra where \rightharpoonup is defined by

$$a \rightharpoonup b := b_2 \mathcal{R}^{-1}(a \otimes b_1 \bullet T(b_3))$$

and $\beta_{\leftarrow} a : b \mapsto b_2 \mathcal{R}^{-1}(T(a) \otimes b_1 \bullet T(b_3))$.

In the examples of [12, Section 6], the morphisms \rightharpoonup are explicitly computed, hence, we just have to collect the results writing them in terms of the \cdot operation. We will only explain the structures $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$, saying which standard Hopf algebras they are the transmutation of, without indicating the \mathcal{R} structures; the standard Hopf algebras correspond to the subadjacent Hopf algebras. We refer the reader to [12, Section 6] for more details about the following examples.

Example 3.29. Let \mathbb{k} be a field of $\text{char}(\mathbb{k}) \neq 2$ and the algebra $(H, \cdot, 1)$ generated by g, x modulo the relations

$$g \cdot g = 1, \quad x \cdot x = k1 - kg, \quad x \cdot g = g \cdot x,$$

where $k \in \mathbb{k}$. Define Δ, ε and S on g and x as

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes 1 + g \otimes x, \quad \varepsilon(g) = 1_{\mathbb{k}}, \quad \varepsilon(x) = 0, \quad S(g) = g, \quad S(x) = -x \cdot g.$$

We have that $(H, \cdot, 1, \Delta, \varepsilon, S)$ is the transmutation of the Sweedler’s Hopf algebra H_4 , see [12, 6.1]. Moreover, $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ is a Yetter–Drinfeld post-Hopf algebra, where \rightharpoonup is given as in the following left table and α_{\rightharpoonup} has convolution inverse β_{\rightharpoonup} explained in the right table:

\rightharpoonup	1	g	x	$x \cdot g$	β_{\rightharpoonup}	1	g	x	$x \cdot g$
1	1	g	x	$x \cdot g$	1	1	g	x	$x \cdot g$
g	1	g	$-x$	$-x \cdot g$	g	1	g	$-x$	$-x \cdot g$
x	0	0	$k(1 - g)$	$k(g - 1)$	x	0	0	$k(g - 1)$	$k(1 - g)$
$x \cdot g$	0	0	$k(g - 1)$	$k(1 - g)$	$x \cdot g$	0	0	$k(g - 1)$	$k(1 - g)$

Let us observe that post-Hopf algebra structures on the Sweedler’s Hopf algebra are classified in [22, Example 2.12]. The Sweedler’s Hopf algebra is not cocommutative and indeed the post-Hopf algebra structures generally differ from the previous Yetter–Drinfeld post-Hopf algebra structures, in which H is a Hopf monoid in the category of Yetter–Drinfeld modules and not a standard Hopf algebra.

Example 3.30. Let \mathbb{k} be a field of $\text{char}(\mathbb{k}) \neq 2$. For a fixed $1 \leq n \in \mathbb{N}$, consider the algebra $(H, \cdot, 1)$ generated by g, x_1, \dots, x_n subject to the relations

$$g \cdot g = 1, \quad x_i \cdot x_j + x_j \cdot x_i = 2A_{ij}(1 - g), \quad x_i \cdot g = g \cdot x_i,$$

where $A = (A_{ij})_{ij}$ is an $n \times n$ symmetric matrix with entries in \mathbb{k} . Define Δ, ε and S on g, x_i as

$$\Delta(g) = g \otimes g, \quad \Delta(x_i) = x_i \otimes 1 + g \otimes x_i, \quad \varepsilon(g) = 1_{\mathbb{k}}, \quad \varepsilon(x_i) = 0, \quad S(g) = g, \quad S(x_i) = -x_i \cdot g.$$

In case $n = 1$, we recover the previous example. We have that $(H, \cdot, 1, \Delta, \varepsilon, S)$ is the transmutation of the Hopf algebra $E(n)$, see [12, 6.2]. Moreover, $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ is a Yetter–Drinfeld post-Hopf algebra, where \rightharpoonup and β_{\rightharpoonup} are defined by the following tables:

\rightharpoonup	1	g	x_j	$x_j \cdot g$	β_{\rightharpoonup}	1	g	x_j	$x_j \cdot g$
1	1	g	x_j	$x_j \cdot g$	1	1	g	x_j	$x_j \cdot g$
g	1	g	$-x_j$	$-x_j \cdot g$	g	1	g	$-x_j$	$-x_j \cdot g$
x_i	0	0	$A_{ij}(1 - g)$	$A_{ij}(g - 1)$	x_i	0	0	$A_{ij}(g - 1)$	$A_{ij}(1 - g)$
$x_i \cdot g$	0	0	$A_{ij}(g - 1)$	$A_{ij}(1 - g)$	$x_i \cdot g$	0	0	$A_{ij}(g - 1)$	$A_{ij}(1 - g)$

Example 3.31. Let $0 \neq q \in \mathbb{C}$ and consider the \mathbb{C} -algebra $(H, \cdot, 1)$ generated by a, b, c, d modulo the relations

$$\begin{aligned} a \cdot b &= q^2 b \cdot a, & b \cdot c - c \cdot b &= (q^2 - 1)(a \cdot a - a \cdot d), \\ a \cdot c &= q^{-2} c \cdot a, & b \cdot d - d \cdot b &= (q^2 - 1)a \cdot b, \\ a \cdot d &= d \cdot a, & c \cdot d - d \cdot c &= (1 - q^2)c \cdot a, \end{aligned}$$

and $a \cdot d - q^{-2}c \cdot b = 1$. Define Δ, ε and S on a, b, c, d as

$$\Delta \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \otimes \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad \varepsilon \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and

$$S \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} q^{-2}d + (1 - q^{-2})a & -q^{-2}b \\ -q^{-2}c & a \end{pmatrix}.$$

We have that $(H, \cdot, 1, \Delta, \varepsilon, S)$ is the transmutation of the Hopf algebra $SL_q(2)$, see [12, 6.3]. Moreover, $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ is a Yetter–Drinfeld post-Hopf algebra where \rightharpoonup is given as in the following table:

\rightharpoonup	a	b	c	d
a	a	$q^{-1}b$	qc	d
b	$(1 - q^{-2})b$	0	$q(1 - q^{-2})(d - a)$	$(1 - q^2)b$
c	0	0	0	0
d	a	qb	$q^{-1}c$	d

Moreover, α_{\rightharpoonup} has convolution inverse β_{\rightharpoonup} given as in the following table:

β_{\rightharpoonup}	a	b	c	d
a	a	qb	$q^{-1}c$	d
b	$q(q^{-2} - 1)b$	0	$(1 - q^2)(d - a)$	$-q(1 - q^2)b$
c	0	0	0	0
d	a	$q^{-1}b$	qc	d

Example 3.32. Let \mathbb{k} be an algebraically closed field of $\text{char}(\mathbb{k}) \neq 2$. Consider the algebra $(H, \cdot, 1)$ generated by a, b, c, d modulo the relations

$$\begin{aligned} a \cdot a &= d \cdot d, & c \cdot c &= \alpha\beta^{-1}b \cdot b, & c \cdot b &= b \cdot c, & a \cdot d &= d \cdot a, \\ a \cdot b &= b \cdot a = a \cdot c = c \cdot a = b \cdot d = d \cdot b = c \cdot d = d \cdot c = 0, \end{aligned}$$

where $\alpha, \beta \in \mathbb{k}$. Define Δ, ε and S on a, b, c, d as

$$\Delta \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \otimes \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \quad \varepsilon \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and

$$S \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} \alpha^2\beta^2(a \cdot d^2) & \alpha^4(b^2 \cdot c) \\ \alpha^4(b \cdot c^2) & \alpha^2\beta^2(d \cdot a^2) \end{pmatrix}.$$

We have that $(H, \cdot, 1, \Delta, \varepsilon, S)$ is the transmutation of the Suzuki Hopf algebra $A_{1,2}^{\nu,\lambda}$, see [12, 6.4]. Moreover, $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$ is a Yetter–Drinfeld post-Hopf algebra where \rightharpoonup is given as in the following table, which coincides also with the table of β_{\rightharpoonup} .

\rightharpoonup	a	b	c	d
a	d	$\alpha^{-1}\beta c$	$\alpha\beta^{-1}b$	a
b	0	0	0	0
c	0	0	0	0
d	d	$\alpha\beta^{-1}c$	$\alpha^{-1}\beta b$	a

Remark 3.33. Notice that, starting from a Hopf algebra $(H, \bullet, 1, \Delta, \varepsilon, T)$, one can obtain a Yetter–Drinfeld brace through the matched pair of actions $(\rightharpoonup, \leftarrow)$ given by the left adjoint action $a \rightharpoonup b := a_1 \bullet b \bullet T(a_2)$ and the right trivial action $a \leftarrow b := a\varepsilon(b)$ as in [12, Lemma 3.19]. Hence, one obtains a Yetter–Drinfeld post-Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup)$, where \cdot and S are defined by

$$a \cdot b := a_1 \bullet T(a_3) \bullet b \bullet T(T(a_2)), \quad S(a) := a_1 \bullet T(a_3) \bullet T(a_2).$$

Recall that $\beta_{\rightharpoonup} a : b \mapsto (T(a) \rightharpoonup b)$. This can be a way to produce other examples of Yetter–Drinfeld post-Hopf algebras, which are not post-Hopf algebras in general.

4. Yetter–Drinfeld relative Rota–Baxter operators

The notion of *Rota–Baxter operator* on a cocommutative Hopf algebra H was introduced in [13] as a morphism of coalgebras $R : H \rightarrow H$ satisfying

$$R(x)R(y) = R(x_1 \text{ad}_L(R(x_2) \otimes y)) = R(x_1 R(x_2) y S(R(x_3))) \quad \text{for all } x, y \in H,$$

where ad_L denotes the left adjoint action $H \otimes H \rightarrow H, x \otimes y \mapsto x_1 y S(x_2)$, then generalised for arbitrary actions with the notion of *relative Rota–Baxter operator* on a cocommutative Hopf algebra [22, Definition 3.1]. Moreover, in [22, Theorem 3.4] an adjunction between the categories of relative Rota–Baxter operators on cocommutative Hopf algebras and cocommutative post-Hopf algebras was provided.

Remark 4.1. In [22, Definition 3.1], H and K are cocommutative Hopf algebras and K is also an object in $\text{Bimon}(H\mathfrak{M})$ (where the braiding considered for $H\mathfrak{M}$ is given by the canonical flip, which corresponds to the quasitriangular structure $1_H \otimes 1_H$ on H). But then, mimicking the proof of [12, Proposition 3.9], one obtains that S_K is in $H\mathfrak{M}$, hence, K is actually an object in $\text{Hopf}(H\mathfrak{M})$.

We introduce a generalisation of the notion of relative Rota–Baxter operator, which we call *Yetter–Drinfeld relative Rota–Baxter operator*. A subcategory of the category of bijective Yetter–Drinfeld relative Rota–Baxter operators will turn out to be equivalent to the category of Yetter–Drinfeld post-Hopf algebras.

Definition 4.2. Let H be a Hopf algebra and K be an object in $\text{Bimon}({}_H^H \mathcal{YD})$, and let the action of H on K be denoted by \rightharpoonup . A coalgebra morphism $R : K \rightarrow H$ is called a *Yetter–Drinfeld relative Rota–Baxter operator* on H with respect to (K, \rightharpoonup) if the following equalities hold, for all $a, b \in K$:

$$R(a) \cdot R(b) = R(a_1 \cdot (R(a_2) \rightharpoonup b)), \tag{4.1}$$

$$\begin{aligned} S_H R(R(a_1) \rightharpoonup b_1) \cdot R(a_2) \cdot R(b_2) \otimes R(R(a_3) \rightharpoonup b_3) \\ = S_H R(R(a_2) \rightharpoonup b_2) \cdot R(a_3) \cdot R(b_3) \otimes R(R(a_1) \rightharpoonup b_1). \end{aligned} \tag{4.2}$$

A morphism of Yetter–Drinfeld relative Rota–Baxter operators from $R : K \rightarrow H$ to $R' : K' \rightarrow H'$ is a pair of morphisms of algebras and coalgebras $(f : H \rightarrow H', g : K \rightarrow K')$ such that

$$fR = R'g, \quad g \rightharpoonup = \rightharpoonup' (f \otimes g). \tag{4.3}$$

The category of Yetter–Drinfeld relative Rota–Baxter operators and morphisms of Yetter–Drinfeld relative Rota–Baxter operators will be denoted by $\mathcal{YDrRB}(\text{Vec}_{\mathbb{k}})$.

Remark 4.3. Since $R(1_K)$ is a group-like element in $G(H)$, it has inverse (given by $S_H R(1_K)$). Moreover, from (4.1), we have

$$R(1_K) \cdot R(1_K) = R(1_K \cdot (R(1_K) \rightharpoonup 1_K)) = R(\varepsilon(R(1_K))1_K) = R(1_K),$$

hence, $R(1_K) = 1_H$.

Remark 4.4. Notice that if K is cocommutative, then (4.2) is automatically satisfied. Moreover, if $R : K \rightarrow H$ is bijective, the cocommutativity of K induces the cocommutativity of H (and vice versa). We point out that a subcategory of bijective Yetter–Drinfeld relative Rota–Baxter operators will turn out to be equivalent to the category of Yetter–Drinfeld post-Hopf algebras and (4.2) will correspond to (2.9).

Even when R is bijective, the morphism $R^{-1}S_H R$ is not an antipode for K in general, since R is not a morphism of algebras, but we can equip K with an antipode in the following way.

Lemma 4.5. *Let $R : K \rightarrow H$ be a bijective Yetter–Drinfeld relative Rota–Baxter operator. Then, the morphism $S_K : K \rightarrow K, a \mapsto (R(a_1) \rightharpoonup R^{-1}S_H R(a_2))$ satisfies*

$$S_K(a_1) \cdot a_2 = \varepsilon(a)1_K = a_1 \cdot S_K(a_2).$$

As a consequence, K is an object in $\text{Hopf}({}_H^H \mathcal{YD})$.

Proof. First, we compute

$$\begin{aligned} R(a_1 \cdot S_K(a_2)) &= R(a_1 \cdot (R(a_2) \rightharpoonup R^{-1}S_H R(a_3))) \stackrel{(4.1)}{=} R(a_1) \cdot RR^{-1}S_H R(a_2) \\ &= R(a)_1 \cdot S_H(R(a)_2) = \varepsilon(R(a))1_H = \varepsilon(a)1_H, \end{aligned}$$

so that $a_1 \cdot S_K(a_2) = \varepsilon(a)1_K$. Moreover, using that \rightarrow is multiplicative, we also have

$$\begin{aligned} R(S_H R(a_1) \rightarrow (S_K(a_2) \cdot a_3)) &= R((S_H R(a_1)_1 \rightarrow S_K(a_2)) \cdot (S_H R(a_1)_2 \rightarrow a_3)) \\ &= R((S_H R(a_2) \rightarrow S_K(a_3)) \cdot (S_H R(a_1) \rightarrow a_4)) \\ &= R((S_H R(a_2) \rightarrow (R(a_3) \rightarrow R^{-1} S_H R(a_4))) \cdot (S_H R(a_1) \rightarrow a_5)) \\ &= R(R^{-1} S_H R(a_2) \cdot (S_H R(a_1) \rightarrow a_3)) \\ &= R(R^{-1} S_H R(a_1)_1 \cdot (R(R^{-1} S_H R(a_1)_2) \rightarrow a_2)) \\ &= RR^{-1} S_H R(a_1) \cdot R(a_2) = S_H(R(a_1)) \cdot R(a_2) \\ &= \varepsilon(a)1_H, \end{aligned}$$

so that $S_H R(a_1) \rightarrow (S_K(a_2) \cdot a_3) = \varepsilon(a)1_K$. Then, we obtain

$$\begin{aligned} S_K(a_1) \cdot a_2 = R(a_1) \rightarrow (S_H R(a_2) \rightarrow (S_K(a_3) \cdot a_4)) &= R(a_1) \rightarrow \varepsilon(a_2)1_K \\ &= \varepsilon(R(a))1_K = \varepsilon(a)1_K \end{aligned}$$

and so, K is an object in $\text{Hopf}(\mathcal{H}_H \mathcal{YD})$ by [12, Proposition 3.9]. ■

We denote by $\text{BYDrRB}(\text{Vec}_{\mathbb{k}})$ the category of bijective Yetter–Drinfeld relative Rota–Baxter operators.

Remark 4.6. We have shown that every bijective Yetter–Drinfeld relative Rota–Baxter operator $R : K \rightarrow H$ is such that K is in $\text{Hopf}(\mathcal{H}_H \mathcal{YD})$. Clearly, one can deduce the same for Yetter–Drinfeld 1-cocycles in the sense of [12, Definition 4.1], so that (20) in [12, Definition 4.1] becomes redundant. We have that R is a bijective Yetter–Drinfeld relative Rota–Baxter operator if and only if R^{-1} is a Yetter–Drinfeld 1-cocycle in the sense of [12, Definition 4.1]; this result generalises [31, Proposition 3.18].

Moreover, denoting by \mathcal{C} the full subcategory of $\text{YDrRB}(\text{Vec}_{\mathbb{k}})$ whose objects are Yetter–Drinfeld relative Rota–Baxter operators $R : K \rightarrow H$ such that K is in $\text{Hopf}(\mathcal{H}_H \mathcal{YD})$, the category $\text{BYDrRB}(\text{Vec}_{\mathbb{k}})$ is a subcategory of \mathcal{C} .

A Yetter–Drinfeld relative Rota–Baxter operator naturally induces relative Rota–Baxter operators on the group $G(H)$ and on the Lie algebra $P(H)$.

Proposition 4.7. *Let $R : K \rightarrow H$ be a Yetter–Drinfeld relative Rota–Baxter operator on H with respect to (K, \rightarrow) . Then, the following statements hold:*

- (1) $R|_{G(K)} : G(K) \rightarrow G(H)$ is a relative Rota–Baxter operator on the group $G(H)$ with respect to $(G(K), \rightarrow)$,
- (2) $R|_{P(K)} : P(K) \rightarrow P(H)$ is a relative Rota–Baxter operator on the Lie algebra $P(H)$ with respect to $(P(K), \rightarrow)$.

The previous result generalises [22, Theorem 3.2] and it has the same proof, which only uses (4.1) and the fact that R is a morphism of coalgebras. Hence, Yetter–Drinfeld relative Rota–Baxter operators are natural candidates in a non-cocommutative setting.

4.1. Connection with Yetter–Drinfeld post-Hopf algebras

A Yetter–Drinfeld post-Hopf algebra produces a Yetter–Drinfeld relative Rota–Baxter operator, as it is shown in the following result that generalises [22, Proposition 3.3] removing the hypothesis of cocommutativity.

Proposition 4.8. *Let (H, \rightharpoonup) be a Yetter–Drinfeld post-Hopf algebra and H_{\rightharpoonup} the subadjacent Hopf algebra defined in Proposition 3.10. Then, the identity map $\text{Id}_H : H \rightarrow H_{\rightharpoonup}$ is a (bijective) Yetter–Drinfeld relative Rota–Baxter operator on H_{\rightharpoonup} with respect to (H, \rightharpoonup) . Moreover, if $g : (H, \rightharpoonup) \rightarrow (H', \rightharpoonup')$ is a morphism of Yetter–Drinfeld post-Hopf algebras, then $(g : H_{\rightharpoonup} \rightarrow H'_{\rightharpoonup'}, g : H \rightarrow H')$ is a morphism of Yetter–Drinfeld relative Rota–Baxter operators from $\text{Id}_H : H \rightarrow H_{\rightharpoonup}$ to $\text{Id}_{H'} : H' \rightarrow H'_{\rightharpoonup'}$. This yields a functor $L : \mathcal{YDPH}(\text{Vec}_{\mathbb{k}}) \rightarrow \mathcal{YDrRB}(\text{Vec}_{\mathbb{k}})$.*

Proof. For any $a, b \in H$, we have

$$\text{Id}_H(a) \bullet_{\rightharpoonup} \text{Id}_H(b) = a \bullet_{\rightharpoonup} b \stackrel{(3.5)}{=} a_1 \cdot (a_2 \rightharpoonup b) = \text{Id}_H(a_1 \cdot (\text{Id}_H(a_2) \rightharpoonup b)),$$

so (4.1) holds true. Moreover, also (4.2) is satisfied:

$$\begin{aligned} S_{\rightharpoonup} \text{Id}_H(\text{Id}_H(a_1) \rightharpoonup b_1) \bullet_{\rightharpoonup} \text{Id}_H(a_2) \bullet_{\rightharpoonup} \text{Id}_H(b_2) \otimes \text{Id}_H(\text{Id}_H(a_3) \rightharpoonup b_3) \\ &= S_{\rightharpoonup}(a_1 \rightharpoonup b_1) \bullet_{\rightharpoonup} a_2 \bullet_{\rightharpoonup} b_2 \otimes (a_3 \rightharpoonup b_3) \\ &\stackrel{(3.7)}{=} (a_1 \leftarrow b_1) \otimes (a_2 \rightharpoonup b_2) \\ &\stackrel{(2.9)}{=} (a_2 \leftarrow b_2) \otimes (a_1 \rightharpoonup b_1) \\ &\stackrel{(3.7)}{=} S_{\rightharpoonup}(a_2 \rightharpoonup b_2) \bullet_{\rightharpoonup} a_3 \bullet_{\rightharpoonup} b_3 \otimes (a_1 \rightharpoonup b_1) \\ &= S_{\rightharpoonup} \text{Id}_H(\text{Id}_H(a_2) \rightharpoonup b_2) \bullet_{\rightharpoonup} \text{Id}_H(a_3) \bullet_{\rightharpoonup} \text{Id}_H(b_3) \otimes \text{Id}_H(\text{Id}_H(a_1) \rightharpoonup b_1). \end{aligned}$$

Thus, $\text{Id}_H : H \rightarrow H_{\rightharpoonup}$ is a (bijective) Yetter–Drinfeld relative Rota–Baxter operator on H_{\rightharpoonup} with respect to (H, \rightharpoonup) .

Let $g : (H, \cdot, 1, \Delta, \varepsilon, S, \rightharpoonup) \rightarrow (H', \cdot', 1', \Delta', \varepsilon', S', \rightharpoonup')$ be a Yetter–Drinfeld post-Hopf algebra morphism, i.e., $g : H \rightarrow H'$ is a morphism of algebras and coalgebras which satisfies $g \rightharpoonup = \rightharpoonup' (g \otimes g)$. Clearly, the pair $(g : H_{\rightharpoonup} \rightarrow H'_{\rightharpoonup'}, g : H \rightarrow H')$ satisfies (4.3). Moreover, we have

$$\begin{aligned} g(x \bullet_{\rightharpoonup} y) &\stackrel{(3.5)}{=} g(x_1 \cdot (x_2 \rightharpoonup y)) = g(x_1) \cdot' g(x_2 \rightharpoonup y) = g(x_1) \cdot' (g(x_2) \rightharpoonup' g(y)) \\ &= g(x) \bullet_{\rightharpoonup'} g(y), \end{aligned}$$

so g is also a morphism of algebras from H_{\rightharpoonup} to $H'_{\rightharpoonup'}$, and the pair $(g : H_{\rightharpoonup} \rightarrow H'_{\rightharpoonup'}, g : H \rightarrow H')$ is a morphism of Yetter–Drinfeld relative Rota–Baxter operators from $\text{Id}_H : H \rightarrow H_{\rightharpoonup}$ to $\text{Id}_{H'} : H' \rightarrow H'_{\rightharpoonup'}$. Clearly, the assignment L respects identities and compositions. ■

Remark 4.9. The previous result can also be deduced in the following way.

Given a Yetter–Drinfeld post-Hopf algebra (H, \dashv) , one obtains the Yetter–Drinfeld brace $(H, \cdot, \bullet_{\dashv}, 1, \Delta, \varepsilon, S, S_{\dashv})$ as in Corollary 3.21. Then, from [12, Theorem 4.2 (ii) to (i)], one obtains the Yetter–Drinfeld 1-cocycle $\text{Id}_H : H_{\dashv} \rightarrow H$, hence, the Yetter–Drinfeld relative Rota–Baxter operator $\text{Id}_H : H \rightarrow H_{\dashv}$.

We can use Proposition 4.8 to obtain examples of Yetter–Drinfeld relative Rota–Baxter operators.

Example 4.10. In Examples 3.29, 3.30, 3.31, and 3.32 we obtained Yetter–Drinfeld post-Hopf algebras that coincide with the transmutation of the Sweedler’s Hopf algebra H_4 , the Hopf algebras $E(n)$, the Hopf algebra $\text{SL}_q(2)$ and the Suzuki Hopf algebra $A_{1,2}^{\nu,\lambda}$, respectively. We denote them as $\underline{H_4}$, $\underline{E(n)}$, $\underline{\text{SL}_q(2)}$, and $\underline{A_{1,2}^{\nu,\lambda}}$. Therefore, using Proposition 4.8, we obtain that

$$\text{Id} : \underline{H_4} \rightarrow H_4, \quad \text{Id} : \underline{E(n)} \rightarrow E(n), \quad \text{Id} : \underline{\text{SL}_q(2)} \rightarrow \text{SL}_q(2), \quad \text{Id} : \underline{A_{1,2}^{\nu,\lambda}} \rightarrow A_{1,2}^{\nu,\lambda}$$

are all examples of Yetter–Drinfeld relative Rota–Baxter operators.

Let \mathcal{D} be the full subcategory of $\text{BYDrRB}(\text{Vec}_{\mathbb{k}})$ whose objects are bijective Yetter–Drinfeld relative Rota–Baxter operators $R : K \rightarrow H$ such that K has coaction given by $(\text{Id}_H \otimes R^{-1})\text{Ad}_L^H R$, where Ad_L^H denotes the left adjoint coaction on H (equivalently, $R : K \rightarrow H$ is left H -colinear by considering H equipped with Ad_L^H).

Remark 4.11. If H is cocommutative, then Ad_L^H is the trivial H -coaction on H , and so, the previous H -coaction on K is trivial. Hence, in this case, $\sigma_{K,K}^{\mathcal{Y}\mathcal{D}}$ becomes the canonical flip $\tau_{K,K}$ and K becomes a standard (cocommutative) Hopf algebra (and an object in $\text{Hopf}(H\mathfrak{M})$). Thus, the subcategory of \mathcal{D} whose objects $R : K \rightarrow H$ are such that H is cocommutative coincides with the category of bijective relative Rota–Baxter operators on cocommutative Hopf algebras, see [22, Definition 3.1].

As it is shown in the following result, we can build a functor $M : \mathcal{D} \rightarrow \mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$.

Proposition 4.12. *Let $R : K \rightarrow H$ be a bijective Yetter–Drinfeld relative Rota–Baxter operator on the Hopf algebra $(H, \cdot, 1, \Delta, \varepsilon, S)$ with respect to (K, \dashv) , where K is in $H\mathfrak{M}$ with coaction given by $(\text{Id}_H \otimes R^{-1})\text{Ad}_L^H R$. Then, $(H, \cdot_R, 1, \Delta, \varepsilon, S_R, \dashv_R)$ is a Yetter–Drinfeld post-Hopf algebra where \cdot_R and S_R are defined, for all $a, b \in H$, by*

$$a \cdot_R b = R(R^{-1}(a) \cdot_K R^{-1}(b)), \quad S_R(a) := R(a_1 \dashv R^{-1}S(a_2)) \tag{4.4}$$

and \dashv_R is defined by

$$a \dashv_R b := R(a \dashv R^{-1}(b)).$$

Proof. Given a bijective Yetter–Drinfeld relative Rota–Baxter operator $R : K \rightarrow H$, we obtain a Yetter–Drinfeld 1-cocycle $R^{-1} : H \rightarrow K$. Thus, we can apply [12, Theorem 4.2 (i) to (ii)] obtaining a Yetter–Drinfeld brace $(H, \cdot_R, \cdot, 1, \Delta, \varepsilon, S_R, S)$, where \cdot_R and S_R

are defined as in (4.4). Then, by Proposition 3.22, we have that $(H, \cdot_R, 1, \Delta, \varepsilon, S_R, \rightarrow_R)$ is a Yetter–Drinfeld post-Hopf algebra where \rightarrow_R is defined by

$$\begin{aligned} a \rightarrow_R b &:= S_R(a_1) \cdot_R (a_2 \cdot b) = R(a_1 \rightarrow R^{-1}S(a_2)) \cdot_R (a_3 \cdot b) \\ &= R(R^{-1}R(a_1 \rightarrow R^{-1}S(a_2))) \cdot_K R^{-1}(a_3 \cdot b) \\ &\stackrel{(!)}{=} R((a_1 \rightarrow R^{-1}S(a_2)) \cdot_K R^{-1}(a_3) \cdot_K (a_4 \rightarrow R^{-1}(b))) \\ &= R(S_K(R^{-1}(a_1)) \cdot_K R^{-1}(a_2) \cdot_K (a_3 \rightarrow R^{-1}(b))) \\ &= R(\varepsilon(a_1)1_K \cdot_K (a_2 \rightarrow R^{-1}(b))) \\ &= R(a \rightarrow R^{-1}(b)), \end{aligned}$$

where the equality marked with (!) follows using that R^{-1} is a Yetter–Drinfeld 1-cocycle. ■

Remark 4.13. Since we know that $(H, \cdot_R, \cdot, 1, \Delta, \varepsilon, S_R, S)$ is a Yetter–Drinfeld brace we immediately have that $H' := (H, \cdot, 1, \Delta, \varepsilon, S)$ is the subadjacent Hopf algebra of $(H, \cdot_R, 1, \Delta, \varepsilon, S_R)$, so that $(H, \cdot_R, 1, \Delta, \varepsilon, S_R)$ is an object in $\text{Hopf}(\overset{H}{H} \mathcal{YD})$.

Let us observe that the functor L given in Proposition 4.8 goes into the category \mathcal{D} . In analogy with [12, Theorem 4.2], using functors L and M , we obtain the following result.

Corollary 4.14. *The category $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$ and the subcategory \mathcal{D} of $\text{BYDrRB}(\text{Vec}_{\mathbb{k}})$ are equivalent.*

Proof. Let $R : K \rightarrow H$ be an object in \mathcal{D} . Then, by Remark 4.13, $LM((R : K \rightarrow H))$ is the Yetter–Drinfeld relative Rota–Baxter operator $\text{Id}_H : (H, \cdot_R, 1, \Delta, \varepsilon, S_R, \rightarrow_R) \rightarrow (H, \cdot, 1, \Delta, \varepsilon, S)$. Since $\text{Id}_H(a) \rightarrow_R R(b) = R(a \rightarrow R^{-1}R(b)) = R(a \rightarrow b)$ and R is a morphism of algebras with respect to the new algebra structure $(H, \cdot_R, 1)$, we have that (Id_H, R) is an isomorphism of Yetter–Drinfeld relative Rota–Baxter operators between $R : K \rightarrow H$ and $LM((R : K \rightarrow H))$ (and it is clearly natural). Hence, $LM \cong \text{Id}$. On the other hand,

$$\begin{aligned} ML((H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow)) &= M(\text{Id}_H : (H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow) \rightarrow (H, \bullet_{\rightarrow}, 1, \Delta, \varepsilon, S_{\rightarrow})) \\ &= (H, \cdot, 1, \Delta, \varepsilon, S, \rightarrow), \end{aligned}$$

since $a_1 \rightarrow S_{\rightarrow}(a_2) = \alpha_{\rightarrow} a_1 (\beta_{\rightarrow} a_2 (S(a_3))) = S(a)$. Thus, $ML = \text{Id}$. ■

If we restrict the previous equivalence, we obtain an equivalence between cocommutative post-Hopf algebras and bijective relative Rota–Baxter operators on cocommutative Hopf algebras.

Remark 4.15. If we restrict the functor L to cocommutative Yetter–Drinfeld post-Hopf algebras, we recover the functor given in [22, Proposition 3.3] from the category of cocommutative post-Hopf algebras to the category of (bijective) relative Rota–Baxter operators on cocommutative Hopf algebras. Moreover, in [22, Theorem 3.4] it is proven that this

functor has a right adjoint. We want to recover this result from a different equivalence between the category of Yetter–Drinfeld post-Hopf algebras and the subcategory \mathcal{D} of the category of bijective Yetter–Drinfeld relative Rota–Baxter operators.

Given a bijective Yetter–Drinfeld relative Rota–Baxter operator $R : K \rightarrow H$ in \mathcal{D} , we want to induce a structure of Yetter–Drinfeld post-Hopf algebra on K .

Proposition 4.16. *Let $R : K \rightarrow H$ be an object in \mathcal{D} . Then, (K, \rightarrow_R) is a Yetter–Drinfeld post-Hopf algebra where \rightarrow_R is given by*

$$a \rightarrow_R b := R(a) \rightarrow b \quad \text{for all } a, b \in K.$$

Let $R : K \rightarrow H$ and $R' : K' \rightarrow H'$ be objects in \mathcal{D} and $(f : H \rightarrow H', g : K \rightarrow K')$ a morphism between them. Then, g is a morphism of Yetter–Drinfeld post-Hopf algebras from (K, \rightarrow_R) to $(K', \rightarrow_{R'})$. This yields a functor $\mathbb{R} : \mathcal{D} \rightarrow \mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$.

Proof. Let us observe that $\rightarrow_R := \rightarrow (R \otimes \text{Id}_K)$. Thus, since R and \rightarrow are morphisms of coalgebras, also \rightarrow_R is a morphism of coalgebras. Moreover, we already know that $(K, \cdot, 1)$ is an algebra and (K, Δ, ε) is a coalgebra and $S_K : K \rightarrow K$ satisfies $a_1 \cdot S_K(a_2) = \varepsilon(a)1_K = S_K(a_1) \cdot a_2$, for all $k \in K$. Moreover, since \rightarrow satisfies (3.1), the same holds for \rightarrow_R :

$$a \rightarrow_R (b \cdot c) = R(a) \rightarrow (b \cdot c) \stackrel{(3.1)}{=} (R(a_1) \rightarrow b) \cdot (R(a_2) \rightarrow c) = (a_1 \rightarrow_R b) \cdot (a_2 \rightarrow_R c).$$

We already know that \rightarrow is an H -action on K , so also (3.2) is satisfied:

$$\begin{aligned} a \rightarrow_R (b \rightarrow_R c) &= R(a) \rightarrow (R(b) \rightarrow c) = (R(a) \cdot R(b)) \rightarrow c \\ &\stackrel{(4.1)}{=} R(a_1 \cdot (R(a_2) \rightarrow b)) \rightarrow c = (a_1 \cdot (a_2 \rightarrow_R b)) \rightarrow_R c. \end{aligned}$$

For all $a, b \in K$, define $(\beta_{\rightarrow_R} a)(b) := S_H(R(a)) \rightarrow b$, so that

$$\begin{aligned} \beta_{\rightarrow_R} a_1 (\alpha_{\rightarrow_R} a_2 (b)) &= S_H R(a_1) \rightarrow (R(a_2) \rightarrow b) = (S_H(R(a_1)) \cdot R(a_2)) \rightarrow b \\ &= \varepsilon(R(a))1 \rightarrow b = \varepsilon(a)b, \end{aligned}$$

hence, $(\beta_{\rightarrow_R} a_1) \circ (\alpha_{\rightarrow_R} a_2) = \varepsilon(a)\text{Id}_K$ for all $a \in K$. Similarly, $(\alpha_{\rightarrow_R} a_1) \circ (\beta_{\rightarrow_R} a_2) = \varepsilon(a)\text{Id}_K$, for all $a \in K$. We already know that $\varepsilon(a \cdot b) = \varepsilon(a)\varepsilon(b)$ for all $a, b \in K$, $\varepsilon(1_K) = 1_{\mathbb{k}}$ and $\Delta(1_K) = 1_K \otimes 1_K$. Moreover, since K is in $\text{Bimon}(\overset{H}{\mathcal{YD}})$ and it is in ${}^H\mathfrak{M}$ with $(\text{Id}_H \otimes R^{-1})\text{Ad}_L^H R$, we have

$$\sigma_{K,K}^{\mathcal{YD}}(a \otimes b) := (R(a_1) \cdot S_H(R(a_3)) \rightarrow b) \otimes a_2,$$

so that

$$\begin{aligned} \Delta(a \cdot b) &= (m \otimes m)(\text{Id}_K \otimes \sigma_{K,K}^{\mathcal{YD}} \otimes \text{Id}_K)(\Delta \otimes \Delta)(a \otimes b) \\ &= a_1 \cdot (R(a_2) \cdot S_H(R(a_4)) \rightarrow b_1) \otimes (a_3 \cdot b_2) \\ &= a_1 \cdot (R(a_2) \rightarrow (S_H(R(a_4)) \rightarrow b_1)) \otimes (a_3 \cdot b_2) \\ &= a_1 \cdot (\alpha_{\rightarrow_R} a_2 (\beta_{\rightarrow_R} a_4 (b_1))) \otimes (a_3 \cdot b_2), \end{aligned}$$

hence, (3.4) holds true. Furthermore, (3.5) and (3.6) become

$$a \bullet_{\rightarrow_R} b := a_1 \cdot (a_2 \rightarrow_R b) = a_1 \cdot (R(a_2) \rightarrow b) \stackrel{(4.1)}{=} R^{-1}(R(a) \cdot R(b))$$

and

$$S_{\rightarrow_R}(a) := \beta_{\rightarrow_R} a_1(S_K(a_2)) = S_H(R(a_1)) \rightarrow S_K(a_2).$$

But, by Lemma 4.5, we know that $S_K(a) = R(a_1) \rightarrow R^{-1}S_H R(a_2)$ and then

$$\begin{aligned} S_{\rightarrow_R}(a) &= S_H(R(a_1)) \rightarrow (R(a_2) \rightarrow R^{-1}S_H R(a_3)) \\ &= (S_H(R(a_1)) \cdot R(a_2)) \rightarrow R^{-1}S_H R(a_3) = R^{-1}S_H R(a). \end{aligned}$$

Thus, $S_{\rightarrow_R} = R^{-1}S_H R$, and since R and R^{-1} are comultiplicative and S_H is anti-comultiplicative, S_{\rightarrow_R} is anti-comultiplicative. Finally, (3.7) becomes

$$\begin{aligned} a \leftarrow_R b &:= S_{\rightarrow_R}(a_1 \rightarrow_R b_1) \bullet_{\rightarrow_R} a_2 \bullet_{\rightarrow_R} b_2 \\ &= R^{-1}(RS_{\rightarrow_R}(a_1 \rightarrow_R b_1) \cdot R(a_2) \cdot R(b_2)) \\ &= R^{-1}(S_H R(R(a_1) \rightarrow b_1) \cdot R(a_2) \cdot R(b_2)). \end{aligned}$$

So, we can compute

$$\begin{aligned} (a_1 \leftarrow_R b_1) \otimes R(a_2 \rightarrow_R b_2) &= R^{-1}(S_H R(R(a_1) \rightarrow b_1) \cdot R(a_2) \cdot R(b_2)) \otimes R(R(a_3) \rightarrow b_3) \\ &\stackrel{(4.2)}{=} R^{-1}(S_H R(R(a_2) \rightarrow b_2) \cdot R(a_3) \cdot R(b_3)) \otimes R(R(a_1) \rightarrow b_1) \\ &= (a_2 \leftarrow_R b_2) \otimes R(a_1 \rightarrow_R b_1). \end{aligned}$$

Thus, since R is injective, also (2.9) is satisfied and (K, \rightarrow_R) is a Yetter–Drinfeld post-Hopf algebra.

Let $(f : H \rightarrow H', g : K \rightarrow K')$ be a morphism of Yetter–Drinfeld relative Rota–Baxter operators from $R : K \rightarrow H$ to $R' : K' \rightarrow H'$. In particular, g is a morphism of algebras and coalgebras; moreover, it satisfies

$$g(a \rightarrow_R b) = g(R(a) \rightarrow b) = f(R(a)) \rightarrow' g(b) = R'(g(a)) \rightarrow' g(b) = g(a) \rightarrow'_{R'} g(b),$$

so g is a morphism of Yetter–Drinfeld post-Hopf algebras from (K, \rightarrow_R) to $(K', \rightarrow'_{R'})$. Clearly, the assignment R respects identities and compositions. ■

Using the functors L and R , we obtain another equivalence between $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$ and the subcategory \mathcal{D} of $\text{BYDrRB}(\text{Vec}_{\mathbb{k}})$ which again restricts to an equivalence between cocommutative post-Hopf algebras and bijective relative Rota–Baxter operators on cocommutative Hopf algebras.

Corollary 4.17. *The categories $\mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$ and \mathcal{D} are equivalent.*

Proof. Given a Yetter–Drinfeld post-Hopf algebra (H, \dashv) we have $RL((H, \dashv)) = R(\text{Id}_H : H \rightarrow H_{\dashv}) = (H, \dashv)$, so $RL = \text{Id}$. On the other hand, given $R : K \rightarrow H$ in \mathcal{D} , we have

$$LR((R : K \rightarrow H)) = L((K, \dashv_R)) = (\text{Id}_K : (K, \dashv_R) \rightarrow K_R).$$

The pair $(R^{-1} : H \rightarrow K_R, \text{Id}_K)$ clearly, satisfies (4.3): indeed, $R^{-1}(a) \dashv_R \text{Id}_K(b) = RR^{-1}(a) \dashv b = \text{Id}_K(a \dashv b)$. Moreover, R^{-1} is a morphism of coalgebras and

$$R^{-1}(a) \bullet_{\dashv_R} R^{-1}(b) = R^{-1}(a_1) \cdot (a_2 \dashv R^{-1}(b)) = R^{-1}(a \cdot b),$$

where the last equality follows since R^{-1} is a Yetter–Drinfeld 1-cocycle. Thus, (R^{-1}, Id_K) is an isomorphism (which is clearly natural), and so, $LR \cong \text{Id}$. ■

Remark 4.18. Notice that the bijectivity of $R : K \rightarrow H$ is not used in the first part of Proposition 4.16. Hence, one can try to obtain a similar result considering the category $\mathcal{YDrRB}(\text{Vec}_{\mathbb{k}})$ of Yetter–Drinfeld relative Rota–Baxter operators and its subcategory \mathcal{C} whose objects $R : K \rightarrow H$ are such that K is in $\text{Hopf}(\mathop{H}\limits^H \mathcal{YD})$.

Define the full subcategory \mathcal{C}' of \mathcal{C} whose objects are injective Yetter–Drinfeld relative Rota–Baxter operators $R : K \rightarrow H$ such that K has H -coaction given by $a \mapsto R(a_1)S_H(R(a_3)) \otimes a_2$.

Remark 4.19. The difference between the categories \mathcal{C}' and \mathcal{D} is simply that an object $R : K \rightarrow H$ in \mathcal{C}' is not necessarily invertible; the category \mathcal{D} is a subcategory of \mathcal{C}' . Moreover, the subcategory of \mathcal{C}' whose objects $R : K \rightarrow H$ are such that H is cocommutative (so also K is cocommutative since R is injective) is the category of injective relative Rota–Baxter operators on cocommutative Hopf algebras, see [22, Definition 3.1].

Notice that the functor L of Proposition 4.8 goes into the category \mathcal{C}' . We can then show the following result.

Proposition 4.20. *Let $R : K \rightarrow H$ be an object in \mathcal{C}' . Then, (K, \dashv_R) is a Yetter–Drinfeld post-Hopf algebra where \dashv_R is defined by $a \dashv_R b := R(a) \dashv b$, for all $a, b \in K$. Consider also the same assignment on morphisms of the functor R given in Proposition 4.16. This yields a functor $R' : \mathcal{C}' \rightarrow \mathcal{YDPH}(\text{Vec}_{\mathbb{k}})$, which coincides with R when it is restricted to \mathcal{D} . Moreover, the functor R' is a right adjoint of the functor L .*

Proof. The first part of the proof of Proposition 4.16 remains true in this setting. Then, we have

$$a \bullet_{\dashv_R} b := a_1 \cdot (R(a_2) \dashv b), \quad S_{\dashv_R}(a) := S_H(R(a_1)) \dashv S_K(a_2).$$

In order to obtain that (K, \dashv_R) is a Yetter–Drinfeld post-Hopf algebra it only remains to prove that S_{\dashv_R} is anti-comultiplicative and the pair $(\dashv_R, \dashleftarrow_R)$ satisfies (2.9), where

$$a \dashleftarrow_R b := S_{\dashv_R}(a_1 \dashv_R b_1) \bullet_{\dashv_R} a_2 \bullet_{\dashv_R} b_2.$$

Observe that, in terms of $\bullet \rightarrow_R$, (4.1) reads $R(a) \cdot R(b) = R(a \bullet \rightarrow_R b)$ and then (4.2) becomes

$$\begin{aligned} S_H R(a_1 \rightarrow_R b_1) \cdot R(a_2 \bullet \rightarrow_R b_2) \otimes R(a_3 \rightarrow_R b_3) \\ = S_H R(a_2 \rightarrow_R b_2) \cdot R(a_3 \bullet \rightarrow_R b_3) \otimes R(a_1 \rightarrow_R b_1). \end{aligned} \tag{4.5}$$

Let us observe that

$$\begin{aligned} R(a_1) \rightarrow S_{\rightarrow_R}(a_2) &= R(a_1) \rightarrow (S_H R(a_2) \rightarrow S_K(a_3)) \\ &= (R(a_1) \cdot S_H(R(a_2))) \rightarrow S_K(a_3) = S_K(a), \end{aligned} \tag{4.6}$$

and so,

$$R(a_1) \cdot R(S_{\rightarrow_R}(a_2)) \stackrel{(4.1)}{=} R(a_1 \cdot (R(a_2) \rightarrow S_{\rightarrow_R}(a_3))) = R(a_1 \cdot S_K(a_2)) = \varepsilon(a)1_H. \tag{4.7}$$

Thus, we obtain

$$R(S_{\rightarrow_R}(a)) = S_H(R(a_1)) \cdot R(a_2) \cdot R(S_{\rightarrow_R}(a_3)) \stackrel{(4.7)}{=} S_H(R(a)), \tag{4.8}$$

i.e., $RS_{\rightarrow_R} = S_H R$. Therefore, since R is injective and comultiplicative and S_H is anti-comultiplicative, S_{\rightarrow_R} is anti-comultiplicative. Moreover, (4.5) becomes

$$\begin{aligned} R(S_{\rightarrow_R}(a_1 \rightarrow_R b_1) \bullet \rightarrow_R a_2 \bullet \rightarrow_R b_2) \otimes R(a_3 \rightarrow_R b_3) \\ = R(S_{\rightarrow_R}(a_2 \rightarrow_R b_2) \bullet \rightarrow_R a_3 \bullet \rightarrow_R b_3) \otimes R(a_1 \rightarrow_R b_1), \end{aligned}$$

which is exactly

$$R(a_1 \leftarrow_R b_1) \otimes R(a_2 \rightarrow_R b_2) = R(a_2 \leftarrow_R b_2) \otimes R(a_1 \rightarrow_R b_1).$$

Since R is injective we get that the pair $(\rightarrow_R, \leftarrow_R)$ satisfies (2.9), hence, (K, \rightarrow_R) is a Yetter–Drinfeld post-Hopf algebra. As in the proof of Proposition 4.16 one concludes that R' is a functor.

Finally, we just have to prove that, for every $R : K \rightarrow H$ in \mathcal{C}' and (H', \rightarrow') in $\mathcal{YDPH}(\text{Vec}_{\mathbb{K}})$, there is a natural bijection between $\text{Hom}(L(H', \rightarrow'), R : K \rightarrow H)$ and $\text{Hom}((H', \rightarrow'), R'(R : K \rightarrow H))$, i.e., between $\text{Hom}(\text{Id}_{H'} : H' \rightarrow H'_{\rightarrow'}, R : K \rightarrow H)$ and $\text{Hom}((H', \rightarrow'), (K, \rightarrow_R))$. Given a morphism $(f : H'_{\rightarrow'} \rightarrow H, g : H' \rightarrow K)$ in $\text{Hom}(\text{Id}_{H'} : H' \rightarrow H'_{\rightarrow'}, R : K \rightarrow H)$ we take g as a morphism in $\text{Hom}((H', \rightarrow'), (K, \rightarrow_R))$. Indeed, g is a morphism of algebras and coalgebras; moreover, since $f = f \text{Id}_{H'} = Rg$, we have

$$g(a \rightarrow' b) = f(a) \rightarrow_K g(b) = R(g(a)) \rightarrow_K g(b) = g(a) \rightarrow_R g(b).$$

On the other hand, given $g : (H', \rightarrow') \rightarrow (K, \rightarrow_R)$ in $\text{Hom}((H', \rightarrow'), (K, \rightarrow_R))$ we take $(Rg : H'_{\rightarrow'} \rightarrow H, g : H' \rightarrow K)$ as a morphism in $\text{Hom}(\text{Id}_{H'} : H' \rightarrow H'_{\rightarrow'}, R : K \rightarrow H)$.

Indeed, $g : H' \rightarrow K$ is a morphism of algebras and coalgebras, so Rg is a morphism of coalgebras. Moreover, Rg is also a morphism of algebras

$$\begin{aligned} Rg(a \bullet_{\rightarrow} b) &= Rg(a_1 \cdot (a_2 \dashrightarrow b)) = R(g(a_1) \cdot g(a_2 \dashrightarrow b)) \\ &= R(g(a_1) \cdot (g(a_2) \dashrightarrow_R g(b))) \\ &= R(g(a_1) \cdot (Rg(a_2) \dashrightarrow g(b))) = Rg(a) \cdot Rg(b) \end{aligned}$$

and (4.3) is trivially satisfied. Clearly, the two assignments are inverse to each other and the bijection is natural. ■

In the cocommutative setting, we obtain an adjunction between cocommutative post-Hopf algebras and injective relative Rota–Baxter operators on cocommutative Hopf algebras. This is exactly the adjunction given in [22, Theorem 3.4], which is given for relative Rota–Baxter operators on cocommutative Hopf algebras not necessarily injective. These can be seen as objects $R : K \rightarrow H$ in \mathcal{C} where K and H are cocommutative and the H -coaction on K is given by $a \mapsto R(a_1)S_H(R(a_3)) \otimes a_2$: an example is given by the extension of a relative Rota–Baxter operator on a Lie algebra to the respective universal enveloping algebras, as it is proved in [22, Theorem 3.6].

Remark 4.21. Assume the hypotheses of the previous result are satisfied. By Proposition 3.10, we have that $(K, \bullet_{\rightarrow_R}, 1, \Delta, \varepsilon, S_{\rightarrow_R})$ is a Hopf algebra, where

$$a \bullet_{\rightarrow_R} b := a_1 \cdot (R(a_2) \dashrightarrow b), \quad S_{\rightarrow_R}(a) := S_H R(a_1) \dashrightarrow S_K(a_2),$$

which we call the *descendent Hopf algebra* K_R in analogy with [22, Corollary 3.5]. We automatically have that $(K, \cdot, \bullet_{\rightarrow_R}, 1, \Delta, \varepsilon, S_K, S_{\rightarrow_R})$ is a Yetter–Drinfeld brace by Corollary 3.21; this result generalises [31, Proposition 3.16]. Moreover, using Corollary 3.25, one obtains a matched pair of actions $(K_R, \dashrightarrow_R, \dashleftarrow_R)$, where $a \dashleftarrow_R b := S_{\rightarrow_R}(a_1 \dashrightarrow_R b_1) \bullet_{\rightarrow_R} a_2 \bullet_{\rightarrow_R} b_2$; this result generalises [22, Corollary 4.10].

Acknowledgements. The author would like to thank A. Ardizzoni for useful comments and D. Ferri for his interest in this paper and a nice discussion regarding possible future developments. Furthermore, he would like to express his gratitude to the referee for meaningful suggestions.

Funding. This paper was written while the author was member of the “National Group for Algebraic and Geometric Structures and their Applications” (GNSAGA-INdAM). He was also partially supported by the project funded by the European Union -NextGenerationEU under NRRP, Mission 4 Component 2 CUP D53D23005960006 - Call PRIN 2022 no. 104 of February 2, 2022 of Italian Ministry of University and Research; Project 2022S97PMY Structures for Quivers, Algebras and Representations (SQUARE).

References

- [1] M. J. H. Al-Kaabi, K. Ebrahimi-Fard, and D. Manchon, [Free post-groups, post-groups from group actions, and post-Lie algebras](#). *J. Geom. Phys.* **198** (2024), article no. 105129 Zbl 1536.16030 MR 4705107
- [2] I. Angiono, C. Galindo, and L. Vendramin, [Hopf braces and Yang–Baxter operators](#). *Proc. Amer. Math. Soc.* **145** (2017), no. 5, 1981–1995 Zbl 1392.16032 MR 3611314
- [3] C. Bai, L. Guo, and X. Ni, [Nonabelian generalized Lax pairs, the classical Yang–Baxter equation and PostLie algebras](#). *Comm. Math. Phys.* **297** (2010), no. 2, 553–596 Zbl 1206.17020 MR 2651910
- [4] C. Bai, L. Guo, Y. Sheng, and R. Tang, [Post-groups, \(Lie-\)Butcher groups and the Yang–Baxter equation](#). *Math. Ann.* **388** (2024), no. 3, 3127–3167 Zbl 1535.22061 MR 4705761
- [5] V. G. Bardakov and V. Gubarev, [Rota–Baxter groups, skew left braces, and the Yang–Baxter equation](#). *J. Algebra* **596** (2022), 328–351 Zbl 1492.17019 MR 4370524
- [6] F. Borceux, [Handbook of categorical algebra. I. Basic category theory](#). Encyclopedia Math. Appl. 50, Cambridge University Press, Cambridge, 1994 Zbl 0803.18001 MR 1291599
- [7] Y. Bruned and F. Katssetsiadis, [Post-Lie algebras in regularity structures](#). *Forum Math. Sigma* **11** (2023), article no. e98 Zbl 1540.60235 MR 4662283
- [8] F. Chapoton and M. Livernet, [Pre-Lie algebras and the rooted trees operad](#). *Internat. Math. Res. Notices* (2001), no. 8, 395–408 Zbl 1053.17001 MR 1827084
- [9] A. Connes and D. Kreimer, [Renormalization in quantum field theory and the Riemann–Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem](#). *Comm. Math. Phys.* **210** (2000), no. 1, 249–273 Zbl 1032.81026 MR 1748177
- [10] C. Curry, K. Ebrahimi-Fard, and B. Owren, [The Magnus expansion and post-Lie algebras](#). *Math. Comp.* **89** (2020), no. 326, 2785–2799 Zbl 1450.34015 MR 4136547
- [11] K. Ebrahimi-Fard, A. Lundervold, and H. Z. Munthe-Kaas, [On the Lie enveloping algebra of a post-Lie algebra](#). *J. Lie Theory* **25** (2015), no. 4, 1139–1165 Zbl 1360.17015 MR 3350091
- [12] D. Ferri and A. Sciandra, [Matched pairs and Yetter–Drinfeld braces](#). *Can. J. Math.* (2025), DOI 10.4153/S0008414X25000239
- [13] M. Goncharov, [Rota–Baxter operators on cocommutative Hopf algebras](#). *J. Algebra* **582** (2021), 39–56 Zbl 1505.17012 MR 4256904
- [14] L. Guarnieri and L. Vendramin, [Skew braces and the Yang–Baxter equation](#). *Math. Comp.* **86** (2017), no. 307, 2519–2534 Zbl 1371.16037 MR 3647970
- [15] J. A. Guccione, J. J. Guccione, and C. Valqui, [Set-theoretic type solutions of the braid equation](#). [v1] 2020, [v5] 2024, arXiv:2008.13494v5
- [16] J. A. Guccione, J. J. Guccione, and L. Vendramin, [Yang–Baxter operators in symmetric categories](#). *Comm. Algebra* **46** (2018), no. 7, 2811–2845 Zbl 1430.16037 MR 3780826
- [17] L. Guo, H. Lang, and Y. Sheng, [Integration and geometrization of Rota–Baxter Lie algebras](#). *Adv. Math.* **387** (2021), article no. 107834 Zbl 1468.17026 MR 4271483
- [18] D. Kreimer, [On the Hopf algebra structure of perturbative quantum field theories](#). *Adv. Theor. Math. Phys.* **2** (1998), no. 2, 303–334 Zbl 1041.81087 MR 1633004
- [19] B. A. Kupershmidt, [What a classical \$r\$ -matrix really is](#). *J. Nonlinear Math. Phys.* **6** (1999), no. 4, 448–488 Zbl 1015.17015 MR 1722068
- [20] Y. Li, [Matched pairs and Yang–Baxter operators](#). 2025, arXiv:2501.11975v2
- [21] Y. Li, Y. Sheng, and R. Tang, [Post-Hopf algebras, relative Rota–Baxter operators and solutions of the Yang–Baxter equation](#). 2022, arXiv:2203.12174v1

- [22] Y. Li, Y. Sheng, and R. Tang, [Post-Hopf algebras, relative Rota–Baxter operators and solutions to the Yang–Baxter equation](#). *J. Noncommut. Geom.* **18** (2024), no. 2, 605–630
Zbl [07828328](#) MR [4721043](#)
- [23] S. Majid, [Transmutation theory and rank for quantum braided groups](#). *Math. Proc. Cambridge Philos. Soc.* **113** (1993), no. 1, 45–70 Zbl [0781.17006](#) MR [1188817](#)
- [24] S. Majid, *Foundations of quantum group theory*. Cambridge University Press, Cambridge, 1995 Zbl [0857.17009](#) MR [1381692](#)
- [25] H. Z. Munthe-Kaas and A. Lundervold, [On post-Lie algebras, Lie–Butcher series and moving frames](#). *Found. Comput. Math.* **13** (2013), no. 4, 583–613 Zbl [1327.17001](#) MR [3085679](#)
- [26] J.-M. Oudom and D. Guin, [On the Lie enveloping algebra of a pre-Lie algebra](#). *J. K-Theory* **2** (2008), no. 1, 147–167 Zbl [1178.17011](#) MR [2434170](#)
- [27] D. E. Radford, *Hopf algebras*. Ser. Knots Everything 49, World Scientific, Hackensack, NJ, 2012 Zbl [1266.16036](#) MR [2894855](#)
- [28] W. Rump, [Braces, radical rings, and the quantum Yang–Baxter equation](#). *J. Algebra* **307** (2007), no. 1, 153–170 Zbl [1115.16022](#) MR [2278047](#)
- [29] M. E. Sweedler, *Hopf algebras*. Mathematics Lecture Note Series, W. A. Benjamin, New York, 1969 Zbl [0194.32901](#) MR [0252485](#)
- [30] B. Vallette, [Homology of generalized partition posets](#). *J. Pure Appl. Algebra* **208** (2007), no. 2, 699–725 Zbl [1109.18002](#) MR [2277706](#)
- [31] H. Zheng, L. Guo, T. Ma, and L. Zhang, [Rota–Baxter operators on cocommutative Hopf algebras and Hopf braces](#). [v1] 2023, [v2] 2024, arXiv:[2304.07684v2](#)

Received 19 August 2024; revised 20 March 2025.

Andrea Sciandra

Department of Mathematics “G. Peano”, University of Turin, Via Carlo Alberto 10, 10123 Torino, Italy; andrea.sciandra@unito.it