



Manor Mendel · Assaf Naor

Markov convexity and local rigidity of distorted metrics

Abstract. It is shown that a Banach space admits an equivalent norm whose modulus of uniform convexity has power-type p if and only if it is Markov p -convex. Counterexamples are constructed to natural questions related to isomorphic uniform convexity of metric spaces, showing in particular that tree metrics fail to have the dichotomy property.

Contents

1. Introduction	287
2. Markov p -convexity and p -convexity coincide	296
3. A doubling space which is not Markov p -convex for any $p \in (0, \infty)$	302
4. Lipschitz quotients	304
5. A dichotomy theorem for vertically faithful embeddings of trees	306
6. Tree metrics do not have the dichotomy property	309
7. Discussion and open problems	335
References	335

1. Introduction

A Banach space $(X, \|\cdot\|_X)$ is said to be *finitely representable* in a Banach space $(Y, \|\cdot\|_Y)$ if there exists a constant $D < \infty$ such that for every finite-dimensional linear subspace $F \subseteq X$ there is a linear operator $T : F \rightarrow Y$ satisfying $\|x\|_X \leq \|Tx\|_Y \leq D\|x\|_X$ for all $x \in F$. In 1976 Ribe [31] proved that if two Banach spaces X and Y are *uniformly homeomorphic*, i.e., there is a bijection $f : X \rightarrow Y$ such that f and f^{-1} are uniformly continuous, then X is finitely representable in Y and vice versa. This remarkable theorem motivated what is known today as the “Ribe program”: the search for purely metric reformulations of basic linear concepts and invariants from the local theory of Banach spaces. This research program was put forth by Bourgain in 1986 [5].

Since its inception, the Ribe program attracted the work of many mathematicians, and led to the development of several satisfactory metric theories that extend important

M. Mendel: Mathematics and Computer Science Department, The Open University of Israel,
1 University Rd., P.O. Box 808, Raanana 43107, Israel; e-mail: manorme@openu.ac.il

A. Naor: Courant Institute of Mathematical Sciences, New York University, 251 Mercer Street,
New York, NY 10012, USA; e-mail: naor@cims.nyu.edu

concepts and results of Banach space theory; see the introduction of [24] for a historical discussion. So far, progress on the Ribe program has come hand-in-hand with applications to metric geometry, group theory, functional analysis, and computer science.

The present paper contains further progress in this direction: we obtain a metric characterization of p -convexity in Banach spaces, derive some of its metric consequences, and construct unexpected counterexamples which indicate that further progress on the Ribe program can uncover nonlinear phenomena that are markedly different from their Banach space counterparts. In doing so, we answer questions posed by Lee–Naor–Peres and Feferman, and improve a theorem of Bates, Johnson, Lindenstrauss, Preiss and Schechtman. These results, which will be explained in detail below, were announced in [23].

For $p \geq 2$, a Banach space $(X, \|\cdot\|_X)$ is said to be p -convex if there exists a norm $\|\!\| \cdot \|\!\|$ which is equivalent to $\|\cdot\|_X$ (i.e., for some $a, b > 0$, $a\|x\|_X \leq \|\!\|x\|\!\| \leq b\|x\|_X$ for all $x \in X$), and a constant $K > 0$ satisfying

$$\|\!\|x\|\!\| = \|\!\|y\|\!\| = 1 \Rightarrow \left\| \frac{x+y}{2} \right\| \leq 1 - K\|\!\|x-y\|\!\|^p. \quad (1)$$

X is called *superreflexive* if it is p -convex for some $p \geq 2$ (historically, this is not the original definition of superreflexivity,¹ but it is equivalent to it due to a deep theorem of Pisier [29], which builds on important work of James [10] and Enflo [7]). For concreteness, we recall (see, e.g., [2]) that L_p is 2-convex for $p \in (1, 2]$ and p -convex for $p \in [2, \infty)$.

Ribe's theorem implies that p -convexity, and hence also superreflexivity, is preserved under uniform homeomorphisms. The first major success of the Ribe program is a famous theorem of Bourgain [5] which gives a metrical characterization of superreflexivity:

Theorem 1.1 (Bourgain's metrical characterization of superreflexivity [5]). *Let B_n be the complete unweighted binary tree of depth n , equipped with the natural graph-theoretical metric. Then a Banach space X is superreflexive if and only if*

$$\lim_{n \rightarrow \infty} c_X(B_n) = \infty. \quad (2)$$

Here, and in what follows, given two metric spaces $(\mathcal{M}, d_{\mathcal{M}})$, $(\mathcal{N}, d_{\mathcal{N}})$, the parameter $c_{\mathcal{M}}(\mathcal{N})$ denotes the smallest bi-Lipschitz distortion with which \mathcal{N} embeds into \mathcal{M} , i.e., the infimum of those $D > 0$ such that there exists a scaling factor $r > 0$ and a mapping $f : \mathcal{N} \rightarrow \mathcal{M}$ satisfying $rd_{\mathcal{N}}(x, y) \leq d_{\mathcal{M}}(f(x), f(y)) \leq Drd_{\mathcal{N}}(x, y)$ for all $x, y \in \mathcal{N}$ (if no such f exists then we set $c_{\mathcal{M}}(\mathcal{N}) = \infty$).

Bourgain's theorem characterizes superreflexivity of Banach spaces in terms of their metric structure, but it leaves open the characterization of p -convexity. The notion of p -convexity is crucial for many applications in Banach space theory and metric geometry,

¹ James' original definition of superreflexivity is that a Banach space X is superreflexive if "its local structure forces reflexivity", i.e., if every Banach space Y that is finitely representable in X must be reflexive. Enflo's renorming theorem states that superreflexivity is equivalent to having an equivalent norm $\|\!\| \cdot \|\!\|$ that is *uniformly convex*, i.e., for every $\varepsilon \in (0, 1)$ there exists $\delta > 0$ such that if $\|\!\|x\|\!\| = \|\!\|y\|\!\| = 1$ and $\|\!\|x-y\|\!\| = \varepsilon$ then $\|\!\|x+y\|\!\| \leq 2-\delta$.

and it turns out that the completion of the Ribe program for p -convexity requires significant additional work beyond Bourgain’s superreflexivity theorem. As a first step in this direction, Lee, Naor and Peres [16] defined a bi-Lipschitz invariant of metric spaces called *Markov convexity*, which is motivated by Ball’s notion of Markov type [1] and Bourgain’s argument in [5].

Definition 1.2 ([16]). Let $\{X_t\}_{t \in \mathbb{Z}}$ be a Markov chain on a state space Ω . Given an integer $k \geq 0$, we denote by $\{\tilde{X}_t(k)\}_{t \in \mathbb{Z}}$ the process which equals X_t for time $t \leq k$, and evolves independently (with respect to the same transition probabilities) for time $t > k$. Fix $p > 0$. A metric space (X, d_X) is called *Markov p -convex with constant Π* if for every Markov chain $\{X_t\}_{t \in \mathbb{Z}}$ on a state space Ω , and every $f : \Omega \rightarrow X$,

$$\sum_{k=0}^{\infty} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_X(f(X_t), f(\tilde{X}_t(t - 2^k)))^p]}{2^{kp}} \leq \Pi^p \cdot \sum_{t \in \mathbb{Z}} \mathbb{E}[d_X(f(X_t), f(X_{t-1}))^p]. \tag{3}$$

The least constant Π for which (3) holds for all Markov chains is called the *Markov p -convexity constant* of X , and is denoted $\Pi_p(X)$. We shall say that (X, d_X) is *Markov p -convex* if $\Pi_p(X) < \infty$.

To gain intuition for Definition 1.2, consider the standard downward random walk starting from the root of the binary tree B_n (with absorbing states at the leaves). For an arbitrary mapping f from B_n to a metric space (X, d_X) , the triangle inequality implies that for each $k \in \mathbb{N}$ we have

$$\sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_X(f(X_t), f(\tilde{X}_t(t - 2^k)))^p]}{2^{kp}} \lesssim_p \sum_{t \in \mathbb{Z}} \mathbb{E}[d_X(f(X_t), f(X_{t-1}))^p], \tag{4}$$

with asymptotic equality (up to constants depending only on p) for $k \leq (\log n)/2$ when $X = B_n$ and f is the identity mapping. On the other hand, if X is a Markov p -convex space then the sum over k of the left-hand side of (4) is *uniformly bounded* by the right-hand side of (4), and therefore Markov p -convex spaces cannot contain B_n with distortion uniformly bounded in n .

We refer to [16] for more information on the notion of Markov p -convexity. In particular, it is shown in [16] that the Markov 2-convexity constant of an arbitrary weighted tree T is, up to constant factors, the Euclidean distortion of T . We refer to [16] for L_p versions of this statement and their algorithmic applications. It was also shown in [16], via a modification of an argument of Bourgain [5], that if a Banach space X is p -convex then it is also Markov p -convex. It was asked in [16] if the converse is also true. Here we answer this question positively:

Theorem 1.3. *A Banach space is p -convex if and only if it is Markov p -convex.*

Thus Markov p -convexity is equivalent to p -convexity in Banach spaces, completing the Ribe program in this case. Our proof of Theorem 1.3 is based on a renorming method of Pisier [29]. It can be viewed as a nonlinear variant of Pisier’s argument, and several subtle changes are required in order to adapt it to a nonlinear condition such as (3).

Results similar to Theorem 1.3 have been obtained for the notions of type and cotype of Banach spaces (see [6, 30, 1, 25, 24, 22]), and have been used to transfer some of the linear theory to the setting of general metric spaces. This led to several applications to problems in metric geometry. Apart from the applications of Markov p -convexity that were obtained in [16], here we show that this invariant is preserved under Lipschitz quotients. The notion of Lipschitz quotient was introduced by Gromov [8, Sec. 1.25]. Given two metric spaces (X, d_X) and (Y, d_Y) , a surjective mapping $f : X \rightarrow Y$ is called a *Lipschitz quotient* if it is Lipschitz, and it is also “Lipschitzly open” in the sense that there exists a constant $c > 0$ such that for every $x \in X$ and $r > 0$,

$$f(B_X(x, r)) \supseteq B_Y(f(x), r/c). \quad (5)$$

Here we show the following result:

Theorem 1.4. *If (X, d_X) is Markov p -convex and (Y, d_Y) is a Lipschitz quotient of X , then Y is also Markov p -convex.*

In [3] Bates, Johnson, Lindenstrauss, Preiss and Schechtman investigated in detail Lipschitz quotients of Banach spaces. Their results imply that if $2 \leq p < q$ then L_q is not a Lipschitz quotient of L_p . Since L_p is p -convex, it is also Markov p -convex. Hence also all of its subsets are Markov p -convex. But L_q is not p -convex, so we deduce that L_q is not a Lipschitz quotient of any subset of L_p . Thus our new “invariant approach” to the above result of [3] significantly extends it. Note that the method of [3] is based on a differentiation argument, and hence it crucially relies on the fact that the Lipschitz quotient mapping is defined on all of L_p and not just on an arbitrary subset of L_p .

In light of Theorem 1.3 it is natural to ask if Bourgain’s characterization of super-reflexivity holds for general metric spaces. Namely, is it true that for any metric space X , if $\lim_{n \rightarrow \infty} c_X(B_n) = \infty$ then X is Markov p -convex for some $p < \infty$? This question was asked in [16], where it was shown that the answer is positive if (X, d_X) is a metric tree. Here we show that in general the answer is negative:

Theorem 1.5. *There is a metric space (X, d_X) which is not Markov p -convex for any $p \in (0, \infty)$, yet $\lim_{n \rightarrow \infty} c_X(B_n) = \infty$. In fact, (X, d_X) can be a doubling metric space, and hence $c_X(B_n) \geq 2^{\kappa n}$ for some constant $\kappa > 0$.*

Theorem 1.5 is in sharp contrast to the previously established metric characterizations of the linear notions of type and cotype. Specifically, it was shown by Bourgain, Milman and Wolfson [6] that any metric space with no nontrivial metric type must contain the Hamming cubes $(\{0, 1\}^n, \|\cdot\|_1)$ with distortion independent of n . An analogous result was obtained in [24] for metric spaces with no nontrivial metric cotype, with the Hamming cube replaced by the ℓ_∞ grid $(\{1, \dots, m\}^n, \|\cdot\|_\infty)$.

Our proof of Theorem 1.5 is based on an analysis of the behavior of a certain Markov chain on the Laakso graphs, a sequence of combinatorial graphs whose definition is recalled in Section 3. As a consequence of this analysis, we obtain the following distortion lower bound:

Theorem 1.6. *For any $p \geq 2$, the Laakso graph of cardinality n incurs distortion $\Omega((\log n)^{1/p})$ in any embedding into a p -convex Banach space.*

Thus, in particular, for $p > 2$ the n -point Laakso graph incurs distortion $\Omega((\log n)^{1/p})$ in any embedding into L_p . The case of L_p embeddings of the Laakso graphs when $1 < p \leq 2$ was already solved in [26, 12, 15, 14] using the uniform 2-convexity property of L_p . But these proofs rely crucially on 2-convexity and do not extend to the case of p -convexity when $p > 2$. Subsequent to the publication of our proof of Theorem 1.6 in the announcement [23], an alternative proof of this fact was recently discovered by Johnson and Schechtman in [11].

1.1. The nonexistence of a metric dichotomy for trees

Bourgain's metrical characterization of superreflexivity yields the following statement:

Theorem 1.7 (Bourgain's tree dichotomy [5]). *For any Banach space $(X, \|\cdot\|_X)$ one of the following two dichotomic possibilities must hold true: either*

- for all $n \in \mathbb{N}$ we have $c_X(B_n) = 1$, or
- there exists $\alpha = \alpha_X > 0$ such that for all $n \in \mathbb{N}$ we have $c_X(B_n) \geq (\log n)^\alpha$.

Thus, there is a gap in the possible rates of growth of the sequence $\{c_X(B_n)\}_{n=1}^\infty$ when X is a Banach space; consequently, if we were told that, say, $c_X(B_n) = O(\log \log n)$, then we would immediately deduce that actually $c_X(B_n) = 1$ for all n . Additional gap results of this type are known due to the theory of nonlinear type and cotype:

Theorem 1.8 (Bourgain–Milman–Wolfson cube dichotomy [6]). *For any metric space (X, d_X) one of the following two dichotomic possibilities must hold true: either*

- for all $n \in \mathbb{N}$ we have $c_X(\{0, 1\}^n, \|\cdot\|_1) = 1$, or
- there exists $\alpha = \alpha_X > 0$ such that for all $n \in \mathbb{N}$ we have $c_X(\{0, 1\}^n, \|\cdot\|_1) \geq n^\alpha$.

Theorem 1.8 is a metric analogue of Pisier's characterization [28] of Banach spaces with trivial Rademacher type. A metric analogue of the Maurey–Pisier characterization [20] of Banach spaces with finite Rademacher cotype yields the following dichotomy result for ℓ_∞ grids:

Theorem 1.9 (Grid dichotomy [24]). *For any metric space (X, d_X) one of the following two dichotomic possibilities must hold true: either*

- for all $n \in \mathbb{N}$ we have $c_X(\{0, \dots, n\}^n, \|\cdot\|_\infty) = 1$, or
- there exists $\alpha = \alpha_X > 0$ such that for all $n \in \mathbb{N}$ we have $c_X(\{0, \dots, n\}^n, \|\cdot\|_\infty) \geq n^\alpha$.

We refer to the survey article [21] for more information on the theory of metric dichotomies.

Note that Theorem 1.7 is stated for Banach spaces, while Theorems 1.8 and 1.9 hold for general metric spaces. One might expect that as in the case of previous progress on the Ribe program, a metric theory of p -convexity would result in a proof that Theorem 1.7 holds when X is a general metric space. Surprisingly, we show here that this is not true:

Theorem 1.10. *There exists a universal constant $C > 0$ with the following property. Assume that $\{s(n)\}_{n=0}^\infty \subseteq [4, \infty)$ is a nondecreasing sequence such that $\{n/s(n)\}_{n=0}^\infty$ is also nondecreasing. Then there exists a metric space (X, d_X) satisfying, for all $n \geq 2$,*

$$s\left(\left\lfloor \frac{n}{40s(n)} \right\rfloor\right)\left(1 - \frac{Cs(n)\log s(n)}{\log n}\right) \leq c_X(B_n) \leq s(n). \tag{6}$$

Thus, assuming that $s(n) = o\left(\frac{\log n}{\log \log n}\right)$, there exists a subsequence $\{n_k\}_{k=1}^\infty$ for which

$$(1 - o(1))s(n_k) \leq c_X(B_{n_k}) \leq s(n_k). \tag{7}$$

Theorem 1.10 shows that unlike the case of Banach spaces, for general metric spaces, $c_X(B_n)$ can have an arbitrarily slow growth rate.

Bourgain, Milman and Wolfson also obtained in [6] the following finitary version of Theorem 1.8:

Theorem 1.11 (Local rigidity of Hamming cubes [6]). *For every $\varepsilon > 0$, $D > 1$ and $n \in \mathbb{N}$ there exists $m = m(\varepsilon, D, n) \in \mathbb{N}$ such that*

$$\lim_{n \rightarrow \infty} m(\varepsilon, D, n) = \infty,$$

and for every metric d on $\{0, 1\}^n$ which is bi-Lipschitz with distortion $\leq D$ to the ℓ_1 (Hamming) metric,

$$c_{(\{0,1\}^n, d)}(\{0, 1\}^m, \|\cdot\|_1) \leq 1 + \varepsilon.$$

We refer to [6] (see also [30]) for bounds on $m(\varepsilon, D, n)$. Informally, Theorem 1.11 says that the Hamming cube $(\{0, 1\}^n, \|\cdot\|_1)$ is *locally rigid* in the following sense: it is impossible to distort the Hamming metric on a sufficiently large hypercube without the resulting metric space containing a hardly distorted copy of an arbitrarily large Hamming cube. Stated in this way, Theorem 1.11 is a metric version of James’ theorem [9] that ℓ_1 is not a distortable space. The analogue of Theorem 1.11 with the Hamming cube replaced by the ℓ_∞ grid $(\{0, \dots, n\}^n, \|\cdot\|_\infty)$ is Matoušek’s BD-Ramsey theorem [19]; see [24] for quantitative results of this type in the ℓ_∞ case. The following variant of Theorem 1.10 shows that a local rigidity statement as above fails to hold true for binary trees; it can also be viewed as a negative solution of the distortion problem for the infinite binary tree (see [27] and [4, Ch. 13, 14] for more information on the distortion problem for Banach spaces).

Theorem 1.12. *Let B_∞ be the complete unweighted infinite binary tree. For every $D \geq 4$ there exists a metric d on B_∞ that is D -equivalent to the original shortest-path metric on B_∞ , yet for every $\varepsilon \in (0, 1)$ and $m \in \mathbb{N}$,*

$$c_{(B_\infty, d)}(B_m) \leq D - \varepsilon \Rightarrow m \leq D^{CD^2/\varepsilon}.$$

The local rigidity problem for binary trees was studied by several mathematicians. In particular, C. Fefferman asked (private communication, 2005) whether $\{B_n\}_{n=1}^\infty$ have the local rigidity property, and Theorem 1.12 answers this question negatively. Fefferman also

proved a partial local rigidity result which is a nonquantitative variant of Theorem 1.14 below (see also Section 5). We are grateful to C. Fefferman for asking us the question that led to the counterexamples of Theorems 1.10 and 1.12, for sharing with us his partial positive results, and for encouraging us to work on these questions. M. Gromov also investigated the local rigidity problem for binary trees, and proved (via different methods) nonquantitative partial positive results in the spirit of Theorem 1.14. We thank M. Gromov for sharing with us his unpublished work on this topic.

The results of Theorems 1.10 and 1.12 are quite unexpected. Unfortunately, their proofs are delicate and lengthy, and as such constitute the most involved part of this article. In order to facilitate the understanding of these constructions, we end the introduction with an overview of the main geometric ideas that are used in their proofs. This is done in Section 1.1 below—we recommend reading this section first before delving into the technical details presented in Section 6.

Overview of the proofs of Theorems 1.10 and 1.12. For $x \in B_\infty$ let $h(x)$ be its depth, i.e., its distance from the root. Also, for $x, y \in B_\infty$ let $\text{lca}(x, y)$ denote their least common ancestor. The tree metric on B_∞ is then given by

$$d_{B_\infty}(x, y) = h(x) + h(y) - 2h(\text{lca}(x, y)).$$

The metric space X of Theorem 1.10 will be B_∞ as a set, with a new metric defined as follows. Given a sequence $\varepsilon = \{\varepsilon_n\}_{n=0}^\infty \subseteq (0, 1]$ we define $d_\varepsilon : B_\infty \times B_\infty \rightarrow [0, \infty)$ by

$$d_\varepsilon(x, y) = |h(y) - h(x)| + 2\varepsilon_{\min\{h(x), h(y)\}}[\min\{h(x), h(y)\} - h(\text{lca}(x, y))].$$

d_ε does not necessarily satisfy the triangle inequality, but under some simple conditions on the sequence $\{\varepsilon_n\}_{n=0}^\infty$ it does become a metric on B_∞ ; see Lemma 6.1. A pictorial description of the metric d_ε is contained in Figure 1. Note that when $\varepsilon_n = 1$ for all n , we have $d_\varepsilon = d_{B_\infty}$. Below we call the metric spaces $(B_\infty, d_\varepsilon)$ horizontally distorted trees, or *H-trees*, for short.

The metric space (X, d_X) of Theorem 1.10 will be $(B_\infty, d_\varepsilon)$, where $\varepsilon_n = 1/s(n)$ for all n . The identity mapping of B_n into the top n -levels of B_∞ has distortion at most $s(n)$,




Fig. 1. The metric d_ε defined on B_∞ . The arrows indicate horizontal contraction by $\varepsilon_{h(x)}$.

and therefore $c_X(B_n) \leq s(n)$. The challenge is to prove the lower bound on $c_X(B_n)$ in (6). Our initial approach to lower-bounding $c_X(B_n)$ was Matoušek’s metric differentiation proof [18] of asymptotically sharp distortion lower bounds for embeddings of B_n into uniformly convex Banach spaces.

Following Matoušek’s terminology [18], for $\delta > 0$ a quadruple of points (x, y, z, w) in a metric space (X, d_X) is called a δ -fork if $y \in \text{Mid}(x, z, \delta) \cap \text{Mid}(x, w, \delta)$, where for $a, b \in X$ the set $\text{Mid}(a, b, \delta) \subseteq X$ of δ -approximate midpoints is defined as the set of all $w \in X$ satisfying $\max\{d_X(x, y), d_X(y, z)\} \leq \frac{1}{2}(1 + \delta) \cdot d_X(x, z)$. The points z, w will be called below the *prongs* of the δ -fork (x, y, z, w) . Matoušek starts with the observation that if X is a uniformly convex Banach space then in any δ -fork in X the distance between the prongs must be much smaller (as $\delta \rightarrow 0$) than $d_X(x, y)$. Matoušek then shows that for all $D > 0$, any distortion D embedding of B_n into X must map some 0-fork in B_n to a δ -fork in X , provided n is large enough (as a function of D and δ). This reasoning immediately implies that $c_X(B_n)$ must be large when X is a uniformly convex Banach space, and a clever argument of Matoušek in [18] turns this qualitative argument into sharp quantitative bounds.

Of course, we cannot hope to use the above argument of Matoušek in order to prove Theorem 1.10, since Bourgain’s tree dichotomy theorem (Theorem 1.7) does hold true for Banach spaces. But perhaps we can mimic this uniform convexity argument for other target metric spaces? On the face of it, H -trees are ideally suited for this purpose, since the horizontal contractions that we introduced shrink distances between the prongs of canonical forks (call $(x, y, z, w) \in B_\infty$ a *canonical fork* if x is an ancestor of y and z, w are descendants of y at depth $h(x) + 2(h(y) - h(x))$). It is for this reason exactly that we defined H -trees.

Unfortunately, the situation is not that simple. It turns out that H -trees do not behave like uniformly convex Banach spaces in terms of the prong-contractions that they impose on δ -forks. H -trees can even contain larger problematic configurations that have several undistorted δ -forks; such an example is depicted in Figure 2.

Thus, in order to prove Theorem 1.10 it does not suffice to use Matoušek’s argument that a bi-Lipschitz embedding of a large enough B_n must send some 0-fork to a δ -fork. But it turns out that this argument applies not only to forks, but also to larger configurations.

Definition 1.13. Let (T, d_T) be a tree with root r , and let (X, d_X) be a metric space. A mapping $f : T \rightarrow X$ is called a D -vertically faithful embedding if there exists a (scaling factor) $\lambda > 0$ satisfying, for any $x, y \in T$ such that x is an ancestor of y ,

$$\lambda d_T(x, y) \leq d_X(f(x), f(y)) \leq D\lambda d_T(x, y). \tag{8}$$

Recall that the *distortion* of a mapping $\phi : \mathcal{M} \rightarrow \mathcal{N}$ between metric spaces $(\mathcal{M}, d_{\mathcal{M}})$ and $(\mathcal{N}, d_{\mathcal{N}})$ is defined as

$$\text{dist}(\phi) := \left(\sup_{\substack{x, y \in \mathcal{M} \\ x \neq y}} \frac{d_{\mathcal{N}}(\phi(x), \phi(y))}{d_{\mathcal{M}}(x, y)} \right) \cdot \left(\sup_{\substack{x, y \in \mathcal{M} \\ x \neq y}} \frac{d_{\mathcal{M}}(x, y)}{d_{\mathcal{N}}(\phi(x), \phi(y))} \right) \in [1, \infty].$$

With this terminology, we can state the following crucial result.




Fig. 2. The metric space on the right is the H -tree $(B_\infty, d_\varepsilon)$, where $\varepsilon_n = \delta$ for all n . The picture describes an embedding of the tree on the left (B_3 minus four leaves) into $(B_\infty, d_\varepsilon)$ with distortion at most 6, yet all ancestor/descendant distances are distorted by at most $1 + O(\delta)$.

Theorem 1.14. *There exists a universal constant $c > 0$ with the following property. Fix an integer $t \geq 2$, $\delta, \xi \in (0, 1)$, and $D \geq 2$, and assume that $n \in \mathbb{N}$ satisfies*

$$n \geq \frac{1}{\xi} D^{c(t \log t)/\delta}. \quad (9)$$

Let (X, d_X) be a metric space and $f : B_n \rightarrow X$ a D -vertically faithful embedding. Then there exists a mapping $\phi : B_t \rightarrow B_n$ with the following properties.

- *If $x, y \in B_t$ are such that x is an ancestor of y , then $\phi(x)$ is an ancestor of $\phi(y)$.*
- *$\text{dist}(\phi) \leq 1 + \xi$.*
- *The mapping $f \circ \phi : B_t \rightarrow X$ is a $(1 + \delta)$ -vertically faithful embedding of B_t in X .*

Theorem 1.14 is essentially due to Matoušek [18]. Matoušek actually proved this statement only for $t = 2$, since this is all that he needed in order to analyze forks. But his proof extends in a straightforward way to any $t \in \mathbb{N}$. Since we will use this assertion with larger t , for the sake of completeness we reprove it, in a somewhat different way, in Section 5. Note that Theorem 1.14 says that $\{B_n\}_{n=1}^\infty$ do have a local rigidity property with respect to vertically faithfully embeddings.

We solve the problem created by the existence of configurations as those depicted in Figure 2 by studying $(1 + \delta)$ -vertically faithful embeddings of B_4 , and arguing that they must contain a large contracted pair of points. This claim, formalized in Lemma 6.27, is proved in Sections 6.2 and 6.3.

We begin in Section 6.2.1 by studying how the metric P_2 (3-point path) can be approximately embedded in $(B_\infty, d_\varepsilon)$. We find that there are essentially only two ways to embed it in $(B_\infty, d_\varepsilon)$, as depicted in Figure 3. We then proceed in Section 6.2.2 to study δ -forks in $(B_\infty, d_\varepsilon)$. Since forks are formed by “stitching” two approximate P_2 metrics

along a common edge (the handle), we can limit the “search space” using the results of Section 6.2.1. We find that there are six possible types of different approximate forks in $(B_\infty, d_\varepsilon)$, only four of which (depicted in Figure 4) do not have highly contracted prongs. Complete binary trees, and in particular B_4 , are composed of forks stitched together, handle to prong. In order to study handle-to-prong stitching, we investigate in Section 6.2.3 how the metric P_3 (4-point path) can be approximately embedded in $(B_\infty, d_\varepsilon)$. This is again done by studying how two P_2 metrics can be stitched together, this time bottom edge to top edge. We find that there are only three different approximate configurations of P_4 in $(B_\infty, d_\varepsilon)$.

Using the machinery described above, we study in Section 6.3 how the different types of forks can be stitched together in embeddings of B_4 into $(B_\infty, d_\varepsilon)$, reaching the conclusion that a large contraction is unavoidable, and thus completing the proof of Lemma 6.27. The proofs of Theorems 1.10 and 1.12 are concluded in Section 6.4.

2. Markov p -convexity and p -convexity coincide

In this section we prove Theorem 1.3, i.e., that for Banach spaces p -convexity and Markov p -convexity are the same properties. We first show that p -convexity implies Markov p -convexity, and in fact it implies a stronger inequality that is stated in Proposition 2.1 below. The slightly weaker assertion that p -convexity implies Markov p -convexity was first proved in [16], based on an argument from [5]. Our argument here is different and simpler.

It was proved in [29] that a Banach space X is p -convex if and only if it admits an equivalent norm $\|\cdot\|$ for which there exists $K > 0$ such that for every $a, b \in X$,

$$2\|a\|^p + \frac{2}{K^p}\|b\|^p \leq \|a + b\|^p + \|a - b\|^p. \tag{10}$$

Proposition 2.1. *Let $\{X_t\}_{t \in \mathbb{Z}}$ be random variables taking values in a set Ω . For every $s \in \mathbb{Z}$ let $\{\tilde{X}_t(s)\}_{t \in \mathbb{Z}}$ be random variables taking values in Ω , with the following property:*

$$\forall r \leq s \leq t, \quad (X_r, X_t) \text{ and } (X_r, \tilde{X}_t(s)) \text{ have the same distribution.} \tag{11}$$

Fix $p \geq 2$ and let $(X, \|\cdot\|)$ be a Banach space whose norm satisfies (10). Then for every $f : \Omega \rightarrow X$ we have

$$\sum_{k=0}^{\infty} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_t) - f(\tilde{X}_t(t - 2^k))\|^p]}{2^{kp}} \leq (4K)^p \sum_{t \in \mathbb{Z}} \mathbb{E}[\|f(X_t) - f(X_{t-1})\|^p]. \tag{12}$$

Remark 2.2. Observe that condition (11) holds when $\{X_t\}_{t \in \mathbb{Z}}$ is a Markov chain on a state space Ω , and $\{\tilde{X}_t(s)\}_{t \in \mathbb{Z}}$ is as in Definition 1.2.

We start by proving a useful inequality that is a simple consequence of (10).

Lemma 2.3. *Let X be a Banach space whose norm satisfies (10). Then for every $x, y, z, w \in X$,*

$$\frac{\|x - w\|^p + \|x - z\|^p}{2^{p-1}} + \frac{\|z - w\|^p}{4^{p-1}K^p} \leq \|y - w\|^p + \|z - y\|^p + 2\|y - x\|^p. \quad (13)$$

Proof. For every $x, y, z, w \in X$, (10) implies that

$$\frac{\|x - w\|^p}{2^{p-1}} + \frac{2}{K^p} \left\| y - \frac{x + w}{2} \right\|^p \leq \|y - x\|^p + \|y - w\|^p,$$

and

$$\frac{\|z - x\|^p}{2^{p-1}} + \frac{2}{K^p} \left\| y - \frac{z + x}{2} \right\|^p \leq \|z - y\|^p + \|y - x\|^p.$$

Summing these two inequalities, and applying the convexity of the map $u \mapsto \|u\|^p$, we see that

$$\begin{aligned} & \|y - w\|^p + \|z - y\|^p + 2\|y - x\|^p \\ & \geq \frac{\|x - w\|^p + \|z - x\|^p}{2^{p-1}} + \frac{4}{K^p} \cdot \frac{\|y - \frac{x+w}{2}\|^p + \|y - \frac{z+x}{2}\|^p}{2} \\ & \geq \frac{\|x - w\|^p + \|z - x\|^p}{2^{p-1}} + \frac{4}{K^p} \cdot \left\| \frac{z - w}{4} \right\|^p, \end{aligned}$$

implying (13). □

Proof of Proposition 2.1. Using Lemma 2.3 we see that for every $t \in \mathbb{Z}$ and $k \in \mathbb{N}$,

$$\begin{aligned} & \frac{\|f(X_t) - f(X_{t-2^k})\|^p + \|f(\tilde{X}_t(t-2^{k-1})) - f(X_{t-2^k})\|^p}{2^{p-1}} + \frac{\|f(X_t) - f(\tilde{X}_t(t-2^{k-1}))\|^p}{4^{p-1}K^p} \\ & \leq \|f(X_{t-2^{k-1}}) - f(X_t)\|^p + \|f(X_{t-2^{k-1}}) - f(\tilde{X}_t(t-2^{k-1}))\|^p \\ & \quad + 2\|f(X_{t-2^{k-1}}) - f(X_{t-2^k})\|^p. \end{aligned}$$

Taking expectation, and using the assumption (11), we get

$$\begin{aligned} & \frac{\mathbb{E}[\|f(X_t) - f(X_{t-2^k})\|^p]}{2^{p-2}} + \frac{\mathbb{E}[\|f(X_t) - f(\tilde{X}_t(t-2^{k-1}))\|^p]}{4^{p-1}K^p} \\ & \leq 2\mathbb{E}[\|f(X_{t-2^{k-1}}) - f(X_t)\|^p] + 2\mathbb{E}[\|f(X_{t-2^{k-1}}) - f(X_{t-2^k})\|^p]. \end{aligned}$$

Dividing by $2^{(k-1)p+2}$ yields

$$\begin{aligned} & \frac{\mathbb{E}[\|f(X_t) - f(X_{t-2^k})\|^p]}{2^{kp}} + \frac{\mathbb{E}[\|f(X_t) - f(\tilde{X}_t(t-2^{k-1}))\|^p]}{2^{(k+1)p}K^p} \\ & \leq \frac{\mathbb{E}[\|f(X_{t-2^{k-1}}) - f(X_t)\|^p]}{2^{(k-1)p+1}} + \frac{\mathbb{E}[\|f(X_{t-2^{k-1}}) - f(X_{t-2^k})\|^p]}{2^{(k-1)p+1}}. \end{aligned}$$

Summing this inequality over $k = 1, \dots, m$ and $t \in \mathbb{Z}$ we get

$$\begin{aligned} & \sum_{k=1}^m \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_t) - f(X_{t-2^k})\|^p]}{2^{kp}} + \sum_{k=1}^m \sum_{t \in \mathbb{Z}} \frac{[\mathbb{E}\|f(X_t) - f(\tilde{X}_t(t - 2^{k-1}))\|^p]}{2^{(k+1)p} K^p} \\ & \leq \sum_{k=1}^m \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_{t-2^{k-1}}) - f(X_t)\|^p]}{2^{(k-1)p+1}} + \sum_{k=1}^m \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_{t-2^{k-1}}) - f(X_{t-2^k})\|^p]}{2^{(k-1)p+1}} \\ & = \sum_{j=0}^{m-1} \sum_{s \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_s) - f(X_{s-2^j})\|^p]}{2^{jp}}. \end{aligned} \tag{14}$$

It is only of interest to prove (12) when $\sum_{t \in \mathbb{Z}} \mathbb{E}[\|f(X_t) - f(X_{t-1})\|^p] < \infty$. By the triangle inequality, this implies that for every $k \in \mathbb{N}$ we have $\sum_{t \in \mathbb{Z}} \mathbb{E}[\|f(X_t) - f(X_{t-2^k})\|^p] < \infty$. We may therefore cancel terms in (14), arriving at the following inequality:

$$\begin{aligned} & \sum_{k=1}^m \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_t) - f(\tilde{X}_t(t - 2^{k-1}))\|^p]}{2^{(k+1)p} K^p} \\ & \leq \sum_{t \in \mathbb{Z}} \mathbb{E}[\|f(X_t) - f(X_{t-1})\|^p] - \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_t) - f(X_{t-2^m})\|^p]}{2^{mp}} \\ & \leq \sum_{t \in \mathbb{Z}} \mathbb{E}[\|f(X_t) - f(X_{t-1})\|^p]. \end{aligned}$$

Equivalently,

$$\sum_{k=0}^{m-1} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[\|f(X_t) - f(\tilde{X}_t(t - 2^k))\|^p]}{2^{kp}} \leq (4K)^p \sum_{t \in \mathbb{Z}} \mathbb{E}[\|f(X_t) - f(X_{t-1})\|^p].$$

Proposition 2.1 now follows by letting $m \rightarrow \infty$. □

We next prove the more interesting direction of the equivalence of p -convexity and Markov p -convexity: a Markov p -convex Banach space is also p -convex.

Theorem 2.4. *Let $(X, \|\cdot\|)$ be a Banach space which is Markov p -convex with constant Π . Then for every $\varepsilon \in (0, 1)$ there exists a norm $\|\!\| \cdot \!\|$ on X such that for all $x, y \in X$,*

$$(1 - \varepsilon)\|x\| \leq \|\!\|x\!\| \leq \|x\|,$$

and

$$\left\| \frac{x+y}{2} \right\|^p \leq \frac{\|\!\|x\!\|^p + \|\!\|y\!\|^p}{2} - \frac{1 - (1 - \varepsilon)^p}{4\Pi^p(p+1)} \cdot \left\| \frac{x-y}{2} \right\|^p.$$

Thus the norm $\|\!\| \cdot \!\|$ satisfies (10) with constant $K = O(\Pi/\varepsilon^{1/p})$.

Proof. The fact that X is Markov p -convex with constant Π implies that for every Markov chain $\{X_t\}_{t \in \mathbb{Z}}$ with values in X , and for every $m \in \mathbb{N}$, we have

$$\sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|X_t - \tilde{X}_t(t - 2^k)\|^p]}{2^{kp}} \leq \Pi^p \sum_{t=1}^{2^m} \mathbb{E}[\|X_t - X_{t-1}\|^p]. \tag{15}$$

For $x \in X$ we shall say that a Markov chain $\{X_t\}_{t=-\infty}^{2^m}$ is an m -admissible representation of x if $X_t = 0$ for $t \leq 0$ and $\mathbb{E}[X_t] = tx$ for $t \in \{1, \dots, 2^m\}$. Fix $\varepsilon \in (0, 1)$, and denote $\eta = 1 - (1 - \varepsilon)^p$. For every $m \in \mathbb{N}$ define

$$\|x\|_m = \inf \left\{ \left(\frac{1}{2^m} \sum_{t=1}^{2^m} \mathbb{E}[\|X_t - X_{t-1}\|^p] - \frac{\eta}{\Pi^p} \cdot \frac{1}{2^m} \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|X_t - \tilde{X}_t(t - 2^k)\|^p]}{2^{kp}} \right)^{1/p} \right\}, \tag{16}$$

where the infimum is taken over all m -admissible representations of x . Observe that an m -admissible representation of x always exists, since we can define $X_t = 0$ for $t \leq 0$ and $X_t = tx$ for $t \in \{1, \dots, 2^m\}$. This example shows that $\|x\|_m \leq \|x\|$. On the other hand, if $\{X_t\}_{t=-\infty}^{2^m}$ is an m -admissible representation of x then

$$\begin{aligned} & \sum_{t=1}^{2^m} \mathbb{E}[\|X_t - X_{t-1}\|^p] - \frac{\eta}{\Pi^p} \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|X_t - \tilde{X}_t(t - 2^k)\|^p]}{2^{kp}} \\ & \stackrel{(15)}{\geq} (1 - \eta) \sum_{t=1}^{2^m} \mathbb{E}[\|X_t - X_{t-1}\|^p] \geq (1 - \varepsilon)^p \sum_{t=1}^{2^m} \|\mathbb{E}[X_t] - \mathbb{E}[X_{t-1}]\|^p \\ & = (1 - \varepsilon)^p \sum_{t=1}^{2^m} \|tx - (t - 1)x\|^p = 2^m (1 - \varepsilon)^p \|x\|^p, \end{aligned}$$

where in the first inequality we used the convexity of the function $z \mapsto \|z\|^p$. In conclusion, we see that for all $x \in X$,

$$(1 - \varepsilon)\|x\| \leq \|x\|_m \leq \|x\|. \tag{17}$$

Now take $x, y \in X$ and fix $\delta \in (0, 1)$. Let $\{X_t\}_{t=-\infty}^{2^m}$ be an admissible representation of x and $\{Y_t\}_{t=-\infty}^{2^m}$ be an admissible representation of y which is stochastically independent of $\{X_t\}_{t=-\infty}^{2^m}$, such that

$$\sum_{t=1}^{2^m} \mathbb{E}[\|X_t - X_{t-1}\|^p] - \frac{\eta}{\Pi^p} \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|X_t - \tilde{X}_t(t - 2^k)\|^p]}{2^{kp}} \leq 2^m (\|x\|_m^p + \delta), \tag{18}$$

and

$$\sum_{t=1}^{2^m} \mathbb{E}[\|Y_t - Y_{t-1}\|^p] - \frac{\eta}{\Pi^p} \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|Y_t - \tilde{Y}_t(t - 2^k)\|^p]}{2^{kp}} \leq 2^m (\|y\|_m^p + \delta). \tag{19}$$

Define a Markov chain $\{Z_t\}_{t=-\infty}^{2^{m+1}} \subseteq X$ as follows. For $t \leq -2^m$ set $Z_t = 0$. With probability $1/2$ let $(Z_{-2^{m+1}}, Z_{-2^m+2}, \dots, Z_{2^{m+1}})$ equal

$$\underbrace{(0, \dots, 0)}_{2^m \text{ times}}, X_1, \dots, X_{2^m}, X_{2^m} + Y_1, \dots, X_{2^m} + Y_{2^m},$$

and with probability $1/2$ let $(Z_{-2^{m+1}}, Z_{-2^m}, \dots, Z_{2^{m+1}})$ equal

$$\underbrace{(0, \dots, 0)}_{2^m \text{ times}}, Y_1, \dots, Y_{2^m}, X_1 + Y_{2^m}, \dots, X_{2^m} + Y_{2^m}.$$

Hence, $Z_t = 0$ for $t \leq 0$, for $t \in \{1, \dots, 2^m\}$ we have $\mathbb{E}[Z_t] = (\mathbb{E}[X_t] + \mathbb{E}[Y_t])/2 = t(x+y)/2$, and for $t \in \{2^m+1, \dots, 2^{m+1}\}$ we have

$$\begin{aligned} \mathbb{E}[Z_t] &= \frac{\mathbb{E}[X_{2^m} + Y_{t-2^m}] + \mathbb{E}[X_{t-2^m} + Y_{2^m}]}{2} \\ &= \frac{2^m x + (t-2^m)y + (t-2^m)x + 2^m y}{2} = t \frac{x+y}{2}. \end{aligned}$$

Thus $\{Z_t\}_{t=-\infty}^{2^{m+1}}$ is an $(m+1)$ -admissible representation of $(x+y)/2$. The definition (16) implies that

$$2^{m+1} \left\| \frac{x+y}{2} \right\|_{m+1}^p \leq \sum_{t=1}^{2^{m+1}} \mathbb{E}[\|Z_t - Z_{t-1}\|^p] - \frac{\eta}{\Pi^p} \sum_{k=0}^{m+1} \sum_{t=1}^{2^{m+1}} \frac{\mathbb{E}[\|Z_t - \tilde{Z}_t(t-2^k)\|^p]}{2^{kp}}. \quad (20)$$

Note that by definition,

$$\sum_{t=1}^{2^{m+1}} \mathbb{E}[\|Z_t - Z_{t-1}\|^p] = \sum_{t=1}^{2^m} \mathbb{E}[\|X_t - X_{t-1}\|^p] + \sum_{t=1}^{2^m} \mathbb{E}[\|Y_t - Y_{t-1}\|^p]. \quad (21)$$

Moreover,

$$\begin{aligned} & \sum_{k=0}^{m+1} \sum_{t=1}^{2^{m+1}} \frac{\mathbb{E}[\|Z_t - \tilde{Z}_t(t-2^k)\|^p]}{2^{kp}} \\ &= \frac{1}{2^{(m+1)p}} \sum_{t=1}^{2^{m+1}} \mathbb{E}[\|Z_t - \tilde{Z}_t(t-2^{m+1})\|^p] + \sum_{k=0}^m \sum_{t=1}^{2^{m+1}} \frac{\mathbb{E}[\|Z_t - \tilde{Z}_t(t-2^k)\|^p]}{2^{kp}}. \quad (22) \end{aligned}$$

We bound each of the terms in (22) separately. Note that by construction we have for every $t \in \{1, \dots, 2^m\}$,

$$Z_t - \tilde{Z}_t(t-2^{m+1}) = Z_t - \tilde{Z}_t(1-2^{m+1}) = \begin{cases} X_t - Y_t & \text{with probability } 1/4, \\ Y_t - X_t & \text{with probability } 1/4, \\ X_t - \tilde{X}_t(1) & \text{with probability } 1/4, \\ Y_t - \tilde{Y}_t(1) & \text{with probability } 1/4. \end{cases}$$

Thus, the first term on the right hand side of (22) can be bounded from below as follows:

$$\begin{aligned} \frac{1}{2^{(m+1)p}} \sum_{t=1}^{2^{m+1}} \mathbb{E}[\|Z_t - \tilde{Z}_t(t - 2^{m+1})\|^p] &\geq \frac{1}{2^{(m+1)p+1}} \sum_{t=1}^{2^m} \mathbb{E}[\|X_t - Y_t\|^p] \\ &\geq \frac{1}{2^{(m+1)p+1}} \sum_{t=1}^{2^m} \|\mathbb{E}[X_t] - \mathbb{E}[Y_t]\|^p = \frac{\|x - y\|^p}{2^{(m+1)p+1}} \sum_{t=1}^{2^m} t^p \geq \frac{2^m \|x - y\|^p}{2^{p+1}(p+1)}. \end{aligned} \quad (23)$$

We now proceed to bound from below the second term on the right hand side of (22). Note first that for every $k \in \{0, \dots, m\}$ and every $t \in \{2^m + 1, \dots, 2^{m+1}\}$ we have

$$\begin{aligned} Z_t - \tilde{Z}_t(t - 2^k) &= \begin{cases} (X_{2^m} + Y_{t-2^m}) - (\tilde{X}_{2^m}(t - 2^k) + \tilde{Y}_{t-2^m}(t - 2^m - 2^k)) & \text{with probability } 1/2, \\ (Y_{2^m} + X_{t-2^m}) - (\tilde{Y}_{2^m}(t - 2^k) + \tilde{X}_{t-2^m}(t - 2^m - 2^k)) & \text{with probability } 1/2. \end{cases} \end{aligned}$$

By Jensen's inequality, if U, V are X -valued independent random variables with $\mathbb{E}[V] = 0$, then $\mathbb{E}[\|U + V\|^p] \geq \mathbb{E}[\|U + \mathbb{E}[V]\|^p] = \mathbb{E}[\|U\|^p]$. Thus, since $\{X_t\}_{t=-\infty}^{2^m}$ and $\{Y_t\}_{t=-\infty}^{2^m}$ are independent,

$$\begin{aligned} \mathbb{E}[\|Y_{t-2^m} - \tilde{Y}_{t-2^m}(t - 2^m - 2^k) + X_{2^m} - \tilde{X}_{2^m}(t - 2^k)\|^p] &\geq \mathbb{E}[\|Y_{t-2^m} - \tilde{Y}_{t-2^m}(t - 2^m - 2^k)\|^p], \end{aligned}$$

and

$$\begin{aligned} \mathbb{E}[\|X_{t-2^m} - \tilde{X}_{t-2^m}(t - 2^m - 2^k) + Y_{2^m} - \tilde{Y}_{2^m}(t - 2^k)\|^p] &\geq \mathbb{E}[\|X_{t-2^m} - \tilde{X}_{t-2^m}(t - 2^m - 2^k)\|^p]. \end{aligned}$$

It follows that for every $k \in \{0, \dots, m\}$ and every $t \in \{2^m + 1, \dots, 2^{m+1}\}$ we have

$$\begin{aligned} \mathbb{E}[\|Z_t - \tilde{Z}_t(t - 2^k)\|^p] &\geq \frac{1}{2} \mathbb{E}[\|X_{t-2^m} - \tilde{X}_{t-2^m}(t - 2^m - 2^k)\|^p] \\ &\quad + \frac{1}{2} \mathbb{E}[\|Y_{t-2^m} - \tilde{Y}_{t-2^m}(t - 2^m - 2^k)\|^p]. \end{aligned} \quad (24)$$

Hence,

$$\begin{aligned} &\sum_{k=0}^m \sum_{t=1}^{2^{m+1}} \frac{\mathbb{E}[\|Z_t - \tilde{Z}_t(t - 2^k)\|^p]}{2^{kp}} \\ &\stackrel{(24)}{\geq} \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\frac{1}{2} \mathbb{E}[\|X_t - \tilde{X}_t(t - 2^k)\|^p] + \frac{1}{2} \mathbb{E}[\|Y_t - \tilde{Y}_t(t - 2^k)\|^p]}{2^{kp}} \\ &\quad + \sum_{k=0}^m \sum_{t=2^m+1}^{2^{m+1}} \frac{\frac{1}{2} \mathbb{E}[\|X_{t-2^m} - \tilde{X}_{t-2^m}(t - 2^m - 2^k)\|^p] + \frac{1}{2} \mathbb{E}[\|Y_{t-2^m} - \tilde{Y}_{t-2^m}(t - 2^m - 2^k)\|^p]}{2^{kp}} \\ &= \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|X_t - \tilde{X}_t(t - 2^k)\|^p]}{2^{kp}} + \sum_{k=0}^m \sum_{t=1}^{2^m} \frac{\mathbb{E}[\|Y_t - \tilde{Y}_t(t - 2^k)\|^p]}{2^{kp}}. \end{aligned} \quad (25)$$

Combining (18)–(23) and (25), and letting δ tend to 0, we see that

$$2^{m+1} \left\| \frac{x+y}{2} \right\|_{m+1}^p \leq 2^m \|x\|_m^p + 2^m \|y\|_m^p - \frac{\eta}{\Pi^p} \cdot \frac{2^m \|x-y\|^p}{2^{p+1}(p+1)},$$

or

$$\left\| \frac{x+y}{2} \right\|_{m+1}^p \leq \frac{\|x\|_m^p + \|y\|_m^p}{2} - \frac{\eta}{4\Pi^p(p+1)} \cdot \left\| \frac{x-y}{2} \right\|^p. \tag{26}$$

For $w \in X$, define

$$\|w\| = \limsup_{m \rightarrow \infty} \|w\|_m.$$


Then a combination of (17) and (26) yields $(1 - \varepsilon)\|x\| \leq \|x\| \leq \|x\|$, and

$$\begin{aligned} \left\| \frac{x+y}{2} \right\|^p &\leq \frac{\|x\|^p + \|y\|^p}{2} - \frac{\eta}{4\Pi^p(p+1)} \cdot \left\| \frac{x-y}{2} \right\|^p \\ &\leq \frac{\|x\|^p + \|y\|^p}{2} - \frac{\eta}{4\Pi^p(p+1)} \cdot \left\| \frac{x-y}{2} \right\|^p. \end{aligned} \tag{27}$$

Note that (27) implies that the set $\{x \in X : \|x\| \leq 1\}$ is convex, so that $\|\cdot\|$ is a norm on X . This concludes the proof of Theorem 2.4. \square

3. A doubling space which is not Markov p -convex for any $p \in (0, \infty)$

Consider the Laakso graphs [12], $\{G_i\}_{i=0}^\infty$, which are defined as follows. G_0 is the graph consisting of one edge of unit length. To construct G_i , take six copies of G_{i-1} and scale their metric by a factor of $1/4$. We glue four of them cyclically by identifying pairs of endpoints, and attach at two opposite gluing points the remaining two copies. Note that each edge of G_i has length 4^{-i} ; we denote the resulting shortest path metric on G_i by d_{G_i} . As shown in [13, Thm. 2.3], the doubling constant of the metric space (G_i, d_{G_i}) is at most 6.



We direct G_m as follows. Define the root of G_m to be (an arbitrarily chosen) one of the two vertices having only one adjacent edge. In the figure this could be the leftmost vertex r . Note that in no edge the two endpoints are at the same distance from the root. The edges of G_m are then directed from the endpoint closer to the root to the endpoint further away from the root. The resulting directed graph is acyclic. We now define $\{X_t\}_{t=0}^{4^m}$ to be the standard random walk on the directed graph G_m , starting from the root. This random walk is extended to $t \in \mathbb{Z}$ by stipulating that $X_t = X_0$ for $t < 0$, and $X_t = X_{4^m}$ for $t > 4^m$.

Proposition 3.1. *For the random walk defined above,*

$$\sum_{k=0}^{2m} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_{G_m}(X_t, \tilde{X}_t(t-2^k))^p]}{2^{kp}} \gtrsim \frac{m}{8^p} \sum_{t \in \mathbb{Z}} \mathbb{E}[d_{G_m}(X_t, X_{t-1})^p]. \quad (28)$$

Proof. For every $t \in \mathbb{Z}$ we have

$$\mathbb{E}[d_{G_m}(X_t, X_{t-1})^p] = \begin{cases} 4^{-mp}, & t \in \{0, \dots, 4^m - 1\}, \\ 0, & \text{otherwise.} \end{cases}$$

Hence,

$$\sum_{t \in \mathbb{Z}} \mathbb{E}[d_{G_m}(X_t, X_{t-1})^p] = 4^{-m(p-1)}. \quad (29)$$

Fix $k \in \{0, \dots, 2m-2\}$ and write $h = \lceil k/2 \rceil$. View G_m as being built from $A = G_{m-h}$, where each edge of A has been replaced by a copy of G_h . Note that for every $i \in \{0, \dots, 4^{m-h-1} + 1\}$, at time $t = (4i+1)4^h$ the walk X_t is at a vertex of G_m which has two outgoing edges, corresponding to distinct copies of G_h . To see this it suffices to show that all vertices of G_{m-h} that are exactly $4i+1$ edges away from the root, have out-degree 2. This holds since G_{m-h} is obtained from G_{m-h-1} by replacing each edge by a copy of G_1 , and each such copy of G_1 contributes one vertex of out-degree 2, corresponding to the vertex labeled b in the figure describing G_1 .

Consider the set of times

$$T_k := \{0, \dots, 4^m - 1\} \cap \bigcup_{i=0}^{4^{m-h-1}+1} [(4i+1)4^h + 4^{h-2}, (4i+1)4^h + 2 \cdot 4^{h-2}].$$

For $t \in T_k$ find $i \in \{0, \dots, 4^{m-h-1} + 1\}$ such that $t \in [(4i+1)4^h + 4^{h-2}, (4i+1)4^h + 2 \cdot 4^{h-2}]$. Since, by the definition of h , we have $t - 2^k \in [(4i+1)4^h - 4^h, (4i+1)4^h]$, the walks $\{X_s\}_{s \in \mathbb{Z}}$ and $\{\tilde{X}_s(t-2^k)\}_{s \in \mathbb{Z}}$ started evolving independently at some vertex lying in a copy of G_h preceding a vertex v of G_m which has two outgoing edges, corresponding to distinct copies of G_h . Thus, with probability at least $1/2$, the walks X_t and $\tilde{X}_t(t-2^k)$ lie on two distinct copies of G_h in G_m , immediately following the vertex v , and at distance at least $4^{h-2} \cdot 4^{-m}$ and at most $2 \cdot 4^{h-2} \cdot 4^{-m}$ from v . Hence, with probability at least $1/2$ we have $d_{G_m}(X_t, \tilde{X}_t(t-2^k)) \geq 2 \cdot 4^{h-2} \cdot 4^{-m} = 2^{2h-3-2m}$, and therefore

$$\frac{\mathbb{E}[d(X_t, \tilde{X}_t(t-2^k))^p]}{2^{kp}} \geq \frac{\frac{1}{2} 2^{(2h-3-2m)p}}{2^{kp}} \geq 2^{-(2m+3)p-1}.$$

We deduce that for all $k \in \{0, \dots, 2m-2\}$,

$$\begin{aligned} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d(X_t, \tilde{X}_t(t-2^k))^p]}{2^{kp}} &\geq \sum_{t \in T_k} \frac{\mathbb{E}[d(X_t, \tilde{X}_t(t-2^k))^p]}{2^{kp}} \geq |T_k| \cdot 2^{-(2m+3)p-1} \\ &\gtrsim 4^{h-2} \cdot 4^{m-h-1} \cdot 2^{-(2m+3)p-1} \gtrsim \frac{1}{8^p} 4^{-m(p-1)}. \end{aligned} \quad (30)$$

A combination of (29) and (30) implies (28). \square

Proof of Theorem 1.5. As explained in [12, 13], by passing to an appropriate Gromov–Hausdorff limit, there exists a doubling metric space (X, d_X) that contains an isometric copy of all the Laakso graphs $\{G_m\}_{m=0}^\infty$. Proposition 3.1 therefore implies that X is not Markov p -convex for any $p \in (0, \infty)$. \square

Proof of Theorem 1.6. Let (X, d_X) be a Markov p -convex metric space, i.e. $\Pi_p(X) < \infty$. Assume that $f : G_m \rightarrow X$ satisfies

$$x, y \in G_m \Rightarrow \frac{1}{A}d_{G_m}(x, y) \leq d_X(f(x), f(y)) \leq Bd_{G_m}(x, y). \tag{31}$$

Let $\{X_t\}_{t \in \mathbb{Z}}$ be the random walk from Proposition 3.1. Then

$$\begin{aligned} \frac{m}{8^p A^p} \sum_{t \in \mathbb{Z}} \mathbb{E}[d_{G_m}(X_t, X_{t-1})^p] &\stackrel{(28)}{\lesssim} \frac{1}{A^p} \sum_{k=0}^{2m} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_{G_m}(X_t, \tilde{X}_t(t-2^k))^p]}{2^{kp}} \\ &\stackrel{(31)}{\leq} \sum_{k=0}^{2m} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_X(f(X_t), f(\tilde{X}_t(t-2^k)))^p]}{2^{kp}} \\ &\stackrel{(3)}{\leq} \Pi_p(X)^p \sum_{t \in \mathbb{Z}} \mathbb{E}[d_X(f(X_t), f(X_{t-1}))^p] \\ &\stackrel{(31)}{\leq} \Pi_p(X)^p B^p \sum_{t \in \mathbb{Z}} \mathbb{E}[d_{G_m}(X_t, X_{t-1})^p]. \end{aligned}$$

Thus $AB \gtrsim m^{1/p} \gtrsim (\log |G_m|)^{1/p}$. \square

4. Lipschitz quotients

Say that a metric space (Y, d_Y) is a D -Lipschitz quotient of a metric space (X, d_X) if there exist $a, b > 0$ with $ab \leq D$ and a mapping $f : X \rightarrow Y$ such that for all $x \in X$ and $r > 0$,

$$B_Y(f(x), r/a) \subseteq f(B_X(x, r)) \subseteq B_Y(f(x), br). \tag{32}$$

Observe that the last inclusion in (32) is equivalent to f being b -Lipschitz.

The following proposition implies Theorem 1.4.

Proposition 4.1. *If (Y, D_Y) is a D -Lipschitz quotient of (X, d_X) then $\Pi_p(Y) \leq D\Pi_p(X)$.*

Proof. Fix $f : X \rightarrow Y$ satisfying (32). Also, fix a Markov chain $\{X_t\}_{t \in \mathbb{Z}}$ on a state space Ω , and a mapping $g : \Omega \rightarrow Y$.

Fix $m \in \mathbb{Z}$ and let Ω^* be the set of finite sequences of elements of Ω starting at time m , i.e., the set of sequences of the form $(\omega_i)_{i=m}^t \in \Omega^{t-m+1}$ for all $t \geq m$. It will be convenient to consider the Markov chain $\{X_t^*\}_{t=m}^\infty$ on Ω^* given by

$$\Pr[X_t^* = (\omega_m, \omega_{m+1}, \dots, \omega_t)] = \Pr[X_m = \omega_m, X_{m+1} = \omega_{m+1}, \dots, X_t = \omega_t].$$

Also, define $g^* : \Omega^* \rightarrow Y$ by $g^*(\omega_1, \dots, \omega_t) = g(\omega_t)$. By definition, $\{g^*(X_t^*)\}_{t=m}^\infty$ and $\{g(X_t)\}_{t=m}^\infty$ are identically distributed.

We next define a mapping $h^* : \Omega^* \rightarrow X$ such that $f \circ h^* = g^*$ and for all $(\omega_m, \dots, \omega_t) \in \Omega^*$,

$$d_X(h^*(\omega_m, \dots, \omega_{t-1}), h^*(\omega_m, \dots, \omega_t)) \leq ad_Y(g(\omega_{t-1}), g(\omega_t)). \tag{33}$$

For $\omega^* \in \Omega^*$, we will define $h^*(\omega^*)$ by induction on the length of ω^* . If $\omega^* = (\omega_m)$, then we fix $h^*(\omega^*)$ to be an arbitrary element in $f^{-1}(g(\omega_m))$. Assume that $\omega^* = (\omega_m, \dots, \omega_{t-1}, \omega_t)$ and that $h^*(\omega_m, \dots, \omega_{t-1})$ has been defined. Set $x = f(h^*(\omega_m, \dots, \omega_{t-1})) = g^*(\omega_m, \dots, \omega_{t-1}) = g(\omega_{t-1})$ and $r = ad_Y(g(\omega_{t-1}), g(\omega_t))$. Since $g(\omega_t) \in B_Y(x, r/a)$, it follows from (32) that there exists $y \in X$ such that $f(y) = g(\omega_t)$, and $d_X(x, y) \leq r$. We then define $h^*((\omega_m, \dots, \omega_{t-1}, \omega_t)) := y$.

Write $X_t^* = X_m^*$ for $t \leq m$. By the Markov p -convexity of (X, d_X) , we have

$$\sum_{k=0}^{\infty} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_X(h^*(X_t^*), h^*(\tilde{X}_t^*(t - 2^k)))^p]}{2^{kp}} \leq \Pi_p(X)^p \sum_{t \in \mathbb{Z}} \mathbb{E}[d_X(h^*(X_t^*), h^*(X_{t-1}^*))^p]. \tag{34}$$

By (33), for every $t \geq m + 1$ we have

$$d_X(h^*(X_t^*), h^*(X_{t-1}^*)) \leq ad_Y(g(X_t), g(X_{t-1})),$$

while for $t \leq m$ we have $d_X(h^*(X_t^*), h^*(X_{t-1}^*)) = 0$. Thus,

$$\sum_{t \in \mathbb{Z}} \mathbb{E}[d_X(h^*(X_t^*), h^*(X_{t-1}^*))^p] \leq a^p \sum_{t \in \mathbb{Z}} \mathbb{E}[d_Y(g(X_t), g(X_{t-1}))^p]. \tag{35}$$

At the same time, using the fact that f is b -Lipschitz and $f \circ h^* = g^*$, we see that if $t \geq m + 2^k$,

$$\begin{aligned} d_X(h^*(X_t^*), h^*(\tilde{X}_t^*(t - 2^k))) &\geq \frac{1}{b} d_Y(f(h^*(X_t^*)), f(h^*(\tilde{X}_t^*(t - 2^k)))) \\ &= \frac{1}{b} d_Y(g^*(X_t^*), g^*(\tilde{X}_t^*(t - 2^k))) = \frac{1}{b} d_Y(g(X_t), g(\tilde{X}_t(t - 2^k))). \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_X(h^*(X_t^*), h^*(\tilde{X}_t^*(t - 2^k)))^p]}{2^{kp}} &\geq \frac{1}{b^p} \sum_{k=0}^{\infty} \sum_{t=m+2^k}^{\infty} \frac{\mathbb{E}[d_Y(g(X_t), g(\tilde{X}_t(t - 2^k)))^p]}{2^{kp}}. \end{aligned} \tag{36}$$

By combining (35) and (36) with (34), and letting m tend to $-\infty$, we get

$$\sum_{k=0}^{\infty} \sum_{t \in \mathbb{Z}} \frac{\mathbb{E}[d_Y(g(X_t), g(\tilde{X}_t(t - 2^k)))^p]}{2^{kp}} \leq (ab\Pi_p(X))^p \sum_{t \in \mathbb{Z}} \mathbb{E}[d_Y(g(X_t), g(X_{t-1}))^p].$$

Since this inequality holds for every Markov chain $\{X_t\}_{t \in \mathbb{Z}}$ and every $g : \Omega \rightarrow Y$, and since $ab \leq D$, we have proved that $\Pi_p(Y) \leq D\Pi_p(X)$, as required. \square

5. A dichotomy theorem for vertically faithful embeddings of trees

In this section we prove Theorem 1.14. The proof naturally breaks into two parts. The first is the following BD Ramsey property of paths (which can be found nonquantitatively in [19], where also the BD Ramsey terminology is explained).

A mapping $\phi : \mathcal{M} \rightarrow \mathcal{N}$ is called a *rescaled isometry* if $\text{dist}(\phi) = 1$, or equivalently there exists $\lambda > 0$ such that $d_{\mathcal{N}}(\phi(x), \phi(y)) = \lambda d_{\mathcal{M}}(x, y)$ for all $x, y \in \mathcal{M}$. For $n \in \mathbb{N}$ let P_n denote the n -path, i.e., the set $\{0, \dots, n\}$ equipped with the metric inherited from the real line.

Proposition 5.1. *Fix $\delta \in (0, 1)$, $D \geq 2$ and $t, n \in \mathbb{N}$ satisfying $n \geq D^{(4t \log t)/\delta}$. If $f : P_n \rightarrow X$ satisfies $\text{dist}(f) \leq D$ then there exists a rescaled isometry $\phi : P_t \rightarrow P_n$ such that $\text{dist}(f \circ \phi) \leq 1 + \delta$.*

Given a metric space (X, d_X) and a nonconstant mapping $f : P_n \rightarrow X$, define

$$T(X, f) := \frac{d_X(f(0), f(n))}{n \max_{i \in \{1, \dots, n\}} d_X(f(i-1), f(i))} = \frac{d_X(f(0), f(n))}{n \|f\|_{\text{Lip}}}.$$

If f is a constant mapping (equivalently $\max_{i \in \{1, \dots, n\}} d_X(f(i-1), f(i)) = 0$) then we set $T(X, f) = 0$. Note that by the triangle inequality we always have $T(X, f) \leq 1$.

Lemma 5.2. *For every $m, n \in \mathbb{N}$ and $f : P_{mn} \rightarrow X$, there exist rescaled isometries $\phi^{(n)} : P_n \rightarrow P_{mn}$ and $\phi^{(m)} : P_m \rightarrow P_{mn}$ such that*

$$T(X, f) \leq T(X, f \circ \phi^{(m)}) \cdot T(X, f \circ \phi^{(n)}).$$

Proof. Fix $f : P_{mn} \rightarrow X$ and define $\phi^{(m)} : P_m \rightarrow P_{mn}$ by $\phi^{(m)}(i) = in$. Then

$$d_X(f(0), f(mn)) \leq T(X, f \circ \phi^{(m)}) m \max_{i \in \{1, \dots, m\}} d_X(f((i-1)n), f(in)). \quad (37)$$

Similarly, for every $i \in \{1, \dots, m\}$ define $\phi_i^{(n)} : P_n \rightarrow P_{mn}$ by $\phi_i^{(n)}(j) = (i-1)n + j$. Then

$$\begin{aligned} & d_X(f((i-1)n), f(in)) \\ & \leq T(X, f \circ \phi_i^{(n)}) n \max_{j \in \{1, \dots, n\}} d_X(f((i-1)n + j - 1), f((i-1)n + j)). \end{aligned} \quad (38)$$

Letting $i \in \{1, \dots, m\}$ be such that $T(X, f \circ \phi_i^{(n)})$ is maximal, and $\phi^{(n)} = \phi_i^{(n)}$, we conclude that

$$\begin{aligned} & d_X(f(0), f(mn)) \\ & \stackrel{(37) \wedge (38)}{\leq} T(X, f \circ \phi^{(m)}) T(X, f \circ \phi^{(n)}) mn \max_{i \in \{1, \dots, mn\}} d_X(f(i-1), f(i)). \quad \square \end{aligned}$$

Lemma 5.3. *For every $f : P_m \rightarrow X$ we have $\text{dist}(f) \geq 1/T(X, f)$.*

Proof. Assuming $a|i - j| \leq d_X(f(i), f(j)) \leq b|i - j|$ for all $i, j \in P_m$, the claim is $bT(X, f) \geq a$. Indeed,

$$am \leq d_X(f(0), f(m)) \leq T(X, f)m \max_{i \in \{1, \dots, m\}} d_X(f(i-1), f(i)) \leq T(X, f)bm. \quad \square$$

Lemma 5.4. Fix $f : P_m \rightarrow X$. If $0 < \varepsilon < 1/m$ and $T(X, f) \geq 1 - \varepsilon$, then $\text{dist}(f) \leq 1/(1 - m\varepsilon)$.

Proof. Denote $b = \max_{i \in \{1, \dots, m\}} d_X(f(i), f(i-1)) > 0$. For every $0 \leq i < j \leq m$ we have $d_X(f(i), f(j)) \leq \sum_{\ell=i+1}^j d_X(f(\ell-1), f(\ell)) \leq b|j - i|$, and

$$\begin{aligned} (1 - \varepsilon)mb &\leq T(X, f)mb = d_X(f(0), f(m)) \\ &\leq d_X(f(0), f(i)) + d_X(f(i), f(j)) + d_X(f(j), f(m)) \leq d_X(f(i), f(j)) + b(m+i-j). \end{aligned}$$

Thus $d_X(f(i), f(j)) \geq b(j - i - m\varepsilon) \geq (1 - m\varepsilon)b|j - i|$. □

Proof of Proposition 5.1. Set $k = \lfloor \log_t n \rfloor$ and denote by I the identity mapping from P_{t^k} to P_n . By Lemma 5.3 we have $T(X, f \circ I) \geq 1/D$. An iterative application of Lemma 5.2 implies that there exists a rescaled isometry $\phi : P_t \rightarrow P_{t^k}$ such that

$$T(X, f \circ I \circ \phi) \geq D^{-1/k} \geq e^{-(2 \log D)/\log_t n} \geq e^{-\delta/(2t)} \geq 1 - \frac{\delta}{2t}.$$

By Lemma 5.4 we therefore have $\text{dist}(f \circ I \circ \phi) \leq 1/(1 - \delta/2) \leq 1 + \delta$. □

The second part of the proof of Theorem 1.14 uses the following combinatorial lemma due to Matoušek [18]. Denote by $T_{k,m}$ the complete rooted tree of height m , in which every non-leaf vertex has k children. For a rooted tree T , denote by $\text{SP}(T)$ the set of all unordered pairs $\{x, y\}$ of distinct vertices of T such that x is an ancestor of y .

Lemma 5.5 ([18, Lem. 5]). Let $m, r, k \in \mathbb{N}$ satisfy $k \geq r^{(m+1)^2}$. Suppose that each of the pairs from $\text{SP}(T_{k,m})$ is colored by one of r colors. Then there exists a copy T' of B_m in this $T_{k,m}$ such that the color of any pair $\{x, y\} \in \text{SP}(T')$ only depends on the levels of x and y .

Proof of Lemma 1.14. Let $f : B_n \rightarrow X$ be a D -vertically faithful embedding, i.e., for some $\lambda > 0$ it satisfies

$$\lambda d_{B_n}(x, y) \leq d_X(f(x), f(y)) \leq D\lambda d_{B_n}(x, y) \tag{39}$$

whenever $x, y \in B_n$ are such that x is an ancestor of y .

Let $k, \ell \in \mathbb{N}$ be auxiliary parameters to be determined later, and define $m = \lfloor n/(k\ell) \rfloor$. We first construct a mapping $g : T_{2^k, m} \rightarrow B_n$ in a top-down manner as follows. If r is the root of $T_{2^k, m}$ then $g(r)$ is defined to be the root of B_n . Having defined $g(u)$, let $v_1, \dots, v_{2^k} \in T_{2^k, m}$ be the children of u , and let $w_1, \dots, w_{2^k} \in B_n$ be the descendants of $g(u)$ at depth k below $g(u)$. For each $i \in \{1, \dots, 2^k\}$ let $g(v_i)$ be an arbitrary descendant of w_i at depth $h(g(u)) + \ell k$. Note that for this construction to be possible we need to have $m\ell k \leq n$, which is ensured by our choice of m .

By construction, if $x, y \in T_{2^k, m}$ and x is an ancestor of y , then $g(x)$ is an ancestor of $g(y)$ and $d_{B_n}(g(x), g(y)) = \ell k d_{T_{2^k, m}}(x, y)$. Also, if $x, y \in T_{2^k, m}$ and $\text{lca}(x, y) = u$, then we have $h(\text{lca}(g(x), g(y))) \in \{h(g(u)), h(g(u)) + 1, \dots, h(g(u)) + k - 1\}$. This implies that

$$((\ell - 1)k + 1)d_{T_{2^k, m}}(x, y) \leq d_{B_n}(x, y) \leq \ell k d_{T_{2^k, m}}(x, y).$$

Thus, assuming $\ell \geq 2$, we have $\text{dist}(g) \leq 1 + 2/\ell$. Moreover, denoting $F = f \circ g$ and using (39), we see that if $x, y \in T_{2^k, m}$ are such that x is an ancestor of y then

$$k\ell\lambda d_{T_{2^k, m}}(x, y) \leq d_X(F(x), F(y)) \leq D\ell k\lambda d_{T_{2^k, m}}(x, y). \quad (40)$$

Color every pair $\{x, y\} \in \text{SP}(T_{2^k, m})$ with the color

$$\chi(\{x, y\}) := \left\lceil \log_{1+\delta/4} \left(\frac{d_X(F(x), F(y))}{k\ell\lambda d_{T_{2^k, m}}(x, y)} \right) \right\rceil \in \{1, \dots, r\},$$

where $r = \lceil \log_{1+\delta/4} D \rceil$. Assuming that

$$2^k \geq r^{(m+1)^2}, \quad (41)$$

by Lemma 5.5 there exists a copy T' of B_m in $T_{2^k, m}$ such that the colors of pairs $\{x, y\} \in \text{SP}(T')$ only depend on the levels of x and y .

Let P be a root-leaf path in T' (isometric to P_m). The mapping $F|_P : P \rightarrow X$ has distortion at most D by (40). Assuming

$$m \geq D^{16(t \log t)/\delta}, \quad (42)$$

by Proposition 5.1 there are $\{x_i\}_{i=0}^t \subseteq P$ such that for some $a, b \in \mathbb{N}$ with $a, a + tb \in [0, m]$, for all i we have $h(x_i) = a + ib$, and for some $\theta > 0$, for all $i, j \in \{0, \dots, t\}$,

$$\theta b|i - j| \leq d_X(F(x_i), F(x_j)) \leq (1 + \delta/4)\theta b|i - j|. \quad (43)$$

Define a rescaled isometry $\varphi : B_t \rightarrow T'$ in a top-down manner as follows: $\varphi(r) = x_0$, and having defined $\varphi(u) \in T'$, if v, w are the children of u in B_t and v', w' are the children of $\varphi(u)$ in T' , the vertices $\varphi(v), \varphi(w)$ are chosen as arbitrary descendants in T' of v', w' (respectively) at depth $h(\varphi(u)) + b$. Consider the mapping $G : B_t \rightarrow X$ given by $G = F \circ \varphi = f \circ g \circ \varphi$. Take $x, y \in B_t$ such that x is an ancestor of y . Write $h(x) = i$ and $h(y) = j$. Thus $h(\varphi(x)) = a + ib$ and $h(\varphi(y)) = a + jb$. It follows that $\{\varphi(x), \varphi(y)\}$ is colored by the same color as $\{x_i, x_j\}$, i.e.,

$$\begin{aligned} \left\lceil \log_{1+\delta/4} \left(\frac{d_X(G(x), G(y))}{k\ell\lambda b d_{B_t}(x, y)} \right) \right\rceil &= \chi(\{\varphi(x), \varphi(y)\}) = \chi(\{x_i, x_j\}) \\ &= \left\lceil \log_{1+\delta/4} \left(\frac{d_X(F(x_i), F(x_j))}{k\ell\lambda b d_{B_t}(x, y)} \right) \right\rceil. \end{aligned}$$

Consequently, using (43) we deduce that

$$\frac{\theta b}{1 + \delta/4} d_{B_t}(x, y) \leq d_X(G(x), G(y)) \leq (1 + \delta/4)^2 \theta b d_{B_t}(x, y).$$

Thus G is a $(1 + \delta/4)^3 \leq 1 + \delta$ vertically faithful embedding of B_t into X .

It remains to determine the values of the auxiliary parameters ℓ, k , which will lead to the desired restriction on n given in (9). First of all, we want to have $\text{dist}(g \circ \varphi) \leq 1 + \xi$. Since φ is a rescaled isometry and (for $\ell \geq 2$) $\text{dist}(g) \leq 1 + 2/\ell$, we choose $\ell = \lceil 2/\xi \rceil \geq 2$. We will choose k so that $4k \leq n\xi$, so that $n/(k\ell) \geq 1$. Since $m = \lfloor n/(k\ell) \rfloor$, we have $m+1 \leq n\xi/k$ and $m \geq n\xi/(4k)$. Recall that $r = \lceil \log_{1+\delta/4} D \rceil \leq 2 \log_{1+\delta/4} D \leq 16D/\delta$. Hence the requirement (41) will be satisfied if

$$2^{k^3} \geq (16D/\delta)^{n^2\xi^2}, \tag{44}$$

and the requirement (42) will be satisfied if

$$n\xi/(4k) \geq D^{16(r \log t)/\delta}. \tag{45}$$

There exists an integer k satisfying both (44) and (45) provided that

$$(n^2\xi^2 \log_2(16D/\delta))^{1/3} + 1 \leq \frac{n\xi}{4D^{16(r \log t)/\delta}},$$

which holds true provided the constant c in (9) is large enough. □

6. Tree metrics do not have the dichotomy property

This section is devoted to the proofs of Theorems 1.10 and 1.12. These proofs were outlined in Section 1.1, and we will use the notation introduced there.

6.1. Horizontally contracted trees

We start with the following lemma which supplies conditions on $\{\varepsilon_n\}_{n=0}^\infty$ ensuring that the H -tree $(B_\infty, d_\varepsilon)$ is a metric space.

Lemma 6.1. *Assume that $\{\varepsilon_n\}_{n=0}^\infty \subseteq (0, 1]$ is nonincreasing and $\{n\varepsilon_n\}_{n=0}^\infty$ is nondecreasing. Then d_ε is a metric on B_∞ .*

Proof. Take $x, y, z \in B_\infty$ and without loss of generality assume that $h(x) \leq h(y)$. We distinguish between the cases $h(z) > h(y)$, $h(x) \leq h(z) \leq h(y)$ and $h(z) < h(x)$.

If $h(z) > h(y)$ then

$$\begin{aligned} & d_\varepsilon(x, z) + d_\varepsilon(z, y) - d_\varepsilon(x, y) \\ &= 2[h(z) - h(y)] + 2\varepsilon_{h(x)} \cdot [h(\text{lca}(x, y)) - h(\text{lca}(x, z))] + 2\varepsilon_{h(y)} \cdot [h(y) - h(\text{lca}(z, y))] \\ &\geq 2\varepsilon_{h(x)} \cdot [h(\text{lca}(x, y)) - h(\text{lca}(x, z))] + 2\varepsilon_{h(y)} \cdot [h(y) - h(\text{lca}(z, y))]. \end{aligned} \tag{46}$$

To show that (46) is nonnegative observe that this is obvious if $h(\text{lca}(x, y)) \geq h(\text{lca}(x, z))$. So assume that $h(\text{lca}(x, y)) < h(\text{lca}(x, z))$. In this case necessarily $h(\text{lca}(z, y)) = h(\text{lca}(x, y))$, so we can bound (46) from below as follows:

$$\begin{aligned} & 2\varepsilon_{h(x)}[h(\text{lca}(x, y)) - h(\text{lca}(x, z))] + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))] \\ & \geq 2\varepsilon_{h(x)}[h(\text{lca}(x, y)) - h(x)] + 2\frac{h(x)}{h(y)}\varepsilon_{h(x)}[h(y) - h(\text{lca}(x, y))] \\ & = 2\varepsilon_{h(x)}h(\text{lca}(x, y))(1 - h(x)/h(y)) \geq 0. \end{aligned}$$

If $h(z) < h(x)$ then

$$\begin{aligned} d_\varepsilon(x, z) + d_\varepsilon(z, y) &= h(x) + h(y) - 2h(z) + 2\varepsilon_{h(z)}[2h(z) - h(\text{lca}(x, z)) - h(\text{lca}(z, y))] \\ &\geq h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(z)] + 2\varepsilon_{h(x)}[2h(z) - h(\text{lca}(x, z)) - h(\text{lca}(y, z))] \\ &= h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) + h(z) - h(\text{lca}(x, z)) - h(\text{lca}(y, z))] \\ &\geq h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(\text{lca}(x, y))] = d_\varepsilon(x, y), \end{aligned}$$

where we used the fact that $h(z) \geq h(\text{lca}(x, z)) + h(\text{lca}(y, z)) - h(\text{lca}(x, y))$, which is true since $h(\text{lca}(x, y)) \geq \min\{h(\text{lca}(x, z)), h(\text{lca}(y, z))\}$.

It remains to deal with the case $h(x) \leq h(z) \leq h(y)$. In this case

$$\begin{aligned} d_\varepsilon(x, z) + d_\varepsilon(z, y) &= h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(\text{lca}(x, z))] + 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))] \\ &\geq h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(\text{lca}(x, z))] + 2\frac{h(x)}{h(z)}\varepsilon_{h(x)}[h(z) - h(\text{lca}(y, z))] \\ &= h(y) - h(x) + 2\varepsilon_{h(x)}[2h(x) - h(\text{lca}(x, z)) - \frac{h(x)}{h(z)}h(\text{lca}(y, z))] \\ &\geq h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(\text{lca}(x, y))]d_\varepsilon(x, y), \end{aligned}$$

which is equivalent to the inequality

$$h(x) \geq h(\text{lca}(x, z)) + \frac{h(x)}{h(z)}h(\text{lca}(y, z)) - h(\text{lca}(x, y)). \quad (47)$$

To prove (47), note that it is true if $h(\text{lca}(x, y)) \geq h(\text{lca}(x, z))$, since clearly $h(\text{lca}(y, z)) \leq h(z)$. If, on the other hand, $h(\text{lca}(x, y)) < h(\text{lca}(x, z))$ then using the assumption that $h(z) \geq h(x)$ it is enough to show that $h(x) \geq h(\text{lca}(x, z)) + h(\text{lca}(y, z)) - h(\text{lca}(x, y))$. Necessarily $h(\text{lca}(x, y)) = h(\text{lca}(y, z))$, so that the required inequality follows from the fact that $h(x) \geq h(\text{lca}(x, z))$. \square

6.2. Geometry of H -trees

6.2.1. Classification of approximate midpoints. From now on we will always assume that $\varepsilon = \{\varepsilon_n\}_{n=0}^\infty$ satisfies, for all $n \in \mathbb{N}$, $\varepsilon_n \geq \varepsilon_{n+1} > 0$ and $(n+1)\varepsilon_{n+1} \geq n\varepsilon_n$. We recall the important concept of *approximate midpoints* which is used frequently in nonlinear functional analysis (see [4] and the references therein).

Definition 6.2 (Approximate midpoints). Let (X, d_X) be a metric space and $\delta \in (0, 1)$. For $x, z \in X$ the set of δ -approximate midpoints of x and z is defined as

$$\text{Mid}(x, z, \delta) = \left\{ y \in X : \max\{d_X(x, y), d_X(y, z)\} \leq \frac{1 + \delta}{2} d_X(x, z) \right\}.$$

From now on, whenever we refer to the set $\text{Mid}(x, z, \delta)$, the underlying metric will always be understood to be d_ε . In what follows, given $\eta > 0$ we shall say that two sequences (u_1, \dots, u_n) and (v_1, \dots, v_n) of vertices in B_∞ are η -near if for every $j \in \{1, \dots, n\}$ we have $d_\varepsilon(u_j, v_j) \leq \eta$. We shall also require the following terminology:

Definition 6.3. An ordered triple (x, y, z) of vertices in B_∞ will be called a *path-type configuration* if $h(z) \leq h(y) \leq h(x)$, x is a descendant of y , and $h(\text{lca}(z, y)) < h(y)$. The triple (x, y, z) will be called a *tent-type configuration* if $h(y) \leq h(z)$, y is a descendant of x , and $h(\text{lca}(x, z)) < h(x)$. These special configurations are described in Figure 3.




Fig. 3. A schematic description of path-type and tent-type configurations.

The following useful theorem will be used extensively in the ensuing arguments. Its proof will be broken down into several elementary lemmas.

Theorem 6.4. Assume that $\delta \in (0, 1/16)$, and the sequence $\varepsilon = \{\varepsilon_n\}_{n=0}^\infty$ satisfies $\varepsilon_n < 1/4$ for all $n \in \mathbb{N}$. Let $x, y, z \in (B_\infty, d_\varepsilon)$ be such that $y \in \text{Mid}(x, z, \delta)$. Then either (x, y, z) or (z, y, x) is $3\delta d_\varepsilon(x, z)$ -near a path-type or tent-type configuration.

In what follows, given a vertex $v \in B_\infty$ we denote the subtree rooted at v by T_v .

Lemma 6.5. Assume that $\varepsilon_n \leq 1/2$ for all n . Fix $a \in B_\infty$ and let $u, v \in B_\infty$ be its children. For every $x, z \in T_u$ such that $h(x) \geq h(z)$ consider the function $D_{x,z} : \{a\} \cup T_v \rightarrow [0, \infty)$ defined by $D_{x,z}(y) = d_\varepsilon(x, y) + d_\varepsilon(z, y)$. Fix an arbitrary vertex $w \in T_v$ such that $h(w) = h(z)$. Then for every $y \in T_v$ we have $D_{x,z}(y) \geq D_{x,z}(w)$.

Proof. By the definition of d_ε we have $D_{x,z}(y) = Q(h(y))$ where

$$Q(k) = \max\{h(x), k\} + \max\{k, h(z)\} - \min\{h(x), k\} - \min\{k, h(z)\} + 2\varepsilon_{\min\{h(x), k\}}[\min\{h(x), k\} - h(a)] + 2\varepsilon_{\min\{k, h(z)\}}[\min\{k, h(z)\} - h(a)].$$

The required result will follow if we show that Q is nonincreasing on $\{h(a), h(a) + 1, \dots, h(z)\}$ and nondecreasing on $\{h(z), h(z) + 1, \dots\}$. If $k \in \{h(a), h(a) + 1, \dots, h(z) - 1\}$ then

$$\begin{aligned} Q(k + 1) - Q(k) &= -2 + 4\varepsilon_{k+1}[k + 1 - h(a)] - 4\varepsilon_k[k - h(a)] \\ &\leq -2 + 4\varepsilon_k[k + 1 - h(a)] - 4\varepsilon_k[k - h(a)] = -2(1 - 2\varepsilon_k) \leq 0. \end{aligned}$$

If $k \in \{h(x), h(x) + 1, \dots\}$ then $Q(k + 1) - Q(k) = 2$, and if $k \in \{h(z), \dots, h(x) - 1\}$ then

$$Q(k + 1) - Q(k) = 2[(k + 1)\varepsilon_{k+1} - k\varepsilon_k] + 2h(a)[\varepsilon_k - \varepsilon_{k+1}] \geq 0.$$

This completes the proof of Lemma 6.5. □

Lemma 6.6. *Assume that $\varepsilon_n < 1/2$ for all $n \in \mathbb{N}$. Fix $\delta \in (0, 1/3)$ and $x, y, z \in B_\infty$ such that $h(x) \geq h(z)$, $y \in \text{Mid}(x, z, \delta)$ and $h(\text{lca}(x, z)) > h(\text{lca}(x, y))$. Then*


$$h(z) + \frac{1 - 3\delta}{2}d_\varepsilon(x, z) \leq h(y) < h(x) \leq h(y) + \frac{1 + 3\delta}{1 - 3\delta}[h(y) - h(z)].$$

Moreover, if $y' \in B_\infty$ is the point on the segment joining x and $\text{lca}(x, y)$ such that $h(y') = h(y)$ then $d_\varepsilon(y, y') \leq \delta d_\varepsilon(x, z)$. Thus (x, y', z) is a path-type configuration which is $\delta d_\varepsilon(x, z)$ -near (x, y, z)

Proof. Write $a = \text{lca}(x, y)$. If u, v are the two children of a , then without loss of generality $x, z \in T_u$ and $y \in T_v$. Let $w \in T_v$ be such that $h(w) = h(z)$.

By Lemma 6.5,

$$\begin{aligned} d_\varepsilon(x, y) + d_\varepsilon(z, y) &\geq d_\varepsilon(x, w) + d_\varepsilon(z, w) \\ &= h(x) - h(z) + 4\varepsilon_{h(z)}[h(z) - h(a)] \\ &\geq 2d_\varepsilon(x, z) - [h(x) - h(z)]. \end{aligned} \tag{48}$$



On the other hand, since $y \in \text{Mid}(x, z, \delta)$, we have

$$d_\varepsilon(x, y) + d_\varepsilon(z, y) \leq (1 + \delta)d_\varepsilon(x, z).$$

Additionally, by the definition of d_ε we know that if $h(y) \leq h(z)$ then

$$\frac{1 + \delta}{2}d_\varepsilon(x, z) \geq d_\varepsilon(x, y) \geq h(x) - h(y) \geq h(x) - h(z).$$

Combining these observations with (48) we get

$$(1 + \delta)d_\varepsilon(x, z) \geq 2d_\varepsilon(x, z) - \frac{1 + \delta}{2}d_\varepsilon(x, z), \tag{49}$$

which is a contradiction since $\delta < 1/3$. Therefore $h(y) > h(z)$. If $h(y) \geq h(x)$ then

$$\frac{1 + \delta}{2}d_\varepsilon(x, z) \geq d_\varepsilon(z, y) \geq h(y) - h(z) \geq h(x) - h(z),$$

so that we arrive at a contradiction as in (49). We have thus shown that $h(z) < h(y) < h(x)$.

Now, since $y \in \text{Mid}(x, z, \delta)$,

$$\begin{aligned} h(x) - h(z) + 2\varepsilon_{h(y)}[h(y) - h(a)] + 2\varepsilon_{h(z)}[h(z) - h(a)] &= d_\varepsilon(x, y) + d_\varepsilon(z, y) \\ &\leq (1 + \delta)d_\varepsilon(x, z) = (1 + \delta)(h(x) - h(z) + 2\varepsilon_{h(z)}[h(z) - h(a)]). \end{aligned}$$

Thus, letting y' be the point on the segment joining x and a such that $h(y') = h(y)$, we see that

$$d_\varepsilon(y, y') = 2\varepsilon_{h(y)}[h(y) - h(a)] \leq \delta(h(x) - h(z) + 2\varepsilon_{h(z)}[h(z) - h(a)]) = \delta d_\varepsilon(x, z).$$

Moreover

$$\begin{aligned} \frac{1 - \delta}{2}d_\varepsilon(x, z) &\leq d_\varepsilon(y, z) = h(y) - h(z) + 2\varepsilon_{h(z)}h(z) - 2\varepsilon_{h(z)}h(a) \\ &\leq h(y) - h(z) + 2\varepsilon_{h(y)}h(y) - 2\varepsilon_{h(y)}h(a) \leq h(y) - h(z) + \delta d_\varepsilon(x, z). \end{aligned}$$

Thus

$$h(y) - h(z) \geq \frac{1 - 3\delta}{2}d_\varepsilon(x, z). \tag{50}$$

Hence,

$$\frac{2}{1 - 3\delta}[h(y) - h(z)] \stackrel{(50)}{\geq} d_\varepsilon(x, z) = [h(x) - h(y)] + [h(y) - h(z)].$$

It follows that


$$h(x) - h(y) \leq \frac{1 + 3\delta}{1 - 3\delta}[h(y) - h(z)]. \quad \square$$

Lemma 6.7. *Assume that $\varepsilon_n < 1/4$ for all $n \in \mathbb{N}$. Fix $\delta \in (0, 1/16)$ and assume that $x, y, z \in B_\infty$ are distinct vertices such that $\text{lca}(x, y) = \text{lca}(x, z)$, and $y \in \text{Mid}(x, z, \delta)$. Then either (x, y, z) or (z, y, x) is $3\delta d_\varepsilon(x, z)$ -near a path-type or tent-type configuration.*

Proof. Denote $a = \text{lca}(x, y)$. Our assumption implies that $h(\text{lca}(z, y)) \geq h(a)$. We perform a case analysis on the relative heights of x, y, z . Assume first that $h(x) \leq h(y)$.

If $h(x) \leq h(y) \leq h(z)$ then

$$\begin{aligned} (1 + \delta)d_\varepsilon(x, z) &\geq d_\varepsilon(x, y) + d_\varepsilon(y, z) \\ &= h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(a)] \\ &\quad + h(z) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))] \\ &= d_\varepsilon(x, z) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))]. \end{aligned} \tag{51}$$



Let y' be the point on the path from $\text{lca}(y, z)$ to z such that $h(y') = h(y)$. Then (51) implies that

$$d_\varepsilon(y, y') = 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))] \leq \delta d_\varepsilon(x, z).$$

Thus the triple (z, y', x) is a configuration of path-type which is $\delta d_\varepsilon(x, z)$ -near (z, y, x) .

If $h(x) \leq h(z) \leq h(y)$ then since

$$\frac{1+\delta}{2}d_\varepsilon(x, z) \geq d_\varepsilon(x, y) = h(y) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(a)]$$

and $d_\varepsilon(x, z) = h(z) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(a)]$ we deduce that

$$-\frac{1-\delta}{2}d_\varepsilon(x, z) \geq d_\varepsilon(x, y) - d_\varepsilon(x, z) = h(y) - h(z) \geq 0.$$

It follows that $x = z$, contrary to our assumption.

If $h(z) < h(x)$ then let z' be the point on the segment joining a and y such that $h(z') = h(z)$. We thus have

$$d_\varepsilon(z, z') = 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))] = d_\varepsilon(z, y) - [h(y) - h(z)].$$

Moreover,



$$\begin{aligned} & 2\varepsilon_{h(z)}[h(\text{lca}(z, y)) - h(a)] \\ &= d_\varepsilon(x, z) - [h(x) - h(z)] - 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(z, y))] \\ &\geq \frac{2}{1+\delta}d_\varepsilon(z, y) - [h(x) - h(z)] - 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(z, y))] \\ &= \frac{2}{1+\delta}(h(y) - h(z) + 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))]) \\ &\quad - (h(y) - h(z) + 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))]) + [h(y) - h(x)] \\ &= \frac{1-\delta}{1+\delta}d_\varepsilon(y, z) + [h(y) - h(x)] \geq \frac{1-\delta}{1+\delta} \cdot \frac{1-\delta}{2}d_\varepsilon(x, z) + [h(y) - h(x)] \\ &\geq \left(\frac{1-\delta}{1+\delta}\right)^2 d_\varepsilon(x, y) + [h(y) - h(x)] \\ &= d_\varepsilon(x, y) + [h(y) - h(x)] - \frac{4\delta}{(1+\delta)^2}d_\varepsilon(x, y) \\ &= 2[h(y) - h(x)] + 2\varepsilon_{h(x)}[h(x) - h(a)] - \frac{4\delta}{(1+\delta)^2}d_\varepsilon(x, y) \\ &\geq 2[h(y) - h(x)] + 2\varepsilon_{h(z)}h(z) - 2\varepsilon_{h(z)}h(a) - \frac{2\delta}{1+\delta}d_\varepsilon(x, z). \end{aligned}$$

Thus

$$\begin{aligned} \frac{2\delta}{1+\delta}d_\varepsilon(x, z) &\geq 2[h(y) - h(x)] + 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))] \\ &= 2[h(y) - h(x)] + d_\varepsilon(z, z'). \end{aligned} \tag{52}$$


Let y' be the point on the path from a to y such that $h(y') = h(x)$. It follows from (52) that the triple (z', y', x) is a configuration of tent-type which is $2\delta d_\varepsilon(x, z)$ -near (z, y, x) .

This completes the proof of Lemma 6.7 when $h(x) \leq h(y)$. The case $h(x) > h(y)$ is proved analogously. Here are the details.



Assume first of all that $h(z) \geq h(x) > h(y)$. Then

$$\begin{aligned}
 d_\varepsilon(x, z) &\geq \frac{2}{1+\delta} d_\varepsilon(z, y) = 2d_\varepsilon(z, y) - \frac{2\delta}{1+\delta} d_\varepsilon(z, y) \\
 &= 2(h(z) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))]) \\
 &\quad - \frac{2\delta}{1+\delta} d_\varepsilon(z, y) \\
 &\geq 2(h(z) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))]) \\
 &\quad - \delta d_\varepsilon(x, z). \tag{53}
 \end{aligned}$$



On the other hand, since $h(x) > h(y)$,

$$\begin{aligned}
 d_\varepsilon(x, z) &= h(z) - h(x) + 2\varepsilon_{h(x)}[h(x) - h(a)] \\
 &\leq h(z) - h(x) + 2\varepsilon_{h(y)}[h(x) - h(a)] \\
 &= (h(x) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(a)]) + h(y) + h(z) - 2h(x) \\
 &\quad + 2\varepsilon_{h(y)}[h(x) - h(y)] \\
 &= d_\varepsilon(x, y) + h(y) + h(z) - 2h(x) + 2\varepsilon_{h(y)}[h(x) - h(y)] \\
 &\leq \frac{1+\delta}{1-\delta} d_\varepsilon(y, z) + h(y) + h(z) - 2h(x) + 2\varepsilon_{h(y)}[h(x) - h(y)] \\
 &= (h(z) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))]) + \frac{2\delta}{1-\delta} d_\varepsilon(y, z) \\
 &\quad + h(y) + h(z) - 2h(x) + 2\varepsilon_{h(y)}[h(x) - h(y)] \\
 &\leq 2[h(z) - h(x)] + 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(z, y))] + \frac{1+\delta}{1-\delta} \delta d_\varepsilon(x, z).
 \end{aligned}$$

Combining this bound with (53), and canceling terms, gives

$$\begin{aligned}
 \frac{2\delta}{1-\delta} d_\varepsilon(x, z) &\geq 2[h(x) - h(y)] - 4\varepsilon_{h(y)}[h(x) - h(y)] + 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(z, y))] \\
 &\geq 2(1 - 2\varepsilon_{h(y)})[h(x) - h(y)] + 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(z, y))] \\
 &> [h(x) - h(y)] + 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(z, y))], \tag{54}
 \end{aligned}$$


where we used the fact that $\varepsilon_{h(y)} < 1/4$. Let x' be the point on the path from x to a such that $h(x') = h(y)$, and let y' be the point on the path from a to z such that $h(y') = h(y)$. Then by (54) $d_\varepsilon(x, x') = h(x) - h(y) \leq 3\delta d_\varepsilon(x, z)$ and

$$d_\varepsilon(y, y') = 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))] \leq 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(z, y))] \leq 3\delta d_\varepsilon(x, z).$$

Thus the triple (z, y', x') is a configuration of path-type which is $3\delta d_\varepsilon(x, z)$ -near (z, y, x) .

If $h(x) > h(z) \geq h(y)$ then

$$\begin{aligned}
 h(z) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))] \\
 &= d_\varepsilon(z, y) \geq \frac{1-\delta}{1+\delta}d_\varepsilon(x, y) \\
 &= d_\varepsilon(x, y) - \frac{2\delta}{1+\delta}d_\varepsilon(x, y) \\
 &= h(x) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(a)] - \frac{2\delta}{1+\delta}d_\varepsilon(x, y).
 \end{aligned}$$




Canceling terms we see that

$$\begin{aligned}
 \frac{2\delta}{1+\delta}d_\varepsilon(x, y) &\geq h(x) - h(z) + 2\varepsilon_{h(y)}[h(\text{lca}(z, y)) - h(a)] \\
 &= h(x) - h(z) + 2\varepsilon_{h(y)}[h(z) - h(a)] - 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] \\
 &\geq h(x) - h(z) + 2\varepsilon_{h(z)}[h(z) - h(a)] - 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] \\
 &= d_\varepsilon(x, z) - 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] \\
 &\geq 2d_\varepsilon(z, y) - \frac{2\delta}{1+\delta}d_\varepsilon(x, z) - 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] \\
 &= 2(h(z) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(z, y))]) - \frac{2\delta}{1+\delta}d_\varepsilon(x, z) \\
 &\quad - 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] \\
 &= 2(1 - 2\varepsilon_{h(y)})[h(z) - h(y)] + 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] - \frac{2\delta}{1+\delta}d_\varepsilon(x, z) \\
 &\geq [h(z) - h(y)] + 2\varepsilon_{h(y)}[h(z) - h(\text{lca}(z, y))] - \frac{2\delta}{1+\delta}d_\varepsilon(x, z) \\
 &= d_\varepsilon(z, y) - \frac{2\delta}{1+\delta}d_\varepsilon(x, z) \\
 &\geq \left(\frac{1-\delta}{2} - \frac{2\delta}{1+\delta}\right)d_\varepsilon(x, z),
 \end{aligned}$$

which is a contradiction since $\delta < 1/16$.

The only remaining case is when $h(x) > h(y) > h(z)$. In this case we proceed as follows:

$$\begin{aligned}
 d_\varepsilon(x, z) &= h(x) - h(z) + 2\varepsilon_{h(z)}[h(z) - h(a)] \\
 &= d_\varepsilon(y, z) + [h(x) - h(y)] + 2\varepsilon_{h(z)}[h(\text{lca}(y, z)) - h(a)] \\
 &\geq d_\varepsilon(x, y) - \frac{2\delta}{1+\delta}d_\varepsilon(x, y) + [h(x) - h(y)] \\
 &\quad + 2\varepsilon_{h(z)}[h(\text{lca}(y, z)) - h(a)] \\
 &\geq h(x) - h(y) + 2\varepsilon_{h(y)}[h(y) - h(a)] - \delta d_\varepsilon(x, z) \\
 &\quad + [h(x) - h(y)] + 2\varepsilon_{h(z)}[h(\text{lca}(y, z)) - h(a)]
 \end{aligned}$$



$$\begin{aligned}
 &\geq 2[h(x) - h(y)] + 2\varepsilon_{h(z)}[h(z) - h(a)] + 2\varepsilon_{h(z)}[h(\text{lca}(y, z)) - h(a)] - \delta d_\varepsilon(x, z) \\
 &= 2[h(x) - h(y)] + 4\varepsilon_{h(z)}[h(z) - h(a)] - 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))] - \delta d_\varepsilon(x, z) \\
 &= 2d_\varepsilon(x, z) - 2d_\varepsilon(z, y) + 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))] - \delta d_\varepsilon(x, z) \\
 &\geq (1 - 2\delta)d_\varepsilon(x, z) + 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))]. \tag{55}
 \end{aligned}$$

Let z' be the point on the path from a to y such that $h(z') = h(z)$. Then

$$d_\varepsilon(z, z') = 2\varepsilon_{h(z)}[h(z) - h(\text{lca}(y, z))] \stackrel{(55)}{\leq} 2\delta d_\varepsilon(x, z).$$

Therefore the triple (z', y, x) is of tent-type and is $2\delta d_\varepsilon(x, z)$ -near (z, y, x) . \square

Proof of Theorem 6.4. It remains to check that for every $x, y, z \in B_\infty$ such that $y \in \text{Mid}(x, z, \delta)$, at least one of the triples (x, y, z) or (z, y, x) satisfies the conditions of Lemma 6.6 or Lemma 6.7.

Indeed, if $h(\text{lca}(x, y)) = h(\text{lca}(x, z))$ then $\text{lca}(x, y) = \text{lca}(x, z)$, so Lemma 6.7 applies. If $h(\text{lca}(x, y)) < h(\text{lca}(x, z))$ then $\text{lca}(z, y) = \text{lca}(x, z)$, so Lemma 6.7 applies to the triple (z, y, x) . If $h(\text{lca}(x, y)) > h(\text{lca}(x, z))$ then $\text{lca}(x, y) = \text{lca}(z, y)$, and so $h(\text{lca}(x, z)) > h(\text{lca}(z, y))$. Hence Lemma 6.6 applies to either the triple (x, y, z) or the triple (z, y, x) . \square

We end this subsection with a short discussion on the distance between tent-type and path-type configurations. It turns out that when $\varepsilon_h \ll \delta$, a δ -midpoint configuration (x, y, z) can be close to a path-type configuration, and at the same time the reverse triple (z, y, x) close to a tent-type configuration (or vice versa). However, it is easy to see that this is the only ‘‘closeness’’ possible.

Lemma 6.8. *Fix $x, y, z \in B_\infty$ with $x \neq y$. Then the following statements are impossible:*

1. (x, y, z) is $\frac{1}{5}d_\varepsilon(x, y)$ -near a path-type configuration and a tent-type configuration.
2. (x, y, z) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a path-type configuration and (z, y, x) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a path-type configuration.
3. (x, y, z) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a tent-type configuration and (z, y, x) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a tent-type configuration.

Proof. For case 1, assume for contradiction that (x, y, z) is $\frac{1}{5}d_\varepsilon(x, y)$ -near a path-type configuration (a_1, b_1, c_1) , and also $\frac{1}{5}d_\varepsilon(x, y)$ -near a tent-type configuration $(\alpha_1, \beta_1, \gamma_1)$. By the definitions of path-type and tent-type configurations, a_1 is a descendant of b_1 , and β_1 is a descendant of α_1 . Hence,

$$h(a_1) - h(b_1) = d_\varepsilon(a_1, b_1) \geq d_\varepsilon(x, y) - d_\varepsilon(x, a_1) - d_\varepsilon(y, b_1) \geq \frac{3}{5}d_\varepsilon(x, y), \tag{56}$$

$$h(\beta_1) - h(\alpha_1) = d_\varepsilon(\alpha_1, \beta_1) \geq d_\varepsilon(x, y) - d_\varepsilon(x, \alpha_1) - d_\varepsilon(y, \beta_1) \geq \frac{3}{5}d_\varepsilon(x, y). \tag{57}$$

By summing (56) and (57) we see that

$$\begin{aligned}
 \frac{4}{5}d_\varepsilon(x, y) &\geq d_\varepsilon(a_1, x) + d_\varepsilon(x, \alpha_1) + d_\varepsilon(b_1, y) + d_\varepsilon(y, \beta_1) \geq d_\varepsilon(a_1, \alpha_1) + d_\varepsilon(b_1, \beta_1) \\
 &\geq h(a_1) - h(\alpha_1) + h(\beta_1) - h(b_1) \stackrel{(56) \wedge (57)}{\geq} \frac{6}{5}d_\varepsilon(x, y),
 \end{aligned}$$

a contradiction.

For case 2, assume for contradiction that (x, y, z) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a path-type configuration (a_2, b_2, c_2) , and also (z, y, x) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a path-type configuration $(\alpha_2, \beta_2, \gamma_2)$. By the definitions of path-type and tent-type configurations, a_2 is a descendant of b_2 and $h(\beta_2) > h(\gamma_2)$. Hence,

$$h(a_2) - h(b_2) = d_\varepsilon(a_2, b_2) \geq d_\varepsilon(x, y) - d_\varepsilon(x, a_2) - d_\varepsilon(y, b_2) \geq \frac{9}{11}d_\varepsilon(x, y), \quad (58)$$

and

$$\begin{aligned} h(\beta_2) - h(\gamma_2) + 2\varepsilon_{h(\gamma_2)}[h(\gamma_2) - h(\text{lca}(\beta_2, \gamma_2))] &= d_\varepsilon(\beta_2, \gamma_2) \\ &\geq d_\varepsilon(x, y) - d_\varepsilon(x, \gamma_2) - d_\varepsilon(y, \beta_2) \geq \frac{9}{11}d_\varepsilon(x, y). \end{aligned} \quad (59)$$

By summing (58) and (59) we see that

$$\begin{aligned} \frac{17}{11}d_\varepsilon(x, y) &\geq d_\varepsilon(a_2, x) + d_\varepsilon(x, \gamma_2) + d_\varepsilon(b_2, y) + d_\varepsilon(y, \beta_2) + d_\varepsilon(\beta_2, y) \\ &\quad + d_\varepsilon(x, y) + d_\varepsilon(x, \gamma_2) \\ &\geq d_\varepsilon(a_2, \gamma_2) + d_\varepsilon(b_2, \beta_2) + d_\varepsilon(\beta_2, \gamma_2) \\ &\geq (h(a_2) - h(\gamma_2)) + (h(\beta_2) - h(b_2)) + 2\varepsilon_{h(\gamma_2)}[h(\gamma_2) - h(\text{lca}(\beta_2, \gamma_2))] \\ &\stackrel{(58) \wedge (59)}{\geq} \frac{18}{11}d_\varepsilon(x, y), \end{aligned}$$

a contradiction.

For case 3, assume for contradiction that (x, y, z) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a tent-type configuration (a_3, b_3, c_3) , and also (z, y, x) is $\frac{1}{11}d_\varepsilon(x, y)$ -near a tent-type configuration $(\alpha_3, \beta_3, \gamma_3)$. Then b_3 is a descendant of a_3 and $h(\gamma_3) > h(\beta_3)$. Hence,

$$h(b_3) - h(a_3) = d_\varepsilon(a_3, b_3) \geq d_\varepsilon(x, y) - d_\varepsilon(x, a_3) - d_\varepsilon(y, b_3) \geq \frac{9}{11}d_\varepsilon(x, y), \quad (60)$$

and

$$\begin{aligned} h(\gamma_3) - h(\beta_3) + 2\varepsilon_{h(\beta_3)}[h(\beta_3) - h(\text{lca}(\beta_3, \gamma_3))] &= d_\varepsilon(\beta_3, \gamma_3) \\ &\geq d_\varepsilon(x, y) - d_\varepsilon(x, \gamma_3) - d_\varepsilon(y, \beta_3) \geq \frac{9}{11}d_\varepsilon(x, y). \end{aligned} \quad (61)$$

Hence,

$$\begin{aligned} \frac{17}{11}d_\varepsilon(x, y) &\geq d_\varepsilon(b_3, y) + d_\varepsilon(y, \beta_3) + d_\varepsilon(x, \alpha_3) + d_\varepsilon(x, \gamma_3) + d_\varepsilon(\beta_3, y) + d_\varepsilon(x, y) \\ &\quad + d_\varepsilon(x, \gamma_3) \\ &\geq d_\varepsilon(b_3, \beta_3) + d_\varepsilon(a_3, \gamma_3) + d_\varepsilon(\beta_3, \gamma_3) \\ &\geq (h(b_3) - h(\beta_3)) + (h(\gamma_3) - h(a_3)) + 2\varepsilon_{h(\beta_3)}[h(\beta_3) - h(\text{lca}(\beta_3, \gamma_3))] \\ &\stackrel{(60) \wedge (61)}{\geq} \frac{18}{11}d_\varepsilon(x, y), \end{aligned}$$

a contradiction. □

6.2.2. Classification of approximate forks. We begin with three “stitching lemmas” that roughly say that given three points $x, x', y \in (B_\infty, d_\varepsilon)$ such that x' is near x , there exists y' near y such that $d_\varepsilon(x', y')$ is close to $d_\varepsilon(x, y)$, and y' relates to x' in B_∞ “in the same way” that y relates to x in B_∞ .

Lemma 6.9. *Let $x, x', y, y' \in B_\infty$ be such that y is an ancestor of x , and y' is an ancestor of x' satisfying $h(x) - h(y) = h(x') - h(y')$. Then $d_\varepsilon(y, y') \leq d_\varepsilon(x, x')$.*

Proof. Assume without loss of generality that $h(x) \geq h(x')$. So,

$$d_\varepsilon(x, x') = h(x) - h(x') + 2\varepsilon_{h(x')}[h(x') - h(\text{lca}(x, x'))].$$

Note that $h(\text{lca}(y, y')) = \min\{h(y'), h(\text{lca}(x, x'))\}$. Hence,

$$\begin{aligned} d_\varepsilon(y, y') &= h(y) - h(y') + 2\varepsilon_{h(y')}[h(y') - h(\text{lca}(y, y'))] \\ &= h(x) - h(x') + 2\varepsilon_{h(y')}[h(y') - \min\{h(y'), h(\text{lca}(x, x'))\}] \\ &= h(x) - h(x') + 2\varepsilon_{h(y')} \max\{0, h(y') - h(\text{lca}(x, x'))\}. \end{aligned} \quad (62)$$

If the maximum in (62) is 0, then

$$d_\varepsilon(y, y') = h(x) - h(x') \leq d_\varepsilon(x, x').$$

If the maximum in (62) equals $h(y') - h(\text{lca}(x, x'))$, then

$$\begin{aligned} d_\varepsilon(y, y') &= h(x) - h(x') + 2\varepsilon_{h(y')}[h(y') - h(\text{lca}(x, x'))] \\ &\leq h(x) - h(x') + 2\varepsilon_{h(x')}[h(x') - h(\text{lca}(x, x'))] = d_\varepsilon(x, x'), \end{aligned}$$

where we used the fact that the sequence $\{\varepsilon_n(n-a)\}_{n=0}^\infty$ is nondecreasing for all $a \geq 0$. \square

Lemma 6.10. *Let $x, x', y \in B_\infty$ be such that $h(y) \leq h(x)$. Then there exists $y' \in B_\infty$ which satisfies $h(y') - h(x') = h(y) - h(x)$ and*

$$d_\varepsilon(y, y') \leq d_\varepsilon(x, x'), \quad (63)$$

$$d_\varepsilon(x, y) - 2d_\varepsilon(x, x') \leq d_\varepsilon(y', x') \leq d_\varepsilon(x, y) + 2d_\varepsilon(x, x'). \quad (64)$$

Proof. Note that (64) follows from (63) by the triangle inequality. Assume first that $h(x) \geq h(x')$. In this case choose y' to be an ancestor of y satisfying $h(y) - h(y') = h(x) - h(x')$. Then

$$d_\varepsilon(y, y') = h(y) - h(y') = h(x) - h(x') \leq d_\varepsilon(x, x').$$

We next assume that $h(x) < h(x')$. If $h(\text{lca}(x, x')) \neq h(\text{lca}(x, y))$ then choose y' to be an arbitrary descendant of y such that $h(y') - h(y) = h(x') - h(x)$. As before, we conclude that $d_\varepsilon(y, y') = h(y') - h(y) = h(x') - h(x) \leq d_\varepsilon(x, x')$.

It remains to deal with the case $h(x') > h(x)$ and $h(\text{lca}(x, y)) = h(\text{lca}(x', x))$, which also implies that $h(\text{lca}(x', y)) > h(\text{lca}(x, y))$. In this case, we choose y' to be an arbitrary point on a branch containing both $\text{lca}(x, y)$ and x such that $h(y') - h(y) = h(x') - h(x)$. Then $\text{lca}(y', y) = \text{lca}(x, x')$, and therefore

$$\begin{aligned} d_\varepsilon(y, y') &= h(y') - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(y, y'))] \\ &= h(x') - h(x) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(x, x'))] \\ &\leq h(x') - h(x) + 2\varepsilon_{h(x)}[h(x) - h(\text{lca}(x, x'))] = d_\varepsilon(x, x'), \end{aligned}$$

proving (63) in the last remaining case. \square

Lemma 6.11. *Let $x, x', y \in B_\infty$ be such that y is a descendant of x . Then for any $y' \in B_\infty$ which is a descendant of x' and satisfies $h(y') - h(x') = h(y) - h(x)$, we have*

$$\begin{aligned} d_\varepsilon(y, y') &\leq d_\varepsilon(x, x') + 2\varepsilon_{\min\{h(y'), h(y)\}}[h(y) - h(x)] \\ &\leq d_\varepsilon(x, x') + 2\varepsilon_{h(y)}[h(y) - h(x) + d_\varepsilon(x, x')]. \end{aligned}$$

Proof. Note that $h(\text{lca}(y, y')) \geq h(\text{lca}(x, x'))$. Assume first that $h(x') \geq h(x)$. Then

$$\begin{aligned} d_\varepsilon(y, y') &= h(y') - h(y) + 2\varepsilon_{h(y)}[h(y) - h(\text{lca}(y, y'))] \\ &= h(x') - h(x) + 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(y, y'))] + 2\varepsilon_{h(y)}[h(y) - h(x)] \\ &\leq h(x') - h(x) + 2\varepsilon_{h(y)}[h(x) - h(\text{lca}(x, x'))] + 2\varepsilon_{h(y)}[h(y) - h(x)] \\ &\leq d_\varepsilon(x, x') + 2\varepsilon_{h(y)}[h(y) - h(x)]. \end{aligned}$$

When $h(x') < h(x)$, we similarly obtain the bound

$$\begin{aligned} d_\varepsilon(y, y') &= h(y) - h(y') + 2\varepsilon_{h(y')}[h(y') - h(\text{lca}(y, y'))] \\ &= h(x) - h(x') + 2\varepsilon_{h(y')}[h(x') - h(\text{lca}(y, y'))] + 2\varepsilon_{h(y')}[h(y') - h(x')] \\ &\leq d_\varepsilon(x, x') + 2\varepsilon_{h(y')}[h(y) - h(x)]. \end{aligned}$$

The last inequality in the statement of Lemma 6.11 is proved by observing that when $h(y') < h(y)$,

$$\begin{aligned} \varepsilon_{h(y')}[h(y) - h(x)] &= \varepsilon_{h(y')}[h(y') - h(x')] \leq \varepsilon_{h(y)}[h(y) - h(x')] \\ &\leq \varepsilon_{h(y)}[h(y) - h(x) + d_\varepsilon(x, x')]. \quad \square \end{aligned}$$

Definition 6.12. For $\delta \in (0, 1)$ and $x, y, z, w \in B_\infty$, the quadruple (x, y, z, w) is called a δ -fork if

$$y \in \text{Mid}(x, z, \delta) \cap \text{Mid}(x, w, \delta).$$

δ -forks in H-trees can be approximately classified using the approximate classification of midpoint configurations of Section 6.2.1. We have four types of midpoint configurations (recall Figure 3):

- path-type, denoted (P) in what follows,
- reverse path-type, denoted (p); (x, y, z) is of type (p) iff (z, y, x) is of type (P),
- tent-type, denoted (T),
- reverse tent-type, denoted (t); (x, y, z) is of type (t) iff (z, y, x) is of type (T).

Thus, there are $\binom{5}{2} = 10$ possible δ -fork configurations in $(B_\infty, d_\varepsilon)$ (choose two out of the five symbols “P”, “p”, “T”, “t”, “X”, where “X” means “the same”). As we shall see, four of these possible configurations are impossible, two of them have large contraction of the prongs of the forks, i.e., $d_\varepsilon(z, w) \ll d_\varepsilon(x, y)$, which immediately implies large distortion, and the rest of the configurations are problematic in the sense that they are not much distorted from the star $K_{1,3}$ (the metric d on four points p, q, r, s given by $d(p, q) = d(q, r) = d(q, s) = 1$ and $d(p, s) = d(p, r) = d(r, s) = 2$). The ten possible δ -fork configurations are summarized in Table 1.

Table 1. The ten possible fork configurations.

Midpoint configuration	Type
(T T)	Type I
(P P)	Type II
(p T)	Type III
(p t)	Type IV
(p p)	prongs contracted
(t t)	prongs contracted
(P p)	impossible
(P t)	possible only as approximate type II
(P T)	impossible
(t T)	impossible

For future reference, we give names to the four problematic configurations:

Definition 6.13. For $\eta, \delta \in (0, 1)$, a δ -fork (x, y, z, w) of (B_∞, d_ϵ) is called

- η -near Type I (configuration (T||T) in Table 1) if both (x, y, z) and (x, y, w) are η -near tent-type configurations;
- η -near Type II (configuration (P||P) in Table 1) if both (x, y, z) and (x, y, w) are η -near path-type configurations;
- η -near Type III (configuration (p||T) in Table 1) if (z, y, x) is η -near a path-type configuration and (x, y, w) is η -near a tent-type configuration, or vice versa;
- η -near Type IV (configuration (p||t) in Table 1) if (z, y, x) is η -near a path-type configuration and (w, y, x) is η -near a tent-type configuration, or vice versa.

A schematic description of the four problematic configurations is contained in Figure 4.




Fig. 4. The four “problematic” types of δ -forks.

The following lemma is the main result of this section.

Lemma 6.14. Fix $\delta \in (0, 1/70)$ and assume that $\epsilon_n < 1/4$ for all $n \in \mathbb{N}$. If (x, y, z, w) is a δ -fork of (B_∞, d_ϵ) then either it is $35\delta d_\epsilon(x, y)$ -near one of the types I, II, III, IV, or $d_\epsilon(z, w) \leq 2(35\delta + \epsilon_{h_0})d_\epsilon(x, y)$, where $h_0 = \min\{h(x), h(y), h(z), h(w)\}$.

Remark 6.15. One can strengthen the statement of Lemma 6.14 so that in the first case the fork (x, y, z, w) is $O(\delta d_\varepsilon(x, y))$ near another fork (x', y', z', w') which is of (i.e. 0-near) one of the types I–IV. This statement is more complicated to prove, and since we do not actually need it in what follows, we opted to use a weaker property which suffices for our purposes, yet simplifies (the already quite involved) proof.

The proof of Lemma 6.14 proceeds by checking that the cases marked in Table 1 as “impossible” or “prongs contracted” are indeed so—see Figure 5 for a schematic description of the latter case.




Fig. 5. The two configurations of δ -forks with large contraction of the prongs.

We begin with the (p||p) configuration.

Lemma 6.16. *Let (x, y, z, w) be a δ -fork of $(B_\infty, d_\varepsilon)$ and assume that both (z, y, x) and (w, y, x) are $\eta d_\varepsilon(x, y)$ -near path-type configurations. Then, assuming that $\max\{\delta, \eta\} < 1/8$ and $\varepsilon_n < 1/4$ for all n , we have*

$$d_\varepsilon(z, w) \leq (9\eta + 6\delta + 2\varepsilon_{h(y)})d_\varepsilon(x, y).$$

Proof. Let (z', y', x') be a path-type configuration that is $\eta d_\varepsilon(x, y)$ -near (z, y, x) , and let (w'', y'', x'') be a path-type configuration that is $\eta d_\varepsilon(x, y)$ -near (w, y, x) . Without loss of generality assume that $h(y'') \geq h(y')$. Let w' be the descendant of y' satisfying $h(w') - h(y') = h(w'') - h(y'')$ such that w' is either an ancestor or an arbitrary descendant of z' . Note that $h(w') \geq h(y)$. Indeed,

$$\begin{aligned} h(w') &= h(y') + h(w'') - h(y'') = h(y') + d_\varepsilon(w'', y'') \\ &\geq h(y) - |h(y) - h(y')| + d_\varepsilon(w, y) - 2\eta d_\varepsilon(x, y) \\ &\geq h(y) - d_\varepsilon(y, y') + \frac{1 - \delta}{1 + \delta} d_\varepsilon(x, y) - 2\eta d_\varepsilon(x, y) \\ &\geq h(y) + \left(\frac{1 - \delta}{1 + \delta} - 3\eta \right) d_\varepsilon(x, y) \geq h(y). \end{aligned}$$

By Lemma 6.11,

$$\begin{aligned} d_\varepsilon(w', w'') &\leq d_\varepsilon(y', y'') + 2\varepsilon_{h(w')} [h(w'') - h(y'')] \leq d_\varepsilon(y', y'') + 2\varepsilon_{h(w')} d_\varepsilon(y'', w'') \\ &\leq 2\eta d_\varepsilon(x, y) + 2\varepsilon_{h(y)} \left(2\eta + \frac{1 + \delta}{1 - \delta} \right) d_\varepsilon(x, y). \end{aligned} \tag{65}$$

Observe that

$$\begin{aligned} \left(\frac{1-\delta}{1+\delta} - 2\eta\right)d_\varepsilon(x, y) &\leq d_\varepsilon(z, y) - 2\eta d_\varepsilon(x, y) \leq d_\varepsilon(z', y') \\ &\leq d_\varepsilon(z, y) + 2\eta d_\varepsilon(x, y) \leq \left(\frac{1+\delta}{1-\delta} + 2\eta\right)d_\varepsilon(x, y), \end{aligned} \quad (66)$$

Since $d_\varepsilon(w', y') = d_\varepsilon(w'', y'')$, we obtain similarly the bounds:

$$\begin{aligned} \left(\frac{1-\delta}{1+\delta} - 2\eta\right)d_\varepsilon(x, y) &\leq d_\varepsilon(w', y') \leq d_\varepsilon(z, y) + 2\eta d_\varepsilon(x, y) \\ &\leq \left(\frac{1+\delta}{1-\delta} + 2\eta\right)d_\varepsilon(x, y). \end{aligned} \quad (67)$$

Hence

$$d_\varepsilon(z', w') = |d_\varepsilon(y', z') - d_\varepsilon(y', w')| \stackrel{(66)\wedge(67)}{\leq} \left(\frac{4\delta}{1-\delta^2} + 4\eta\right)d_\varepsilon(x, y). \quad (68)$$

So, in conclusion,

$$\begin{aligned} d_\varepsilon(z, w) &\leq d_\varepsilon(z, z') + d_\varepsilon(w, w'') + d_\varepsilon(w'', w') + d_\varepsilon(z', w') \\ &\stackrel{(65)\wedge(68)}{\leq} \left(8\eta + \frac{4\delta}{1-\delta^2} + 2\left(2\eta + \frac{1+\delta}{1-\delta}\right)\varepsilon_{h(y)}\right)d_\varepsilon(x, y) \\ &\leq (9\eta + 6\delta + 2\varepsilon_{h(y)})d_\varepsilon(x, y). \end{aligned} \quad \square$$

We next consider the (t|t) configuration.

Lemma 6.17. *Let (x, y, z, w) be a δ -fork of $(B_\infty, d_\varepsilon)$. Assume that both (z, y, x) and (w, y, x) are $\eta d_\varepsilon(x, y)$ -near tent-type configurations. Then, assuming that $\max\{\delta, \eta\} < 1/4$, we have*

$$d_\varepsilon(z, w) \leq (8\eta + 5\delta)d_\varepsilon(x, y).$$

Proof. Let (z', y', x') be a tent-type configuration that is $\eta d_\varepsilon(x, y)$ -near (z, y, x) , and let (w'', y'', x'') be a tent-type configuration that is $\eta d_\varepsilon(x, y)$ -near (w, y, x) . Assume without loss of generality that $h(y'') - h(w'') \geq h(y') - h(z')$. Let \tilde{w} be a point on the path between w'' and y'' such that $h(y'') - h(\tilde{w}) = h(y') - h(z')$. Then

$$\begin{aligned} d_\varepsilon(w'', \tilde{w}) &= h(y'') - h(w'') - (h(y') - h(z')) = d_\varepsilon(y'', w'') - d_\varepsilon(y', z') \\ &\leq d_\varepsilon(y, w) - d_\varepsilon(y, z) + 4\eta d_\varepsilon(x, y) \\ &\leq \left(\frac{1+\delta}{1-\delta} - \frac{1-\delta}{1+\delta} + 4\eta\right)d_\varepsilon(x, y). \end{aligned} \quad (69)$$

By Lemma 6.9 we have $d_\varepsilon(\tilde{w}, z') \leq d_\varepsilon(y', y'') \leq 2\eta d_\varepsilon(x, y)$. Hence we conclude that

$$d_\varepsilon(y, z) \leq d_\varepsilon(z, z') + d_\varepsilon(\tilde{w}, z') + d_\varepsilon(\tilde{w}, w'') + d_\varepsilon(w'', w) \stackrel{(69)}{\leq} \left(\frac{4\delta}{1-\delta^2} + 8\eta\right)d_\varepsilon(x, y). \quad \square$$

Lemma 6.18. *Let (x, y, z, w) be a δ -fork of B_∞ . Assume that (x, y, z) is $\eta d_\varepsilon(x, y)$ -near a path-type configuration. Assume also that $\delta < 1/30$, $\eta < 1/10$, and $\varepsilon_n < 1/4$ for all n . Then (x, y, w) is $(2\eta + 21\delta)d_\varepsilon(x, y)$ -near a path-type configuration, i.e., (x, y, z, w) is $(2\eta + 21\delta)d_\varepsilon(x, y)$ -near a type II configuration.*

Proof. Let (x', y', z') be a path-type configuration which is $\eta d_\varepsilon(x, y)$ -near (x, y, z) . By Theorem 6.4, either (x, y, w) or (w, y, x) must be $3\delta d_\varepsilon(x, z) \leq \frac{6}{1-\delta}d_\varepsilon(x, y) \leq 7\delta d_\varepsilon(x, y)$ -near either a path-type configuration or a tent-type configuration.

Suppose first that (x, y, w) is $7\delta d_\varepsilon(x, y)$ -near a tent-type configuration (x'', y'', w'') . In this case, x'' is an ancestor of y'' and $h(y'') - h(x'') = d_\varepsilon(x'', y'') \geq (1 - 14\delta)d_\varepsilon(x, y)$. At the same time, y' is an ancestor of x' and $h(x') - h(y') = d_\varepsilon(x', y') \geq (1 - 2\eta)d_\varepsilon(x, y)$. So,

$$\begin{aligned} 2(\eta + 7\delta)d_\varepsilon(x, y) &\geq d_\varepsilon(y'', y') + d_\varepsilon(x', x'') \geq h(y'') - h(x'') + h(x') - h(y') \\ &\geq 2(1 - \eta - 7\delta)d_\varepsilon(x, y), \end{aligned}$$

which is a contradiction since $\eta + 7\delta < 1/2$.

Next suppose that (w, y, x) is $7\delta d_\varepsilon(x, y)$ -near a path-type configuration (w'', y'', x'') . Then $|h(x') - h(x'')| \leq d_\varepsilon(x', x'') \leq (\eta + 7\delta)d_\varepsilon(x, y)$. So,

$$\begin{aligned} (\eta + 7\delta)d_\varepsilon(x, y) &\geq d_\varepsilon(y', y'') \geq h(y'') - h(y') \\ &= (h(y'') - h(x'')) + (h(x'') - h(x')) + h(x') - h(y') \\ &> 0 - (\eta + 7\delta)d_\varepsilon(x, y) + (1 - 2\eta)d_\varepsilon(x, y), \end{aligned}$$

which is a contradiction.

Lastly, suppose that (w, y, x) is $7\delta d_\varepsilon(x, y)$ -near a tent-type configuration (w'', y'', x'') . Note that $|h(y') - h(y'')| \leq d_\varepsilon(y', y'') \leq (\eta + 7\delta)d_\varepsilon(x, y)$. So, $h(y') \geq h(y'') - (\eta + 7\delta)d_\varepsilon(x, y)$. Also,

$$\begin{aligned} h(y'') - h(w'') = d_\varepsilon(y'', w'') &\geq d_\varepsilon(y, w) - 14\delta d_\varepsilon(x, y) \geq \left(\frac{1-\delta}{1+\delta} - 14\delta\right)d_\varepsilon(x, y) \\ &\geq (\eta + 7\delta)d_\varepsilon(x, y). \end{aligned}$$

Consider the point \bar{w} defined as the ancestor of y' at distance $h(y'') - h(w'') - (\eta + 7\delta)d_\varepsilon(x, y)$ from y' . Let also w''' be the ancestor of y'' at distance $h(y'') - h(w'') - (\eta + 7\delta)d_\varepsilon(x, y)$ from y'' . By Lemma 6.9, we have $d_\varepsilon(\bar{w}, w''') \leq d_\varepsilon(y', y'') \leq (\eta + 7\delta)d_\varepsilon(x, y)$. Therefore,

$$d_\varepsilon(\bar{w}, w) \leq d_\varepsilon(\bar{w}, w''') + d_\varepsilon(w''', w'') + d_\varepsilon(w'', w) \leq (2\eta + 21\delta)d_\varepsilon(x, y).$$

Hence (x, y, w) is $(2\eta + 21\delta)d_\varepsilon(x, y)$ -near the path-type configuration (x', y', \bar{w}) . \square

Lemma 6.19. *Let (x, y, z, w) be a δ -fork of B_∞ . Assume that (x, y, z) is $\eta d_\varepsilon(x, y)$ -near a tent-type configuration. Assume also that $\eta < 1/10$ and $\varepsilon_n < 1/4$ for all n . Then (w, y, x) cannot be $\eta d_\varepsilon(x, y)$ -near a tent-type configuration.*

Proof. Let (x', y', z') be a tent type configuration that is $\eta d_\varepsilon(x, y)$ -near (x, y, z) . Suppose for contradiction that there exists a tent type configuration (w'', y'', x'') that is $\eta d_\varepsilon(x, y)$ -near (w, y, x) . Note that $h(y'') \geq h(y') - d_\varepsilon(y', y'') \geq h(y') - 2\eta d_\varepsilon(x, y)$ and $h(y') - h(x') \geq (1 - 2\eta)d_\varepsilon(x, y) > 2\eta d_\varepsilon(x, y)$. Let x^* be the ancestor of y' at distance $h(y') - h(x') - 2\eta d_\varepsilon(x, y)$ from y' , and let \tilde{x} be the ancestor of y'' at distance $h(y') - h(x') - 2\eta d_\varepsilon(x, y)$ from y'' . An application of Lemma 6.9 yields the estimate $d_\varepsilon(\tilde{x}, x^*) \leq d_\varepsilon(y', y'') \leq 2\eta d_\varepsilon(x, y)$. But, since $h(x'') \geq h(y'')$, we also know that $d_\varepsilon(\tilde{x}, x'') \geq h(y'') - h(\tilde{x}) = d_\varepsilon(y', x') - 2\eta d_\varepsilon(x, y)$. Hence,

$$\begin{aligned} 2\eta d_\varepsilon(x, y) &\geq d_\varepsilon(\tilde{x}, x^*) \geq d_\varepsilon(\tilde{x}, x'') - d_\varepsilon(x^*, x') - d_\varepsilon(x', x'') \geq d_\varepsilon(x', y') - 6\eta d_\varepsilon(x, y) \\ &\geq (1 - 8\eta)d_\varepsilon(x, y), \end{aligned}$$

which is a contradiction, since $\eta < 1/10$. □

Proof of Lemma 6.14. Since (x, y, z, w) is a δ -fork, by Theorem 6.4 both (x, y, z) and (x, y, w) are $7\delta d_\varepsilon(x, y)$ -near a tent-type configuration, a path-type configuration, or the corresponding reverse configurations. We have ten possible combinations of these pairs, as appearing in Table 1. By applying Lemmas 6.18 and 6.19 with $\eta = 7\delta$, we rule out three of these configurations, and a fourth configuration is possible but only as $35\delta d_\varepsilon(x, y)$ -near a type II configuration.

We are left with six possible configurations. By applying Lemmas 6.16 and 6.17 with $\eta = 7\delta$ we conclude that in two of those configurations we have $d_\varepsilon(w, z) \leq (69\delta + 2\varepsilon_{h_0})d_\varepsilon(x, y)$, and the rest are configurations that are $7\delta d_\varepsilon(x, y)$ -near one of the types I–IV. □

6.2.3. *Classification of approximate 3-paths.* We start with the following natural notion:

Definition 6.20. For $x_0, x_1, x_2, x_3 \in B_\infty$ the quadruple (x_0, x_1, x_2, x_3) is called a $(1+\delta)$ -approximate P_3 if there exists $L > 0$ such that for every $0 \leq i \leq j \leq 3$ we have

$$(j - i)L \leq d_\varepsilon(x_i, x_j) \leq (1 + \delta)(j - i)L.$$

Note that in this case $x_1 \in \text{Mid}(x_0, x_2, \delta)$ and $x_2 \in \text{Mid}(x_1, x_3, \delta)$.

As in the case of δ -forks, there are ten possible concatenations of two midpoints configurations (path-type or tent-type): P-P, P-p, P-T, P-t, p-P, p-T, p-t, T-T, T-t, t-T (the midpoint configurations p-p, P-p, t-p, T-p, p-P, t-P, T-P, t-t, T-t, t-T are respectively such concatenations with the order of x_0, x_1, x_2, x_3 reversed). We will rule out some of these possibilities, and obtain some stronger properties for the rest. See Table 2.

As in the case of δ -forks, it will be beneficial to give names to three special types of approximate 3-paths:

Definition 6.21. For $x_0, x_1, x_2, x_3 \in B_\infty$ and $\eta > 0$, a quadruple (x_0, x_1, x_2, x_3) is called:


- η -near a type A configuration if both (x_0, x_1, x_2) and (x_1, x_2, x_3) are η -near path-type configurations,

Table 2. The possible configurations of 3-paths.

Midpoint configuration	Reverse configuration	Type
(P-P)	(p-p)	type A
(P-p)	(P-p)	impossible
(P-T)	(t-p)	impossible
(P-t)	(T-p)	type B
(p-P)	(p-P)	impossible
(p-T)	(t-P)	type C
(p-t)	(T-P)	impossible
(T-T)	(t-t)	possible only as type C
(T-t)	(T-t)	impossible
(t-T)	(t-T)	impossible

- η -near a type B configuration if (x_0, x_1, x_2) is η -near a path-type configuration, and (x_3, x_2, x_1) is η -near tent-type configuration,
- η near type C configuration if (x_2, x_1, x_0) is η -near a path-type configuration, and (x_1, x_2, x_3) is η -near a tent-type configuration.

See also Figure 6.

**Fig. 6.** The three possible types of approximate 3-paths.

The following lemma is the main result of this subsection.

Lemma 6.22. Assume that $\varepsilon_n < 1/4$ for all n and fix $\delta < 1/200$. Assume that (x_0, x_1, x_2, x_3) is a $(1+\delta)$ -approximate P_3 . Then either (x_0, x_1, x_2, x_3) or (x_3, x_2, x_1, x_0) is $35\delta d_\varepsilon(x_0, x_1)$ -near a configuration of type A, B or C.

The proof of Lemma 6.22 is again a case analysis that examines all ten possible ways (up to symmetry) to concatenate two midpoint configurations. The proof is divided into a few lemmas according to the cases, and is completed at the end of this subsection.

Lemma 6.23. Assume that $\varepsilon_n < 1/4$ for all n and that (x_0, x_1, x_2, x_3) is a $(1+\delta)$ -approximate P_3 such that (x_0, x_1, x_2) is $\eta d_\varepsilon(x_0, x_1)$ -near a path-type configuration. If $\max\{\delta, \eta\} < 1/200$ then either (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_0, x_1)$ -near a path-type configuration (type A), or (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_0, x_1)$ -near a tent-type configuration (type B).

Proof. Due to Theorem 6.4 we only need to rule out the possibility that (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_1, x_2)$ -near a path-type configuration, or that (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_1, x_2)$ -near a tent-type configuration. Let (x'_0, x'_1, x'_2) be a path-type configuration that is $\eta d_\varepsilon(x_0, x_1)$ -near (x_0, x_1, x_2) .

Suppose first that (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_1, x_2)$ -near the path-type configuration (x''_3, x''_2, x''_1) . Since $h(x'_1) \geq h(x'_2)$ and $h(x''_2) \geq h(x''_1)$ we have

$$\begin{aligned} |h(x_1) - h(x_2)| &\leq |h(x_1) - h(x'_1)| + (h(x'_1) - h(x'_2)) + |h(x'_2) - h(x_2)| \\ &\leq d_\varepsilon(x_1, x'_1) + (h(x'_1) - h(x'_2)) + d_\varepsilon(x'_2, x_2) \\ &\leq 2\eta d_\varepsilon(x_0, x_1) + (h(x'_1) - h(x'_2)), \end{aligned} \quad (70)$$

and similarly,

$$\begin{aligned} |h(x_1) - h(x_2)| &\leq |h(x_1) - h(x''_1)| + (h(x''_2) - h(x''_1)) + |h(x''_2) - h(x_2)| \\ &\leq d_\varepsilon(x_1, x''_1) + (h(x''_2) - h(x''_1)) + d_\varepsilon(x'_2, x_2) \\ &\leq 14\delta d_\varepsilon(x_0, x_1) + (h(x''_2) - h(x''_1)). \end{aligned} \quad (71)$$

By summing (70) and (71) we obtain the bound

$$2|h(x_1) - h(x_2)| \leq (2\eta + 14\delta)d_\varepsilon(x_0, x_1) + d_\varepsilon(x'_1, x''_1) + d_\varepsilon(x'_2, x''_2) \leq (4\eta + 28\delta)d_\varepsilon(x_0, x_1).$$

Thus

$$|h(x_1) - h(x_2)| \leq (2\eta + 14\delta)d_\varepsilon(x_0, x_1). \quad (72)$$

Since x'_0 is a descendant of x'_1 ,

$$\begin{aligned} |h(x_0) - h(x_1) - d_\varepsilon(x_0, x_1)| &\leq |h(x'_0) - h(x'_1) - d_\varepsilon(x_0, x_1)| + 2\eta d_\varepsilon(x_0, x_1) \\ &= |d_\varepsilon(x'_0, x'_1) - d_\varepsilon(x_0, x_1)| + 2\delta d_\varepsilon(x_0, x_1) \leq 4\eta d_\varepsilon(x_0, x_1). \end{aligned} \quad (73)$$

Similarly, since x''_3 is a descendant of x''_2 ,

$$\begin{aligned} |h(x_3) - h(x_2) - d_\varepsilon(x_0, x_1)| &\leq |h(x''_3) - h(x''_2) - d_\varepsilon(x_0, x_1)| + 14\delta d_\varepsilon(x_1, x_2) \\ &= |d_\varepsilon(x''_3, x''_2) - d_\varepsilon(x_0, x_1)| + 14\delta d_\varepsilon(x_0, x_1) \leq 28\delta d_\varepsilon(x_0, x_1). \end{aligned} \quad (74)$$

Hence,

$$\begin{aligned} |h(x''_3) - h(x'_0)| &\leq |h(x''_3) - h(x_3)| + |h(x_3) - h(x_2) - d_\varepsilon(x_0, x_1)| + |h(x_2) - h(x_1)| \\ &\quad + |h(x_0) - h(x_1) - d_\varepsilon(x_0, x_1)| + |h(x_0) - h(x'_0)| \\ &\stackrel{(72) \wedge (73) \wedge (74)}{\leq} d_\varepsilon(x''_3, x_3) + 28\delta d_\varepsilon(x_0, x_1) + (2\eta + 14\delta)d_\varepsilon(x_0, x_1) + 4\eta d_\varepsilon(x_0, x_1) \\ &\quad + d_\varepsilon(x_0, x'_0) \\ &\leq (49\delta + 7\eta)d_\varepsilon(x_0, x_1). \end{aligned} \quad (75)$$

We record for future reference the following consequence of (72) and (75):

$$\begin{aligned}
\min\{h(x'_0), h(x'_3)\} - \min\{h(x'_1), h(x'_2)\} &\leq \max\{h(x'_0) - h(x'_1), h(x'_0) - h(x'_2)\} \\
&\stackrel{(75)}{\leq} \max\{d_\varepsilon(x'_0, x'_1), h(x'_3) - h(x'_2) + (49\delta + 7\eta)d_\varepsilon(x_0, x_1)\} \\
&\leq \max\{(1 + 2\eta)d_\varepsilon(x_0, x_1), d_\varepsilon(x'_3, x'_2) + (49\delta + 7\eta)d_\varepsilon(x_0, x_1)\} \\
&\leq (1 + 64\delta + 7\eta)d_\varepsilon(x_0, x_1). \tag{76}
\end{aligned}$$

We next claim that

$$\text{lca}(x'_0, x'_3) = \text{lca}(x'_1, x'_2). \tag{77}$$

Indeed, since x'_1 is an ancestor of x'_0 and x'_2 is an ancestor of x'_3 , if $\text{lca}(x'_0, x'_3) \neq \text{lca}(x'_1, x'_2)$ then either x'_1 is a descendant of x'_2 , or x'_2 is a descendant of x'_1 . If x'_1 is a descendant of x'_2 then

$$\begin{aligned}
(\eta + 7\delta)d_\varepsilon(x_0, x_1) &\geq d_\varepsilon(x'_1, x'_1) \geq d_\varepsilon(x'_2, x'_1) \geq d_\varepsilon(x_2, x_1) - (\eta + 7\delta)d_\varepsilon(x_0, x_1) \\
&\geq \frac{1}{1 + \delta}d_\varepsilon(x_0, x_1) - (\eta + 7\delta)d_\varepsilon(x_0, x_1),
\end{aligned}$$

which is a contradiction since $\delta, \eta < 1/200$. Similarly, if x'_2 is a descendant of x'_1 then

$$(\eta + 7\delta)d_\varepsilon(x_0, x_1) \geq d_\varepsilon(x'_2, x'_2) \geq d_\varepsilon(x'_2, x'_1) \geq \frac{1}{1 + \delta}d_\varepsilon(x_0, x_1) - (\eta + 7\delta)d_\varepsilon(x_0, x_1),$$

leading once more a contradiction. This proves (77).

Now,

$$\begin{aligned}
\frac{3}{1 + \delta}d_\varepsilon(x_0, x_1) &\leq d_\varepsilon(x_0, x_3) \\
&\leq d_\varepsilon(x'_3, x'_0) + (\eta + 7\delta)d_\varepsilon(x_0, x_1) \\
&\stackrel{(75)}{\leq} 2\varepsilon_{\min\{h(x'_0), h(x'_3)\}}[\min\{h(x'_0), h(x'_3)\} - h(\text{lca}(x'_0, x'_3))] + (8\eta + 56\delta)d_\varepsilon(x_0, x_1) \\
&\stackrel{(77)}{=} 2\varepsilon_{\min\{h(x'_0), h(x'_3)\}}[\min\{h(x'_1), h(x'_2)\} - h(\text{lca}(x'_1, x'_2))] + (8\eta + 56\delta)d_\varepsilon(x_0, x_1) \\
&\quad + 2\varepsilon_{\min\{h(x'_0), h(x'_3)\}}[\min\{h(x'_0), h(x'_3)\} - \min\{h(x'_1), h(x'_2)\}] \\
&\stackrel{(76)}{\leq} 2\varepsilon_{\min\{h(x'_1), h(x'_2)\}}[\min\{h(x'_1), h(x'_2)\} - h(\text{lca}(x'_1, x'_2))] \\
&\quad + \left(8\eta + 56\delta + \frac{1 + 64\delta + 7\eta}{2}\right)d_\varepsilon(x_0, x_1) \tag{78}
\end{aligned}$$

$$\begin{aligned}
&\leq d_\varepsilon(x'_1, x'_2) + (1/2 + 88\delta + 12\eta)d_\varepsilon(x_0, x_1) \\
&\leq (3/2 + 96\delta + 13\eta)d_\varepsilon(x_0, x_1), \tag{79}
\end{aligned}$$

where in (78) we used $\min\{h(x'_0), h(x'_3)\} \geq \min\{h(x'_1), h(x'_2)\}$ and $\varepsilon_{\min\{h(x'_0), h(x'_3)\}} < 1/4$. Since $\max\{\eta, \delta\} < 1/200$, the bound (79) is a contradiction.

Next suppose (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_0, x_1)$ -near a tent-type configuration (x_1'', x_2'', x_3'') . Since $h(x_2') \leq h(x_1')$ and $h(x_2'') \geq h(x_1'')$, we have

$$\begin{aligned} |h(x_1) - h(x_2)| &\leq |h(x_1) - h(x_1')| + (h(x_1') - h(x_2')) + |h(x_2') - h(x_2)| \\ &\leq d_\varepsilon(x_1, x_1') + (h(x_1') - h(x_2')) + (h(x_2'') - h(x_1'')) + d_\varepsilon(x_2', x_2) \\ &\leq d_\varepsilon(x_1, x_1') + d_\varepsilon(x_1', x_1'') + d_\varepsilon(x_2', x_2'') + d_\varepsilon(x_2', x_2) \\ &\leq (4\eta + 14\delta)d_\varepsilon(x_0, x_1). \end{aligned} \quad (80)$$

On the other hand, x_1'' is an ancestor of x_2'' , and therefore

$$\begin{aligned} \left(\frac{1}{1+\delta} - 14\delta\right)d_\varepsilon(x_0, x_1) &\leq d_\varepsilon(x_1'', x_2'') = h(x_2'') - h(x_1'') \\ &\leq |h(x_1) - h(x_2)| + 14\delta d_\varepsilon(x_0, x_1) \stackrel{(80)}{\leq} (4\eta + 28\delta)d_\varepsilon(x_0, x_1), \end{aligned} \quad (81)$$

which is a contradiction since $\max\{\eta, \delta\} < 1/200$. \square

Lemma 6.24. *Assume that $\varepsilon_n < 1/4$ for all n and that (x_0, x_1, x_2, x_3) is a $(1 + \delta)$ -approximate P_3 such that (x_2, x_1, x_0) is $\eta d_\varepsilon(x_0, x_1)$ -near a path-type configuration. If $\max\{\delta, \eta\} < 1/200$ then either (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_0, x_1)$ -near a path-type configuration (reverse type A), or (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_0, x_1)$ -near a tent-type configuration (type C).*

Proof. Let (x_2', x_1', x_0') be a path-type configuration that is $\eta d_\varepsilon(x_0, x_1)$ -near (x_2, x_1, x_0) . First, assume for contradiction that (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_1, x_2)$ -near a tent-type configuration (x_3'', x_2'', x_1'') . Then $h(x_1'') \geq h(x_2'')$, whereas $h(x_2') - h(x_1') = d_\varepsilon(x_2', x_1')$. Arguing as in (80), it follows that $|h(x_1) - h(x_2)| \leq (2\eta + 28\delta)d_\varepsilon(x_0, x_1)$, and we arrive at a contradiction by arguing similarly to (81).

Next, assume for contradiction that (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_1, x_2)$ -near a path-type configuration (x_1'', x_2'', x_3'') . Then $h(x_1'') - h(x_2'') = d_\varepsilon(x_1'', x_2'')$, whereas $h(x_2') - h(x_1') = d_\varepsilon(x_2', x_1')$. By summing these two identities, we arrive at a contradiction as follows:

$$\begin{aligned} \left(\frac{2}{1+\delta} - 2\eta - 14\delta\right)d_\varepsilon(x_0, x_1) &\leq d_\varepsilon(x_1', x_2') + d_\varepsilon(x_1'', x_2'') \\ &= (h(x_2') - h(x_2'')) + (h(x_1'') - h(x_1')) \leq d_\varepsilon(x_2', x_2'') + d_\varepsilon(x_1', x_1'') \\ &\leq (2\eta + 14\delta)d_\varepsilon(x_0, x_1). \end{aligned} \quad \square$$

Lemma 6.25. *Assume that $\varepsilon_n < 1/4$ for all n and that (x_0, x_1, x_2, x_3) is a $(1 + \delta)$ -approximate P_3 such that (x_0, x_1, x_2) is $\eta d_\varepsilon(x_0, x_1)$ -near a tent-type configuration. If $\max\{\delta, \eta\} < 1/200$ then either (x_2, x_1, x_0) is $(14\delta + 3\eta)d_\varepsilon(x_0, x_1)$ -near a path-type configuration and (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_1, x_2)$ -near a tent-type configuration (type C), or (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_0, x_1)$ -near a path-type configuration (reverse type B).*

Proof. Let (x_0', x_1', x_2') be a tent-type configuration that is $\eta d_\varepsilon(x_0, x_1)$ -near (x_0, x_1, x_2) . First, suppose that (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_1, x_2)$ -near a tent-type configuration (x_1'', x_2'', x_3'') . Note that $|h(x_1') - h(x_1'')| \leq d_\varepsilon(x_1', x_1'') \leq (\eta + 7\delta)d_\varepsilon(x_0, x_1)$. So, let x_0'' be an ancestor of x_1'' at distance $h(x_1') - h(x_0'') - (\eta + 7\delta)d_\varepsilon(x_0, x_1) \in [0, h(x_1'')]$ from x_1'' , and let

x_0^* be an ancestor of x_1' at distance $h(x_1') - h(x_0') - (\eta + 7\delta)d_\varepsilon(x_0, x_1)$ from x_1' . Then $h(x_1') - h(x_0^*) = h(x_1'') - h(x_0'')$ and $d_\varepsilon(x_0^*, x_0') \leq (\eta + 7\delta)d_\varepsilon(x_0, x_1)$. By Lemma 6.9,

$$\begin{aligned} d_\varepsilon(x_0, x_0'') - (2\eta + 7\delta)d_\varepsilon(x_0, x_1) &\leq d_\varepsilon(x_0, x_0'') - d_\varepsilon(x_0^*, x_0') - d_\varepsilon(x_0', x_0) \leq d_\varepsilon(x_0^*, x_0'') \\ &\leq d_\varepsilon(x_1', x_1'') \leq (\eta + 7\delta)d_\varepsilon(x_0, x_1). \end{aligned}$$

Hence (x_2'', x_1'', x_0'') is a path-type configuration that is $(14\delta + 3\eta)d_\varepsilon(x_0, x_1)$ -near (x_2, x_1, x_0) .

Next assume for contradiction that (x_3, x_2, x_1) is $7\delta d_\varepsilon(x_1, x_2)$ near a tent-type configuration (x_3', x_2', x_1') . Then

$$\begin{aligned} (1 - 15\delta)d_\varepsilon(x_0, x_1) &\leq \left(\frac{1}{1 + \delta} - 14\delta\right)d_\varepsilon(x_0, x_1) \leq h(x_2'') - h(x_3'') \\ &\leq (1 + 15\delta)d_\varepsilon(x_0, x_1), \end{aligned} \quad (82)$$

and

$$\begin{aligned} (1 - \delta - 2\eta)d_\varepsilon(x_0, x_1) &\leq \left(\frac{1}{1 + \delta} - 2\eta\right)d_\varepsilon(x_0, x_1) \leq h(x_1') - h(x_0') \\ &\leq (1 + \delta + 2\eta)d_\varepsilon(x_0, x_1). \end{aligned} \quad (83)$$

So, let $x_3^{\#\#}$ be an ancestor of x_2'' at distance $h(x_2'') - h(x_3'') - (16\delta + 2\eta)d_\varepsilon(x_0, x_1) \in [0, h(x_2'')]$ from x_2'' , and let $x_0^\#$ be an ancestor of x_1' at distance $h(x_2'') - h(x_3'') - (16\delta + 2\eta)d_\varepsilon(x_0, x_1) \in [0, h(x_1')]$ from x_1' . Then

$$d_\varepsilon(x_3'', x_3^{\#\#}) \leq (16\delta + 2\eta)d_\varepsilon(x_0, x_1), \quad (84)$$

and

$$\begin{aligned} d_\varepsilon(x_0', x_0^\#) &= |h(x_0') - h(x_0^\#)| = |h(x_0') - (h(x_1') - h(x_2'') + h(x_3'') + (16\delta + 2\eta)d_\varepsilon(x_0, x_1))| \\ &\stackrel{(82) \wedge (83)}{\leq} 2(16\delta + 2\eta)d_\varepsilon(x_0, x_1). \end{aligned} \quad (85)$$

Moreover, $h(x_1) - h(x_0^\#) = h(x_2'') - h(x_3^{\#\#})$, so by Lemma 6.9 we have

$$\begin{aligned} \left(\frac{3}{1 + \delta} - 55\delta - 7\eta\right)d_\varepsilon(x_0, x_1) &\leq d_\varepsilon(x_0, x_3) - (55\delta + 7\eta)d_\varepsilon(x_0, x_1) \stackrel{(84) \wedge (85)}{\leq} d_\varepsilon(x_0^\#, x_3^{\#\#}) \\ &\leq d_\varepsilon(x_1', x_2'') \leq (1 + 8\delta + \eta)d_\varepsilon(x_0, x_1), \end{aligned}$$

which is a contradiction since $\max\{\delta, \eta\} < 1/200$.

Lastly, assume for contradiction that (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_1, x_2)$ -near a path-type configuration (x_1'', x_2'', x_3'') . Then since $h(x_1') \leq h(x_2')$ we have

$$\begin{aligned} \left(\frac{1}{1 + \delta} - 14\delta\right)d_\varepsilon(x_0, x_1) &\leq d_\varepsilon(x_1'', x_2'') = h(x_1'') - h(x_2'') \\ &\leq (h(x_1') - h(x_2'')) + (h(x_2') - h(x_1')) \\ &\leq d_\varepsilon(x_1', x_1'') + d_\varepsilon(x_2'', x_2') \leq (14\delta + 2\eta)d_\varepsilon(x_0, x_1), \end{aligned}$$

a contradiction. \square

Lemma 6.26. *Assume that $\varepsilon_n < 1/4$ for all n and that (x_0, x_1, x_2, x_3) is a $(1 + \delta)$ -approximate P_3 such that (x_2, x_1, x_0) is $\eta d_\varepsilon(x_0, x_1)$ -near a tent-type configuration. If $\max\{\delta, \eta\} < 1/200$ then (x_1, x_2, x_3) cannot be $7\delta d_\varepsilon(x_0, x_1)$ near a tent-type configuration.*

Proof. Let (x'_2, x'_1, x'_0) be a tent-type configuration that is $\eta d_\varepsilon(x_0, x_1)$ -near (x_2, x_1, x_0) . Suppose for contradiction that (x_1, x_2, x_3) is $7\delta d_\varepsilon(x_0, x_1)$ -near a tent-type configuration (x''_1, x''_2, x''_3) . Then $h(x'_1) - h(x'_2) = d_\varepsilon(x'_1, x'_2)$, whereas $h(x''_2) - h(x''_1) = d_\varepsilon(x''_1, x''_2)$. Taking the sum of these two inequalities we conclude that

$$d_\varepsilon(x''_1, x''_2) + d_\varepsilon(x'_1, x'_2) \leq d_\varepsilon(x'_1, x''_1) + d_\varepsilon(x'_2, x''_2) \leq (2\eta + 14\delta)d_\varepsilon(x_0, x_1).$$

At the same time, $(\frac{2}{1+\delta} - 2\eta - 14\delta)d_\varepsilon(x_0, x_1) \leq d_\varepsilon(x''_1, x''_2) + d_\varepsilon(x'_1, x'_2)$, which leads to the desired contradiction. \square

Proof of Lemma 6.22. Since (x_0, x_1, x_2, x_3) is a $(1 + \delta)$ -approximate P_3 , we have $x_1 \in \text{Mid}(x_0, x_2, \delta)$, and $x_2 \in \text{Mid}(x_1, x_3, \delta)$. Since the assumptions of Theorem 6.4 hold, we can apply Lemmas 6.23–6.26 with $\eta = 7\delta$ to conclude that either (x_0, x_1, x_2, x_3) or (x_3, x_2, x_1, x_0) must be $35\delta d_\varepsilon(x_0, x_1)$ -near a configuration of type A, B or C. \square

6.3. Nonembeddability of vertically faithful B_4

In what follows we need some standard notation on trees. As before, B_n is the complete binary tree of height n ; the root of B_n is denoted by r . Denote by $I(B_n)$ the set of internal vertices of B_n , i.e., vertices of B_n which are not the root or a leaf. For a vertex v in $\{r\} \cup I(B_n)$ we denote by v_0 and v_1 its children. For $\alpha \in \{0, 1\}^*$ (the set of finite sequences of ‘0’ and ‘1’) and $a \in \{0, 1\}$ we denote $v_{\alpha a} = (v_\alpha)_a$.

The aim of the current section is to prove the following lemma.

Lemma 6.27. *Fix $0 < \delta < 1/400$ and let $f : B_4 \rightarrow (B_\infty, d_\varepsilon)$ be a $(1 + \delta)$ -vertically faithful embedding. Then the distortion of f satisfies*

$$\text{dist}(f) \geq \frac{1}{500\delta + \varepsilon_{h_0}},$$

where $h_0 = \min_{x \in B_4} h(f(x))$.

The proof of Lemma 6.27 is by a contradiction. By Lemma 6.14, assuming the distortion of f is small, all the δ -forks in the $(1 + \delta)$ -vertically faithful embedding must be of types I–IV. By exploring the constraints implied by Lemma 6.22 on how those δ -forks can be “stitched” together, we reach the conclusion that they are sufficiently severe to force any vertically faithful embedding of B_4 to have a large contraction, and therefore high distortion.

Fix $f : B_4 \rightarrow (B_\infty, d_\varepsilon)$. For $u \in I(B_4)$ we denote by $\mathcal{F}(u)$ the fork in which u is the center point, i.e., if v is the parent of u in B_4 , then

$$\mathcal{F}(u) := (f(v), f(u), f(u_0), f(u_1)).$$

We shall assume from now on that f satisfies the assumptions of Lemma 6.27, i.e., f satisfies (8) with $D = 1 + \delta$ for some $\delta < 1/400$ and $\lambda > 0$.

Lemma 6.28. Fix $u \in B_4$ with $h(u) \in \{1, 2\}$. If the fork $\mathcal{F}(u)$ is $37\delta\lambda$ -near a type I or type III configuration, then there exists $w \in I(B_4)$ satisfying

$$d_\varepsilon(f(w_0), f(w_1)) \leq (170\delta + \varepsilon_{h_0}) \cdot 2\lambda. \quad (86)$$

Proof. Let v be the parent of u . Hence, $(f(v), f(u), f(u_0), f(u_1))$ is $35\delta(1 + \delta)\lambda$ -near a type I or a type III configuration. Assume first that $(f(v), f(u), f(u_0), f(u_1))$ is $37\delta\lambda$ -near a type I configuration. If both $(f(u_0), f(u), f(v))$ and $(f(u_1), f(u), f(v))$ were $37\delta\lambda$ -near a path type configuration then by Lemma 6.16 (with $\eta = 37\delta$) we would have

$$d_\varepsilon(f(u_0), f(u_1)) \leq (339\delta + 2\varepsilon_{h_0})(1 + \delta)\lambda \leq (170\delta + \varepsilon_{h_0}) \cdot 2\lambda, \quad (87)$$

proving (86) with $w = u$. The same conclusion holds when $(f(v), f(u), f(u_0), f(u_1))$ is $37\delta\lambda$ -near a type III configuration: in this case without loss of generality $(f(v), f(u), f(u_0))$ is $37\delta\lambda$ -near a tent-type configuration and $(f(u_1), f(u), f(v))$ is $37\delta\lambda$ -near a path-type configuration. Using Lemma 6.16 as above we would arrive at the conclusion (87) if $(f(u_0), f(u), f(v))$ were $37\delta\lambda$ -near a path-type type configuration. Thus, in both the type I and type III cases of Lemma 6.28 we may assume that $(f(v), f(u), f(u_0))$ is $37\delta\lambda$ -near a tent-type configuration, and that, by Lemma 6.8, $(f(v), f(u), f(u_0))$ is not $37\delta\lambda$ -near a path-type configuration, and $(f(u_0), f(u), f(v))$ is not $37\delta\lambda$ -near a path-type configuration or a tent-type configuration.

By Lemma 6.22 (and Table 2) $(f(u_{0c}), f(u_0), f(u), f(v))$ must be $35\delta(1 + \delta)\lambda$ -near a type B configuration for both $c \in \{0, 1\}$. This means that $(f(u_{0c}), f(u_0), f(u))$ are both $35\delta(1 + \delta)\lambda$ -near a path-type configuration, and so by Lemma 6.16 (with $\eta = 35\delta(1 + \delta)$) we deduce that $d_\varepsilon(f(u_{00}), f(u_{01})) \leq (170\delta + \varepsilon_{h_0}) \cdot 2\lambda$. \square

Lemma 6.29. Fix $u \in B_4$ with $h(u) \in \{1, 2\}$. If $\mathcal{F}(u)$ is $37\delta\lambda$ -near a type II configuration then for both $b \in \{0, 1\}$ either $\mathcal{F}(u_b)$ is $99\delta\lambda$ -near a type II configuration, or $d_\varepsilon(f(u_{b0}), f(u_{b1})) \leq 400\delta\lambda$.

Proof. Let v be the parent of u . For both $c \in \{0, 1\}$ we know that $(f(v), f(u), f(u_0), f(u_{0c}))$ is a $(1 + \delta)$ -approximate P_3 , and therefore by Lemma 6.22 either $(f(v), f(u), f(u_0), f(u_{0c}))$ or $(f(u_{0c}), f(u_0), f(u), f(v))$ is $35\delta(1 + \delta)\lambda$ -near a configuration of type A, B or C. Note that since $(f(v), f(u), f(u_0))$ is assumed to be $37\delta\lambda$ -near a path-type configuration, we rule out the possibility that $(f(v), f(u), f(u_0), f(u_{0c}))$ is $35\delta(1 + \delta)\lambda$ -near a configuration of type C, since otherwise both $(f(v), f(u), f(u_0))$ and $(f(u_0), f(u), f(v))$ would be $37\delta\lambda$ -near path-type configurations, contradicting Lemma 6.8. For the same reason we rule out the possibility that $(f(u_{0c}), f(u_0), f(u), f(v))$ is $35\delta(1 + \delta)\lambda$ -near a configuration of type A or type B. An inspection of the three remaining possibilities shows that either $(f(u), f(u_0), f(u_{0c}))$ is $37\delta\lambda$ -near a path-type configuration, or $(f(u_{0c}), f(u_0), f(u))$ is $37\delta\lambda$ -near a tent-type configuration.

Now,

- If for both $c \in \{0, 1\}$ the triple $(f(u), f(u_0), f(u_{0c}))$ is $37\delta\lambda$ -near a path-type configuration, then $\mathcal{F}(u_0)$ is $37\delta\lambda$ -near a type II configuration.
- If for both $c \in \{0, 1\}$ the triple $(f(u_{0c}), f(u_0), f(u))$ is $37\delta\lambda$ -near a tent-type configuration, then by Lemma 6.17 we have $d_\varepsilon(f(u_{01}), f(u_{00})) \leq 400\delta$.

- By Lemma 6.18, the only way that $(f(u), f(u_0), f(u_{00}))$ could be $37\delta\lambda$ -near a path type configuration while at the same time $(f(u_{01}), f(u_0), f(u))$ is $37\delta\lambda$ -near a tent-type configuration (or vice versa), is that $\mathcal{F}(u_0)$ is $99\delta\lambda$ -near a type II configuration. \square

Lemma 6.30. Fix $u \in B_4$ with $h(u) \in \{1, 2\}$. If $\mathcal{F}(u)$ is $35\delta(1 + \delta)\lambda$ -near a type IV configuration, then there exists $b \in \{0, 1\}$ such that $\mathcal{F}(u_b)$ is $37\delta\lambda$ -near a type II configuration.

Proof. Let v be the parent of u . Without loss of generality $(f(u_0), f(u), f(v))$ is $35\delta(1 + \delta)\lambda$ -near a tent-type configuration. By Lemma 6.22 (using Lemma 6.8 to rule out the remaining possibilities), this means that for both $c \in \{0, 1\}$ the quadruple $(f(u_{0c}), f(u_0), f(u), f(v))$ is $35\delta(1 + \delta)^2\lambda$ -near a type C configuration, and therefore $\mathcal{F}(u_0)$ is $35\delta(1 + \delta)^2\lambda$ near a type II configuration. \square

Lemma 6.31. Fix $u \in B_4$ with $h(u) \in \{0, 1, 2\}$. If $\mathcal{F}(u_0)$ and $\mathcal{F}(u_1)$ are both $99\delta\lambda$ -near a type II configuration then $d_\varepsilon(f(u_0), f(u_1)) \leq 1000\delta\lambda$.

Proof. By our assumptions, $(f(u), f(u_0), f(u_{00}))$ is $99\delta\lambda$ -near a path type configuration (u', u'_0, u'_{00}) and $(f(u), f(u_1), f(u_{10}))$ is $99\delta\lambda$ -near a path-type configuration (u'', u''_1, u''_{10}) . We may assume without loss of generality that $h(u'') - h(u'_1) \leq h(u') - h(u'_0)$. We may therefore consider the ancestor u_1^* of u' such that $h(u') - h(u_1^*) = h(u'') - h(u'_1)$, implying in particular that $h(u_1^*) \geq h(u'_0)$ (recall that u'_0 is an ancestor of u' , and u'_1 is ancestor of u''). By Lemma 6.9 we have

$$d_\varepsilon(u_1^*, u''_1) \leq d_\varepsilon(u', u'') \leq 198\delta\lambda. \tag{88}$$

Hence,

$$\begin{aligned} h(u') - h(u_1^*) &= d_\varepsilon(u', u_1^*) \stackrel{(88)}{\geq} d_\varepsilon(u', u''_1) - 198\delta\lambda \geq d_\varepsilon(f(u), f(u_1)) - 394\delta\lambda \\ &\geq (1 - 394\delta)\lambda. \end{aligned} \tag{89}$$

But we also know that

$$h(u') - h(u'_0) = d_\varepsilon(u', u'_0) \leq d_\varepsilon(f(u), f(u_0)) + 198\delta\lambda \leq (1 + 200\delta)\lambda. \tag{90}$$

It follows from (89) and (90) that $d_\varepsilon(u'_0, u_1^*) = h(u_1^*) - h(u'_0) \leq 601\delta\lambda$. Therefore,

$$d_\varepsilon(f(u_1), f(u_0)) \leq d_\varepsilon(f(u_0), u'_0) + d_\varepsilon(u'_0, u_1^*) + d_\varepsilon(u_1^*, u''_1) + d_\varepsilon(u''_1, f(u_1)) = 1000\delta\lambda. \tag{91}$$

Proof of Lemma 6.27. We may assume that for all $u \in I(B_4)$ the fork $\mathcal{F}(u)$ is $35\delta(1 + \delta)\lambda$ -near a configuration of type I, II, III, or IV. Indeed, otherwise the proof is complete by Lemma 6.14. If $\mathcal{F}(r_0)$ is $35\delta(1 + \delta)\lambda$ -near a type I or type III configuration, then by Lemma 6.28 the proof is complete. If $\mathcal{F}(r_0)$ is $35\delta(1 + \delta)\lambda$ -near a type IV configuration then by Lemma 6.30 there exists $b \in \{0, 1\}$ such that $\mathcal{F}(r_{0b})$ is $37\delta\lambda$ -near a type II configuration. It therefore remains to deal with the case in which for some $u \in \{r_0, r_{0b}\}$ the fork $\mathcal{F}(u)$ is $37\delta\lambda$ -near a type II configuration. Applying Lemma 6.29, either we are done, or both $\mathcal{F}(u_0)$ and $\mathcal{F}(u_1)$ are $99\delta\lambda$ -near a type II configuration, but then by Lemma 6.31 the proof of Lemma 6.27 is complete. \square

6.4. Nonembeddability of binary trees

We are now in a position to complete the proof of Theorem 1.10.

Proof of Theorem 1.10. Write $\varepsilon_n = 1/s(n)$ and $\varepsilon = \{\varepsilon_n\}_{n=0}^\infty$. Thus $\{\varepsilon_n\}_{n=0}^\infty$ is nonincreasing, $\{n\varepsilon_n\}_{n=0}^\infty$ is nondecreasing, and $\varepsilon_n \leq 1/4$. We can therefore choose the metric space $(X, d_X) = (B_\infty, d_\varepsilon)$. The identity embedding of B_n into the top n -levels of B_∞ shows that $c_X(B_n) \leq s(n)$. It remains to prove the lower bound on $c_X(B_n)$. To this end take an arbitrary injection $f : B_n \rightarrow X$ satisfying $\text{dist}(f) \leq s(n)$, and we will now prove that

$$\text{dist}(f) \geq s\left(\left\lfloor \frac{n}{40s(n)} \right\rfloor\right) \left(1 - \frac{Cs(n) \log s(n)}{\log n}\right). \tag{91}$$

By adjusting the constant C in (91), we may assume below that n is large enough, say, $n \geq 100$. Write $h_0 = \lfloor n/(40s(n)) \rfloor$ and define $X_{>h_0} = \{x \in B_\infty : h(x) > h_0\}$. We claim that there exists a complete binary subtree $T \subseteq B_n$ of height at least $\lceil n/3 \rceil$ such that we have $f(T) \subseteq X_{>h_0}$. Indeed, let $h_{\min} = \min\{h(x) : x \in f(B_n)\}$ and $h_{\max} = \max\{h(x) : x \in f(B_n)\}$. If $h_{\min} > h_0$ then $f(B_n) \subseteq X_{>h_0}$, and we can take $T = B_n$. So assume that $h_{\min} < h_0$. Since f is an injection it must satisfy $h_{\max} \geq n$. Hence $\|f\|_{\text{Lip}} \geq \frac{h_{\max} - h_{\min}}{2n} \geq \frac{n - h_0}{2n} \geq \frac{1}{4}$. Since $\text{dist}(f) \leq s(n)$ we conclude that $\|f^{-1}\|_{\text{Lip}} \leq 4s(n)$. It follows that, since $\text{diam}(X \setminus X_{>h_0}) \leq 2h_0$, we have $\text{diam}(f^{-1}(X \setminus X_{>h_0})) \leq 8h_0s(n) \leq n/5$. If the top $\lceil n/3 \rceil$ levels of B_n are mapped into $X_{>h_0}$ then we are done, so assume that there exists $u \in f^{-1}(X \setminus X_{>h_0})$ of depth at most $\leq \lceil n/3 \rceil$. In this case $f^{-1}(X \setminus X_{>h_0})$ must be contained in the first $\lceil n/3 \rceil + n/5 < 2n/3 - 1$ levels of B_n , so we can take T to be any subtree of B_n contained in the last $\lceil n/3 \rceil$ levels of B_n .

Fix $\delta \in (0, 1)$. By Theorem 1.14 (with $t = 4$, $D = s(n)$ and $\xi = \delta$), there exists a universal constant $\kappa > 0$ such that if $n \geq s(n)^{\kappa/\delta}$ then there is a mapping $\phi : B_4 \rightarrow B_n$ with $\text{dist}(\phi) \leq 1 + \delta$ such that $f \circ \phi$ is a $(1 + \delta)$ -vertically faithful embedding of B_4 into $X_{>h_0}$. Choosing $\delta = \kappa(\log s(n))/\log n$, by increasing C in (91) if necessary, we may assume that $\delta < 1/400$. Lemma 6.27 then implies

$$(1 + \delta) \text{dist}(f) \geq \text{dist}(f \circ \phi) \geq \frac{1}{500\delta + \varepsilon_{h_0}} = \frac{1}{500\kappa \frac{\log s(n)}{\log n} + \frac{1}{s(\lfloor n/(40s(n)) \rfloor)}}.$$

The deduction of (7) from (6) is a simple exercise: if $s(n) = o((\log n)/\log \log n)$ then we have $(s(n) \log s(n))/\log n = o(1)$. The desired claim will then follow once we check that

$$\limsup_{n \rightarrow \infty} \frac{s(\lfloor n/(40s(n)) \rfloor)}{s(n)} = 1. \tag{92}$$

Indeed, if (92) failed then there would exist $\varepsilon_0 \in (0, 1)$ and $n_0 \in \mathbb{N}$ such that for all $n \geq n_0$,

$$s(\lfloor n/\log n \rfloor) \leq s(\lfloor n/(40s(n)) \rfloor) \leq (1 - \varepsilon_0)s(n). \tag{93}$$

Iterating (93), it would follow that $s(n_j) \geq n_j^{\Omega(1)}$ for some subsequence $\{n_j\}_{j=1}^\infty$, a contradiction. \square

Proof of Theorem 1.12. The proof is identical to the above argument: all one has to notice is that when $s(n) = D$ for all $n \in \mathbb{N}$, the resulting metric d_ε on B_∞ is D -equivalent to the original shortest path metric on B_∞ . In this case, if $c_X(B_n) \leq D - \varepsilon$ then the bound (91) implies that $n \leq D^{CD^2/\varepsilon}$. \square

7. Discussion and open problems

A very interesting problem that arises naturally from Theorem 1.3 and is also a part of the Ribe program, is to find a metric characterization of q -smoothness. A Banach space $(X, \|\cdot\|_X)$ is called q -smooth if it admits an equivalent norm $\|\!\| \cdot \|\!$ such that there is a constant $S > 0$ satisfying

$$\|\!\|x\|\! = 1 \wedge y \in X \Rightarrow \frac{\|\!\|x+y\|\! + \|\!\|x-y\|\!}{2} \leq 1 + S\|\!\|y\|\!^q.$$

A Banach space X is p -convex if and only if its dual space X^* is q -smooth, where $1/p + 1/q = 1$ [17]. It is known that a Banach space X is p -convex for some $p < \infty$ (i.e., superreflexive) if and only if it is q -smooth for some $q > 1$ (this follows from [10, 29]). Hence Bourgain's metric characterization of superreflexivity can be viewed as a statement about uniform smoothness as well. However, we still lack a metric characterization of the more useful notion of q -smoothness. Trees are natural candidates for finite metric obstructions to p -convexity, but it is unclear what would be the possible finite metric witnesses to the "non- q -smoothness" of a metric space.

H -trees are geometric objects that are quite simple combinatorially, yet as we have seen, they have interesting bi-Lipschitz properties. It would therefore be of interest to investigate the geometry of H -trees in its own right. In particular, what is the L_1 distortion of an H -tree? How close can an H -tree be to a metric of negative type?

Acknowledgments. Research of M. Mendel was supported by ISF grant 221/07, BSF grant 2006009, and a gift from Cisco research center.

Research of A. Naor was supported by NSF grants CCF-0635078 and CCF-0832795, BSF grant 2006009, and the Packard Foundation.

References

- [1] Ball, K.: Markov chains, Riesz transforms and Lipschitz maps. *Geom. Funct. Anal.* **2**, 137–172 (1992) [Zbl 0788.46050](#) [MR 1159828](#)
- [2] Ball, K., Carlen, E. A., Lieb, E. H.: Sharp uniform convexity and smoothness inequalities for trace norms. *Invent. Math.* **115**, 463–482 (1994) [Zbl 0803.47037](#) [MR 1262940](#)
- [3] Bates, S., Johnson, W. B., Lindenstrauss, J., Preiss, D., Schechtman, G.: Affine approximation of Lipschitz functions and nonlinear quotients. *Geom. Funct. Anal.* **9**, 1092–1127 (1999) [Zbl 0954.46014](#) [MR 1736929](#)
- [4] Benyamini, Y., Lindenstrauss, J.: *Geometric Nonlinear Functional Analysis*. Vol. 1. *Amer. Math. Soc. Colloq. Publ.* 48, Amer. Math. Soc., Providence, RI (2000) [Zbl 0946.46002](#) [MR 1727673](#)

- [5] Bourgain, J.: The metrical interpretation of superreflexivity in Banach spaces. *Israel J. Math.* **56**, 222–230 (1986) [Zbl 0643.46013](#) [MR 0880292](#)
- [6] Bourgain, J., Milman, V., Wolfson, H.: On type of metric spaces. *Trans. Amer. Math. Soc.* **294**, 295–317 (1986) [Zbl 0617.46024](#) [MR 0819949](#)
- [7] Enflo, P.: Banach spaces which can be given an equivalent uniformly convex norm. In: *Proceedings of the International Symposium on Partial Differential Equations and the Geometry of Normed Linear Spaces* (Jerusalem, 1972), *Israel J. Math.* **13**, 281–288 (1972) [Zbl 0259.46012](#) [MR 0336297](#)
- [8] Gromov, M.: *Metric Structures for Riemannian and non-Riemannian Spaces*. Modern Birkhäuser Classics, Birkhäuser Boston, Boston, MA (2007) [Zbl 1113.53001](#) [MR 2307192](#)
- [9] James, R. C.: Uniformly non-square Banach spaces. *Ann. of Math. (2)* **80**, 542–550 (1964) [Zbl 0132.08902](#) [MR 0173932](#)
- [10] James, R. C.: Some self-dual properties of normed linear spaces. In: *Symposium on Infinite-Dimensional Topology* (Baton Rouge, LA, 1967), *Ann. of Math. Stud.* 69, Princeton Univ. Press, Princeton, NJ, 159–175 (1972) [Zbl 0233.46025](#) [MR 0454600](#)
- [11] Johnson, W. B., Schechtman, G.: Diamond graphs and super-reflexivity. *J. Topol. Anal.* **1**, 177–189 (2009) [Zbl 1183.46022](#) [MR 2541760](#)
- [12] Laakso, T. J.: Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbb{R}^N . *Bull. London Math. Soc.* **34**, 667–676 (2002) [Zbl 1029.30014](#) [MR 1924353](#)
- [13] Lang, U., Plaut, C.: Bilipschitz embeddings of metric spaces into space forms. *Geom. Dedicata* **87**, 285–307 (2001) [Zbl 1024.54013](#) [MR 1866853](#)
- [14] Lee, J. R., Mendel, M., Naor, A.: Metric structures in L_1 : dimension, snowflakes, and average distortion. *Eur. J. Combin.* **26**, 1180–1190 (2005) [Zbl 1106.68086](#) [MR 2163751](#)
- [15] Lee, J. R., Naor, A.: Embedding the diamond graph in L_p and dimension reduction in L_1 . *Geom. Funct. Anal.* **14**, 745–747 (2004) [Zbl 1069.46005](#) [MR 2084978](#)
- [16] Lee, J. R., Naor, A., Peres, Y.: Trees and Markov convexity. *Geom. Funct. Anal.* **18**, 1609–1659 (2009) [Zbl 1171.05318](#) [MR 2481738](#)
- [17] Lindenstrauss, J.: On the modulus of smoothness and divergent series in Banach spaces. *Michigan Math. J.* **10**, 241–252 (1963) [Zbl 0115.10001](#) [MR 0169061](#)
- [18] Matoušek, J.: On embedding trees into uniformly convex Banach spaces. *Israel J. Math.* **114**, 221–237 (1999) [Zbl 0948.46011](#) [MR 1738681](#)
- [19] Matoušek, J.: Ramsey-like properties for bi-Lipschitz mappings of finite metric spaces. *Comment. Math. Univ. Carolin.* **33**, 451–463 (1992) [Zbl 0769.05093](#) [MR 1209287](#)
- [20] Maurey, B., Pisier, G.: Séries de variables aléatoires vectorielles indépendantes et propriétés géométriques des espaces de Banach. *Studia Math.* **58**, 45–90 (1976) [Zbl 0344.47014](#) [MR 0443015](#)
- [21] Mendel, M.: Metric dichotomies. In: *Limits of Graphs in Group Theory and Computer Science*, EPFL Press, Lausanne, 59–76 (2009) [Zbl pre05722684](#) [MR 2562140](#)
- [22] Mendel, M., Naor, A.: Scaled Enflo type is equivalent to Rademacher type. *Bull. London Math. Soc.* **39**, 493–498 (2007) [Zbl 1126.46008](#) [MR 2331580](#)
- [23] Mendel, M., Naor, A.: Markov convexity and local rigidity of distorted metrics [extended abstract]. In: *Computational Geometry (SCG'08)*, ACM, New York, 49–58 (2008) [Zbl 1192.52010](#) [MR 2504270](#)
- [24] Mendel, M., Naor, A.: Metric cotype. *Ann. of Math. (2)* **168**, 247–298 (2008) [Zbl 1187.46014](#) [MR 2415403](#)
- [25] Naor, A., Schechtman, G.: Remarks on non linear type and Pisier's inequality. *J. Reine Angew. Math.* **552**, 213–236 (2002) [Zbl 1033.46013](#) [MR 1940437](#)
- [26] Newman, I., Rabinovich, Y.: A lower bound on the distortion of embedding planar metrics into Euclidean space. *Discrete Comput. Geom.* **29**, 77–81 (2003) [Zbl 1050.68154](#) [MR 1946795](#)

-
- [27] Odell, E., Schlumprecht, T.: The distortion problem. *Acta Math.* **173**, 259–281 (1994)
[Zbl 0828.46005](#) [MR 1301394](#)
- [28] Pisier, G.: Sur les espaces de Banach qui ne contiennent pas uniformément de l_n^1 . *C. R. Acad. Sci. Paris Sér. A-B* **277**, A991–A994 (1973) [Zbl 0271.46011](#) [MR 0333673](#)
- [29] Pisier, G.: Martingales with values in uniformly convex spaces. *Israel J. Math.* **20**, 326–350 (1975) [Zbl 0344.46030](#) [MR 0394135](#)
- [30] Pisier, G.: Probabilistic methods in the geometry of Banach spaces. In: *Probability and Analysis (Varenna, 1985)*, Lecture Notes in Math. 1206, Springer, Berlin, 167–241 (1986)
[Zbl 0606.60008](#) [MR 0864714](#)
- [31] Ribe, M.: On uniformly homeomorphic normed spaces. *Ark. Mat.* **14**, 237–244 (1976)
[Zbl 0336.46018](#) [MR 0440340](#)