

Modules of minimal dimension over completed Weyl algebras

Feliks Rączka

Abstract. We study the category of modules of minimal dimension over completed Weyl algebras in equal characteristic zero. In particular, we prove finiteness of de Rham cohomology of such modules.

1. Introduction

This is the first of two papers devoted to the study of D -modules on rigid analytic varieties over the field $\mathbb{C}((z))$ of formal Laurent series (and, more generally, over discretely valued nonarchimedean fields of equal characteristic zero). Here, we study modules of minimal dimension over completed Weyl algebras and we prove that their de Rham cohomology groups are finite dimensional. In the sequel [11] we use this result to prove its global analogue: on a (quasi-compact and quasi-separated) smooth rigid analytic variety the de Rham cohomology groups of a holonomic D -module have finite dimensions.

Let K be a discretely valued nonarchimedean field, and let $\mathfrak{o}_K$ be its valuation ring. We also fix a uniformizer $\varpi \in \mathfrak{o}_K$. Let $\mathbb{W}_n(\mathfrak{o}_K)$ denote the n -th Weyl algebra over $\mathfrak{o}_K$, i.e., the noncommutative $\mathfrak{o}_K$ -algebra generated by $x_1, \dots, x_n, \partial_1, \dots, \partial_n$ with relations $[x_i, x_j] = [\partial_i, \partial_j] = 0$ and $[\partial_i, x_j] = \delta_{ij}$. We set

$$\hat{\mathcal{D}}_n^{\circ} \stackrel{\text{def}}{=} \varprojlim_{s \geq 1} \frac{\mathbb{W}_n(\mathfrak{o}_K)}{\varpi^{s+1} \mathbb{W}_n(\mathfrak{o}_K)} \quad (1.1)$$

and define the n -th *completed Weyl algebra* over K as

$$\hat{\mathcal{D}}_n \stackrel{\text{def}}{=} \hat{\mathcal{D}}_n^{\circ} \otimes_{\mathfrak{o}_K} K. \quad (1.2)$$

Algebraic properties of completed Weyl algebras have been studied by many authors, for example by L. Narváez Macarro in [9], and more recently by A. Pangalos in [10]. Our objective is to study de Rham cohomology of $\hat{\mathcal{D}}_n$ -modules of minimal dimension. We start by recalling these notions. The *de Rham complex* of a left $\hat{\mathcal{D}}_n$ -module M is

$$\mathrm{DR}^s(M) = \bigoplus_{|I|=s} M \cdot dx_I, \quad (1.3)$$

where $0 \leq s \leq n$ is an integer, $I = (1 \leq i_1 < \cdots < i_s \leq n)$ is a multi-index, and $dx_I = dx_{i_1} \wedge \cdots \wedge dx_{i_s}$. The differential δ is given by

$$\delta(m \cdot dx_I) = \sum_{i=1}^n \partial_i m \cdot dx_i \wedge dx_I. \quad (1.4)$$

The *de Rham cohomology* of M is the cohomology of this complex

$$H_{dR}^i(M) := H^i(\mathrm{DR}^\bullet(M)). \quad (1.5)$$

Although we will not need this interpretation in what follows, we remark that the n -th Tate algebra $T_n = K\langle x_1, \dots, x_n \rangle$ is a $\hat{\mathcal{D}}_n$ -module in a natural way and that (1.5) may be also described as

$$H_{dR}^i(M) = \mathrm{Ext}_{\hat{\mathcal{D}}_n}^i(T_n, M).$$

Given a left and right noetherian ring R of homological dimension n we say that a left R -module M is of *minimal dimension* if it is finitely generated and $\mathrm{Ext}_R^i(M, R) = 0$ for $i < n$. This is the algebraization of the geometric notion of holonomicity in the sense that if R is a ring of differential operators of the affine algebra of a smooth complex algebraic variety X then left R -modules of minimal dimension correspond to holonomic D -modules on X . It is easy to show that the ring $\hat{\mathcal{D}}_n$ is left and right noetherian (Lemma 2.3), and it has been shown by A. Pangalos that if K is of equal characteristic zero then it has homological dimension n (Lemma 2.4). Hence, it is meaningful to consider $\hat{\mathcal{D}}_n$ -modules of minimal dimension.

The main result of this paper is the following theorem.

Theorem 1.1. *Let K be a discretely valued nonarchimedean field of equal characteristic zero and let M be a left $\hat{\mathcal{D}}_n$ -module of minimal dimension. Then $\dim_K H_{dR}^i(M) < \infty$ for all i .*

Let k be the residue field of $\mathfrak{o}_K$. The idea for the proof of Theorem 1.1 is to study $\hat{\mathcal{D}}_n^\circ$ -modules and compare their properties on the generic and the special fiber, i.e., after tensoring with K and k respectively. Consider the ring

$$\bar{\mathcal{D}}_n \stackrel{\mathrm{def}}{=} \hat{\mathcal{D}}_n^\circ / \varpi \hat{\mathcal{D}}_n^\circ = \hat{\mathcal{D}}_n^\circ \otimes_{\mathfrak{o}_K} k.$$

It is isomorphic to the n -th Weyl algebra over k , and if $\mathrm{char} k = 0$ then $\bar{\mathcal{D}}_n$ is the ring of algebraic differential operators of the affine n -space $\mathbb{A}_k^n$. If M is a finitely generated left $\hat{\mathcal{D}}_n$ -module then a *lattice* in M is a finitely generated $\hat{\mathcal{D}}_n^\circ$ -submodule $L \subset M$ such that $L \otimes_{\mathfrak{o}_K} K = M$. We write $\bar{L}$ for the left $\bar{\mathcal{D}}_n$ -module $L \otimes_{\mathfrak{o}_K} k$ and call it the *reduction* of L . A more refined version of Theorem 1.1 is the following theorem.

Theorem 1.2. *Let K be a discretely valued nonarchimedean field of equal characteristic zero, and let M be a finitely generated left $\hat{\mathcal{D}}_n$ -module. Then the following conditions are equivalent:*

- (1) M is of minimal dimension.
- (2) There exists a lattice $L \subset M$ such that $\bar{L}$ is a $\bar{\mathcal{D}}_n$ -module of minimal dimension.
- (3) For any lattice $L \subset M$ the reduction $\bar{L}$ is a $\bar{\mathcal{D}}_n$ -module of minimal dimension.

Moreover, if these equivalent conditions are satisfied, then

- (A) The semisimplification of $\bar{L}$ does not depend on L .
- (B) We have $\dim_K H_{dR}^i(M) < \infty$ for all i and the equality $\chi_{dR}(M) = \chi_{dR}(\bar{L})$ holds.

Here and later,

$$\chi_{dR}(M) = \sum (-1)^i \dim_K H_{dR}^i(M)$$

is the Euler characteristic for de Rham cohomology and $\chi_{dR}(\bar{L})$ is the Euler characteristic for de Rham cohomology of the holonomic $\mathcal{D}_{\mathbb{A}_k^n}$ -module $\bar{L}$, which is known to be finite since $\text{char } k = 0$.

We briefly explain why Theorem 1.1 does not hold when K is of mixed characteristic. If K has equal characteristic zero, the Tate algebra T_n is a $\hat{\mathcal{D}}_n$ -module of minimal dimension. We do not know if it is always of minimal dimension when K has mixed characteristic, but it is the case for $n = 1$ by [10, Prop. 2.2.3]. The de Rham complex of $T_1 = K\langle x \rangle$ is

$$\frac{d}{dx} : K\langle x \rangle \rightarrow K\langle x \rangle.$$

If $\text{char } k = p > 0$ then $|p| < 1$ and any formal power series of form

$$\sum_{n \geq 0} a_n p^n x^{p^n - 1}, \quad (1.6)$$

where $|a_n| = 1$, is an element of $K\langle x \rangle$. These elements cannot be integrated with respect to x in the sense that the formal power series

$$\int \sum_{n \geq 0} a_n p^n x^{p^n - 1} = \sum_{n \geq 0} a_n x^{p^n}$$

is not convergent for $|x| = 1$ and thus not an element of $K\langle x \rangle$. We conclude that $\dim_K H_{dR}^1(K\langle x \rangle)$ is infinite. Note that if the residue characteristic of K is zero, then $|p| = 1$ and power series of form (1.6) are not elements of $K\langle x \rangle$. More generally, in the case of equal characteristic zero, if $f \in K[[x]]$ and $\frac{df}{dx} \in K\langle x \rangle$ then $f \in K\langle x \rangle$, i.e., every element of $K\langle x \rangle$ can be integrated with respect to x and $H_{dR}^1(K\langle x \rangle) = 0$. Therefore Theorem 1.1 works in this simple case.

The paper is organized as follows. In Section 2 we recall basic results about completed Weyl algebras and modules of minimal dimension. In Section 3 we discuss some more sophisticated properties of modules over $\mathfrak{o}_K$ -algebras. In Section 4 we apply results from previous sections to give the proof of Theorem 1.2.

2. Preliminaries

From now till the end of this paper K is a fixed discretely valued nonarchimedean field of equal characteristic zero. We let $\mathfrak{o}_K \subset K$ be the valuation ring and k the residue field. We fix a uniformizer $\varpi \in \mathfrak{o}_K$. Although we will not use it, we note that by the Cohen structure theorem a choice of a uniformizer gives an isomorphism $K = k((\varpi))$.

2.1. Modules of minimal dimension

In this subsection we recall basic properties of modules of minimal dimension and basic properties of holonomic $\mathcal{D}_{\mathbb{A}_k^n}$ -modules.

Let R be a (not necessarily commutative) ring. We recall that the projective dimension of an R -module M (written $pd(M)$) is the infimum of lengths of its projective resolutions. Then we define the left global dimension of R as

$$\text{l.gl.dim}(R) = \sup \{pd(M) : M \text{ is a left } R\text{-module}\}.$$

The right global dimension (written $\text{r.gl.dim}(R)$) is defined in the same way. If R is left and right noetherian then by [15, Exer. 4.1.1] left and right global dimensions of R are equal and we define the *global dimension* of R as

$$\text{gl.dim}(R) = \text{l.gl.dim}(R) = \text{r.gl.dim}(R).$$

Now, let R be a left and right noetherian ring of finite global dimension $\text{gl.dim}(R) = n$. Following [8, Sec. 1.2] we say that a finitely generated left (resp. right) R -module M is of *minimal dimension* if

$$\inf \{i : \text{Ext}_R^i(M, R) \neq 0\} = n.$$

For such module we set

$$M^* = \text{Ext}_R^n(M, R).$$

The following lemma is well known. Since it is an important ingredient in the proof of Theorem 1.2 we sketch the proof for completeness.

Lemma 2.1. *Let M be a left (resp. right) R -module of minimal dimension. Then M^* is a right (resp. left) R -module of minimal dimension and $M^{**} = M$.*

Proof. It is well known that if P is a finitely generated projective left (resp. right) module, then its dual $P^\vee = \text{Hom}_R(P, R)$ is a finitely generated projective right (resp. left) module and the natural map $P \rightarrow P^{\vee\vee}$ is an isomorphism. Let M be a left (resp. right) R -module of minimal dimension. Since R is noetherian and of finite global dimension, M admits a finite projective resolution by finitely generated projective modules. Let $P_\bullet$ be such resolution and let $Q_\bullet = \text{Hom}_R(P_\bullet, R)[n]$. First of all, we have $H_i(Q_\bullet) = \text{Ext}_R^{n-i}(M, R)$ and

therefore $Q_\bullet$ is a projective resolution of M^* . By reflexivity of finite projective modules we have $P_\bullet = \text{Hom}_R(Q_\bullet, R)[n]$ and therefore

$$\text{Ext}_R^i(M^*, R) = H_{n-i}(P_\bullet) = \begin{cases} 0 & \text{if } i \neq n, \\ M & \text{if } i = n. \end{cases}$$

This shows that M^* is of minimal dimension and that $M^{**} = M$. ■

If $R = \mathbb{W}_n(k)$ for some field k of characteristic zero, then the category of left R -modules of minimal dimension coincides with the category of holonomic $\mathcal{D}_{\mathbb{A}_k^n}$ -modules. This category is very well understood and for example the following properties of holonomic modules are well known (de Rham complex and de Rham cohomology of a $\mathbb{W}_n(k)$ -module M is defined by formulas (1.3) and (1.5) of the introduction).

Lemma 2.2. *Let M, M', M'' be finitely generated left $\mathbb{W}_n(k)$ -modules. Then*

- (a) *If $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ is a short exact sequence then M is of minimal dimension if and only if M' and M'' are of minimal dimension.*
- (b) *$\text{Ext}_{\mathbb{W}_n(k)}^n(M, \mathbb{W}_n(k))$ is a right $\mathbb{W}_n(k)$ -module of minimal dimension.*
- (c) *If M is of minimal dimension then it has finite length as a $\mathbb{W}_n(k)$ -module.*
- (d) *If M is of minimal dimension then $\dim_k H_{dR}^i(M) < \infty$ for all i .*

The same holds with right and left replaced.

Proof. Properties (a), (b) and (c) are discussed in [8, Sec. 1.2]. Property (a) is discussed after Def. 1.2.4 of op. cit. and (b) is a consequence of Thm. 1.2.2 of op. cit. See also [7, 4.2.17] for the proof of (b). Property (c) is [8, Prop. 1.2.5]. The last statement is a special case of the classical theorem of Bernstein which states that higher direct images in the derived category of $\mathcal{D}$ -modules preserve holonomicity. We refer to [2, Ch. 1, Thm. 6.1] for the proof of this theorem in the case of Weyl algebras and to [3, Thm. 3.2.3] for the proof in full generality. ■

2.2. Completed Weyl Algebras

In this subsection, we discuss some basic properties of the completed Weyl Algebras, i.e., rings $\hat{\mathcal{D}}_n^\circ$ and $\hat{\mathcal{D}}_n$ defined by formulas (1.1) and (1.2) of the introduction.

Lemma 2.3. *Both $\hat{\mathcal{D}}_n^\circ$ and $\hat{\mathcal{D}}_n$ are left and right noetherian.*

Proof. First of all, the ring $\mathbb{W}_n(\mathfrak{o}_K)$ is left and right noetherian. Indeed, the associated graded ring of the Bernstein filtration $F_n \mathbb{W}_n(\mathfrak{o}_K) = \bigoplus_{|\alpha|+|\beta| \leq n} a_{\alpha\beta} x^\alpha \partial^\beta$ is the polynomial ring in $2n$ variables over $\mathfrak{o}_K$. Since the valuation on K is discrete $\mathfrak{o}_K$ is noetherian and therefore so is any polynomial ring over $\mathfrak{o}_K$. We can apply [3, Prop. D.1.4] which states that if the associated graded ring is noetherian then so is the original ring. Now, it is well known in the commutative case that for a noetherian ring R its I -adic completion is again noetherian. While this needs not be the case for noncommutative rings, it follows

from [6, Prop. 2.1] that the Theorem remains true if I is a two sided ideal generated by a single central element. Because $\mathbb{W}(\mathfrak{o}_K)$ is left and right noetherian and ϖ is central we conclude that $\widehat{\mathcal{D}}_n^\circ$ is left and right noetherian. Then $\widehat{\mathcal{D}}_n$ is left and right noetherian because it is a localization of $\widehat{\mathcal{D}}_0$ at ϖ . This proves the first part of the lemma. ■

The following result is taken from the Ph.D. thesis of A. Pangalos [10].

Lemma 2.4 (A. Pangalos). $\text{gl.dim}(\widehat{\mathcal{D}}_n) = n$.

Proof. [10, Prop. 3.1.3] gives a bound $\text{gl.dim}(\widehat{\mathcal{D}}_n) \geq n$ and Prop. 4.3.6 in op.cit. gives a bound $\text{gl.dim}(\widehat{\mathcal{D}}_n) \leq n$. ■

We will also need the following properties of $\widehat{\mathcal{D}}_n$ -lattices. Recall that $\bar{L} = L \otimes_{\mathfrak{o}_K} k$.

Lemma 2.5. *Let L be a finitely generated left $\widehat{\mathcal{D}}_n^\circ$ -module. Then:*

- (a) L is complete in the ϖ -adic topology.
- (b) If $\bar{L} = 0$ then $L = 0$.

Proof. Since ϖ is central, we can use the same reasoning as in the case of commutative noetherian rings. Since by [13, p. 413] the Artin–Rees lemma holds for finitely generated left $\widehat{\mathcal{D}}_n^\circ$ -modules, we can proceed as in [1, Ch. 10] to check that if $I = (\varpi)$ and L is a finitely generated left $\widehat{\mathcal{D}}_n^\circ$ -module, then

$$\widehat{L}^I = \widehat{\mathcal{D}}_n^{\circ I} \otimes_{\widehat{\mathcal{D}}_n^\circ} L = \widehat{\mathcal{D}}_n^\circ \otimes_{\widehat{\mathcal{D}}_n^\circ} L = L.$$

This proves the first statement of the lemma, and the second statement follows from Nakayama’s lemma for separated modules because $L = \widehat{L}^I$ is separated. ■

3. Algebras over discrete valuation rings

It is possible that Lemmas 3.1, 3.2 and 3.3 below are known to the experts but we are not aware of any published proof in the form that we need. We will only use these results in case of the ring $\widehat{\mathcal{D}}_n$ to prove Theorem 1.2 but since the proofs would not become easier nor shorter after restricting to this special case we present them in a more general setting.

For the purpose of this section we assume that B_0 is a (not necessarily commutative) ring and $\pi \in B_0$ is a central element that is not a zero divisor. We set $B = B_0[\pi^{-1}]$ and $\bar{B} = B_0/\pi B_0$. Because π is not a zero divisor the natural map $B_0 \rightarrow B$ is injective and we may write

$$B = \bigcup_{n \in \mathbb{Z}} \pi^n B_0.$$

A model example of this situation is when π is a uniformizer of some discrete valuation ring $\mathcal{O}$ and B_0 is a flat $\mathcal{O}$ -algebra.

3.1. Künneth type short exact sequences

Lemma 3.1. *Let M be a right B_0 -module that is π -torsion-free and has a finite projective resolution by finitely generated modules. Then for each $i \geq 0$ there are short exact sequences of left $\bar{B}$ -modules*

$$0 \rightarrow \bar{B} \otimes_{B_0} \text{Ext}_{B_0}^i(M, B_0) \rightarrow \text{Ext}_{\bar{B}}^i(M \otimes_{B_0} \bar{B}, \bar{B}) \rightarrow \text{Tor}_1^{B_0}(\bar{B}, \text{Ext}_{B_0}^{i+1}(M, B_0)) \rightarrow 0.$$

The same holds with the left and right replaced and obvious modifications.

Proof. Note that $\bar{B}$ has a projective resolution

$$0 \rightarrow B_0 \xrightarrow{\times \pi} B_0 \rightarrow \bar{B} \rightarrow 0, \quad (3.1)$$

and therefore for any right B_0 -module M we have $\text{Tor}_i^{B_0}(M, \bar{B}) = 0$ for $i \geq 2$ and

$$\text{Tor}_1^{B_0}(M, \bar{B}) = \{m \in M : m\pi = 0\}.$$

It follows that if M is π -torsion-free and if

$$P^\bullet = [0 \rightarrow P^{-n} \rightarrow \dots \rightarrow P^{-1} \rightarrow P^0 \rightarrow 0]$$

is a projective resolution of M by finitely generated modules then

$$\bar{P}^\bullet = P^\bullet \otimes_{B_0} \bar{B}$$

is a projective resolution of $M \otimes_{B_0} \bar{B}$. Set

$$Q_\bullet = \text{Hom}_{B_0}(P^\bullet, B_0). \quad (3.2)$$

This is a complex of finitely generated projective left B_0 -modules and we have

$$H_i(Q_\bullet) = \text{Ext}_{B_0}^{-i}(M, B_0). \quad (3.3)$$

On the other hand,

$$\text{Ext}_{\bar{B}}^{-i}(M \otimes_{B_0} \bar{B}, \bar{B}) = H_i(\text{Hom}(\bar{P}^\bullet, \bar{B})) = H_i(\bar{B} \otimes_{B_0} Q_\bullet). \quad (3.4)$$

Here the first equality holds because $\bar{P}^\bullet$ is a projective resolution of $M \otimes_{B_0} \bar{B}$ and the second equality holds because for any finitely generated projective right B_0 -module P we have natural isomorphisms of left $\bar{B}$ -modules

$$\bar{B} \otimes_{B_0} \text{Hom}_{B_0}(P, B_0) = \text{Hom}_{B_0}(P, \bar{B}) = \text{Hom}_{\bar{B}}(P \otimes_{B_0} \bar{B}, \bar{B}).$$

Consider the following claim: *If $Q_\bullet$ is a bounded chain complex of finitely generated projective left B_0 -modules then there exist exact sequences of left $\bar{B}$ -modules*

$$0 \rightarrow \bar{B} \otimes_{B_0} H_j(Q_\bullet) \rightarrow H_j(\bar{B} \otimes_{B_0} Q_\bullet) \rightarrow \text{Tor}_1^{B_0}(\bar{B}, H_{j-1}(Q_\bullet)) \rightarrow 0. \quad (3.5)$$

Once we have proven the claim we are done with the proof because of equalities (3.3) and (3.4). The fastest way to show existence of exact sequences (3.5) is to use Künneth's spectral sequence [15, Thm. 5.6.4] (see also [12, Thm. 10.90] for the formulation over noncommutative rings)

$$E_{i,j}^2 = \mathrm{Tor}_i^{B_0}(\bar{B}, H_j(Q_\bullet)) \implies H_{i+j}(\bar{B} \otimes_{B_0} Q_\bullet). \quad (3.6)$$

Because of (3.1) we have $\mathrm{Tor}^i(\bar{B}, -) = 0$ for $i \neq 0, 1$ and the above spectral sequence degenerates to short exact sequences

$$0 \rightarrow E_{0,j}^2 \rightarrow H_j(\bar{B} \otimes_{B_0} Q_\bullet) \rightarrow E_{1,j-1}^2 \rightarrow 0. \quad (3.7)$$

The problem with this approach is that in the literature existence of the spectral sequence (3.6) is usually formulated with $\bar{B}$ replaced by an arbitrary right B_0 -module. As a consequence, one needs to additionally check that maps in sequences (3.7) are $\bar{B}$ -linear and not merely additive (which is usually the case for a tensor product of a left and a right module over a noncommutative ring). Alternatively, we can notice that if $d_\bullet$ is a differential in $Q_\bullet$ then as in the proof of [15, Thm. 3.6.1] we have the short exact sequence of complexes

$$0 \rightarrow \ker d_\bullet \otimes_{B_0} \bar{B} \rightarrow Q_\bullet \otimes_{B_0} \bar{B} \rightarrow \mathrm{im} d_\bullet \otimes_{B_0} \bar{B} \rightarrow 0. \quad (3.8)$$

This is again a consequence of description of $\mathrm{Tor}^i(-, \bar{B})$. Based on this observation we can copy the proof of [15, Thm. 3.6.1] to prove our claim. Then $\bar{B}$ -linearity is clear because the arrows in short exact sequences come from the long exact sequence associated to (3.8). ■

3.2. Reduction of lattices

Lemma 3.2 below is a simple generalization of classical results that appear in many branches of mathematics. For example, in algebraic geometry a variant of Lemma 3.2 for vector bundles with integrable connections is due to O. Gabber and may be found in a book of Katz [4, Variant 2.5.2]. More recently, a similar argument was used by A. Langer in [5]. There is also a variant of Lemma 3.2 in representation theory of finite groups over fields of positive characteristic [14, Thm. 2.2.3].

Our definition of a lattice given in the introduction can be formulated in a more general setting as follows. A *lattice* in a finitely generated B -module M is a finitely generated B_0 -submodule $L \subset M$ such that $B \otimes_{B_0} L = L[\pi^{-1}] = M$. We set $\bar{L} = L/\pi L$. Recall that a module N is of *finite length* if it has finite composition series $0 = N_0 \subset N_1 \subset \dots \subset N_r = N$ where the factors N_i/N_{i-1} are simple modules. The module $N^{\mathrm{ss}} = \bigoplus_{i=1}^r N_i/N_{i-1}$ does not depend on the choice of the composition series and is called the *semisimplification* of N . With the above notation, we prove the following.

Lemma 3.2. *Let M be a finitely generated left B -module, and let $L_1, L_2 \subset M$ be two lattices. If $\bar{L}_1$ has finite length, then so does $\bar{L}_2$ and they have isomorphic semisimplifications.*

Proof. Since $B = \bigcup_{n \in \mathbb{Z}} \pi^n B_0$, there exist integers $n, m \in \mathbb{Z}$ with $\pi^n L_2 \subset L_1 \subset \pi^m L_2$, and since $\overline{\pi^k L_i}$ is isomorphic to $\bar{L}_i$, we may assume that

$$L_2 \subset L_1 \subset \pi^{-n} L_2, \quad (3.9)$$

where $n \geq 1$. We prove the lemma by induction on n and we do the inductive step first. Assume that $n \geq 2$ and that the statement is true for $n - 1$. Then it is also true for n because we have containments

$$L_2 \subset L_1 \cap \pi^{-n+1} L_2 \subset \pi^{-n+1} L_2,$$

and

$$L_1 \cap \pi^{-n+1} L_2 \subset L_1 \subset \pi^{-1} (L_1 \cap \pi^{-n+1} L_2).$$

Now, we deal with the case $n = 1$. We have

$$L_1 \subset \pi^{-1} L_2 \subset \pi^{-1} L_1. \quad (3.10)$$

Taking reductions of (3.9) (for $n = 1$) and of (3.10) gives exact sequences

$$\bar{L}_2 \xrightarrow{\varphi} \bar{L}_1 \xrightarrow{\psi} \bar{L}_2$$

and

$$\bar{L}_1 \xrightarrow{\psi} \bar{L}_2 \xrightarrow{\varphi} \bar{L}_1,$$

where φ (resp. ψ) is the map induced by the inclusion $L_2 \subset L_1$ (resp. $L_1 \subset \pi^{-1} L_2$). From the above, we obtain short exact sequences

$$0 \rightarrow \text{im } \varphi \rightarrow \bar{L}_1 \rightarrow \text{im } \psi \rightarrow 0 \quad (3.11)$$

and

$$0 \rightarrow \text{im } \psi \rightarrow \bar{L}_2 \rightarrow \text{im } \varphi \rightarrow 0. \quad (3.12)$$

If $0 \rightarrow N_1 \rightarrow N \rightarrow N_2 \rightarrow 0$ is a short exact sequence of modules then N has finite length if and only if N_1 and N_2 have finite length, and If this is a case, then $N^{\text{ss}} = N_1^{\text{ss}} \oplus N_2^{\text{ss}}$. The result now follows from existence of short exact sequences (3.11) and (3.12). ■

3.3. Euler characteristic of a perfect complex over a complete discrete valuation ring

For the purpose of the next lemma we will need stronger assumptions. Let B_0 be a complete discrete valuation ring with the residue field ℓ and the quotient field B (so $\ell = \bar{B}$ in the previous notation). We also fix a uniformizer $\pi \in B_0$. Recall that a B_0 -module M is separated for the π -adic topology if $\bigcap_{n \geq 0} \pi^n M = \{0\}$ and is complete if

$$M = \varprojlim_{n \geq 0} M / \pi^{n+1} M.$$

In particular, complete modules are separated.

Lemma 3.3. *Let $(C^\bullet, d^\bullet)$ be a cochain complex of complete (for the π -adic topology), torsion-free B_0 -modules and assume that all ℓ -vector spaces $H^i(C^\bullet \otimes_{B_0} \ell)$ have finite dimensions. Then*

- (1) *All B_0 -modules $H^i(C^\bullet)$ are finitely generated. In particular, all B -vector spaces $H^i(C^\bullet \otimes_{B_0} B)$ have finite dimensions.*
- (2) *If $C^\bullet$ is bounded, then*

$$\sum (-1)^i \dim_\ell H^i(C^\bullet \otimes_{B_0} \ell) = \sum (-1)^i \dim_L H^i(C^\bullet \otimes_{B_0} B), \quad (3.13)$$

i.e., the Euler characteristic of $C^\bullet$ on the special and the generic fibers are equal.

We need the following variant of Nakayama's lemma.

Lemma 3.4. *Let*

$$V \rightarrow W \rightarrow Q \rightarrow 0$$

be an exact sequence of B_0 -modules. Assume that V complete, W is separated, and $Q \otimes_{B_0} \ell$ is finitely generated. Then Q is finitely generated.

Proof. Since the tensor product is right exact, we have a commutative diagram with exact rows.

$$\begin{array}{ccccccc} V & \xrightarrow{\varphi} & W & \xrightarrow{\psi} & Q & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ \bar{V} & \xrightarrow{\bar{\varphi}} & \bar{W} & \xrightarrow{\bar{\psi}} & \bar{Q} & \longrightarrow & 0. \end{array}$$

Pick generators $\bar{q}_1, \dots, \bar{q}_n \in \bar{Q} = Q \otimes_{B_0} \ell$ and let $q_1, \dots, q_n \in Q$ denote lifts of these elements to Q . Let $w_1, \dots, w_n \in W$ satisfy $\psi(w_i) = b_i$. To prove the lemma it suffices to show that for any $x \in W$ there exist $r_1, \dots, r_n \in B_0$ and $v \in V$ such that $w = \sum_{i=1}^n r_i w_i + \psi(v)$. Since $\bar{W}$ is generated mod $\text{im } \bar{\varphi}$ by $\bar{w}_1, \dots, \bar{w}_n$, there exist $r_1^0, \dots, r_n^0 \in B_0$, $v_0 \in V$ and $x_1 \in W$ such that

$$x = \sum_{i=1}^n r_i^0 w_i + \varphi(v_0) + \pi x_1.$$

We can repeat this process for x_1 to find inductively elements $r_i^0, r_i^1, r_i^2, \dots \in B_0$, $v_0, v_1, \dots \in V$ and $x_1, x_2, \dots \in W$ such that for every $m \geq 1$

$$x = \sum_{i=1}^n \left(\sum_{j=0}^m \pi^j r_i^j \right) w_i + \sum_{j=0}^m \pi^j \varphi(v_j) + \pi^{m+1} x_{m+1}.$$

Since B_0 is complete, there exist $r_i = \lim_{m \rightarrow \infty} \sum_{j=0}^m \pi^j r_i^j$. Since V is complete, there exists $v = \lim_{m \rightarrow \infty} \sum_{j=0}^m \pi^j v_j$ and therefore $\varphi(v) = \lim_{m \rightarrow \infty} \sum_{j=0}^m \pi^j \varphi(v_j)$. Finally, since W is separated, we have

$$x - \sum_{i=1}^n r_i w_i - \varphi(v) \in \bigcap_{m \geq 1} \pi^m W = \{0\},$$

and therefore $x = \sum_{i=1}^n r_i w_i + \varphi(v)$. ■

Proof of Lemma 3.3. Recall that a module over a discrete valuation ring is flat if and only if it is torsion-free. In particular, images of $d^\bullet$ are also flat and we may invoke the Künneth formula [15, Thm. 3.6.1] to obtain exact sequences

$$0 \rightarrow H^i(C^\bullet) \otimes_{B_0} \ell \rightarrow H^i(C^\bullet \otimes_{B_0} \ell) \rightarrow \mathrm{Tor}_1^{B_0}(H^{i+1}(C^\bullet), \ell) \rightarrow 0 \quad (3.14)$$

To prove the first statement of the lemma we consider the exact sequences

$$C^{n-1} \xrightarrow{d^{n-1}} \ker d^n \rightarrow H^n(C^\bullet) \rightarrow 0. \quad (3.15)$$

By (3.14) and the initial assumptions, dimensions $\dim_\ell H^n(C) \otimes_{B_0} \ell$ are finite. Moreover, by assumption C^n are complete and thus $\ker d^n$ are separated modules as they are submodules of complete (hence separated) modules. Therefore we may apply Lemma 3.4 to sequences (3.15) to conclude the first part of the lemma.

For the second part, recall that it follows from the classification of finitely generated modules over discrete valuation rings that if M is such module then

$$\dim_\ell M \otimes_{B_0} \ell - \dim_B M \otimes_{B_0} B = \dim_k \mathrm{Tor}_1^{B_0}(M, \ell). \quad (3.16)$$

Since $- \otimes_{B_0} B$ is the same as localization at π , it is flat and we have

$$H^i(C^\bullet) \otimes_{B_0} B = H^i(C^\bullet \otimes_{B_0} B). \quad (3.17)$$

The first part of the lemma together with (3.14), (3.16), and (3.17) imply formula (3.13). First, observe that

$$\begin{aligned} \dim_\ell H^i(C^\bullet \otimes_{B_0} \ell) + \dim_B H^{i+1}(C^\bullet \otimes_{B_0} B) \\ = \dim_\ell H^i(C^\bullet) \otimes_{B_0} \ell - \dim_\ell H^{i+1}(C^\bullet) \otimes_{B_0} \ell. \end{aligned}$$

Now, let $N > 0$ be an integer such that $C^k = 0$ for $|k| \geq N$. Taking an alternating sum of the above equalities we obtain

$$\begin{aligned} \chi(C^\bullet \otimes_{B_0} \ell) - \chi(C^\bullet \otimes_{B_0} B) &= \sum_{i=-N}^{N-1} (\dim_\ell H^i(C^\bullet \otimes_{B_0} \ell) + \dim_B H^{i+1}(C^\bullet \otimes_{B_0} B)) \\ &= \dim_\ell H^{-N}(C^\bullet) \otimes_{B_0} \ell - \dim_\ell H^N(C^\bullet) \otimes_{B_0} \ell \\ &= 0. \end{aligned} \quad \blacksquare$$

4. Proof of Theorem 1.2

We now use lemmas proven in the previous section and some well-known properties of holonomic $\mathcal{D}$ -modules on affine spaces to prove Theorem 1.2.

Proof of Theorem 1.2. First we prove that $(1) \Rightarrow (2) \Rightarrow (3) \Rightarrow (1)$. We show condition (A) as a part of the second implication. Then we show (B).

$(1) \Rightarrow (2)$. This is the most tricky part of the proof. It suffices to prove that for any right $\widehat{\mathcal{D}}_n$ -module N of minimal dimension its dual $N^* = \text{Ext}_{\widehat{\mathcal{D}}_n}^n(N, \widehat{\mathcal{D}}_n)$ has a lattice with reduction of minimal dimension. Indeed, by Lemma 2.1 we then may take $N = M^*$ which is of minimal dimension and satisfies $N^* = M^{**} = M$. Let $V \subset N$ be some lattice (*a priori* with reduction that is possibly not of minimal dimension). By Lemma 3.1 applied to $B_0 = \widehat{\mathcal{D}}_n^\circ$ and $\pi = \varpi$ we have an inclusion

$$0 \rightarrow \bar{\mathcal{D}} \otimes_{\widehat{\mathcal{D}}_n^\circ} \text{Ext}_{\widehat{\mathcal{D}}_n^\circ}^d(V, \widehat{\mathcal{D}}_n^\circ) \rightarrow \text{Ext}_{\bar{\mathcal{D}}_n}^n(\bar{V}, \bar{\mathcal{D}}_n).$$

The key observation is that the module on the left is of minimal dimension. Indeed, since $\bar{\mathcal{D}}_n = \mathbb{W}_n(k)$ and $\bar{V}$ is finitely generated, by part (b) of Lemma 2.2 we know that the module on the right hand side is of minimal dimension. Therefore so is the module on the left hand side by part (a) of the same lemma. Now set

$$T = \{m \in \text{Ext}_{\widehat{\mathcal{D}}_n^\circ}^n(V, \widehat{\mathcal{D}}_n^\circ) : \varpi^k m = 0 \text{ for some } k\}.$$

This is a left $\widehat{\mathcal{D}}_n^\circ$ -module because ϖ is central in $\widehat{\mathcal{D}}_n^\circ$. We define L as the quotient of $\text{Ext}_{\widehat{\mathcal{D}}_0}^n(V, \widehat{\mathcal{D}}_0)$ by T , so that it fits into a short exact sequence.

$$0 \rightarrow T \rightarrow \text{Ext}_{\widehat{\mathcal{D}}_0}^n(V, \widehat{\mathcal{D}}_0) \rightarrow L \rightarrow 0. \quad (4.1)$$

We will show that L is the desired lattice, i.e., that

- (a) L is a finitely generated $\widehat{\mathcal{D}}_n^\circ$ -module.
- (b) $K \otimes_{\mathfrak{o}_K} L = N^*$ and the natural map $L \rightarrow N^*$ is injective.
- (c) $\bar{\mathcal{D}}_n \otimes_{\widehat{\mathcal{D}}_n^\circ} L$ has minimal dimension.

To show (a) recall from Lemma 2.3 that $\widehat{\mathcal{D}}_n^\circ$ is left and right noetherian. From noetherianity we conclude that because V is finitely generated so is $\text{Ext}_{\widehat{\mathcal{D}}_n^\circ}^n(V, \widehat{\mathcal{D}}_n^\circ)$, and therefore L is also finitely generated because by (4.1) it is a quotient of a finitely generated module. To show (b) we note that $K \otimes_{\mathfrak{o}_K} -$ coincides with the localization at ϖ . As $\widehat{\mathcal{D}}_n^\circ$ is noetherian and V is finitely generated, we have

$$K \otimes_{\mathfrak{o}_K} \text{Ext}_{\widehat{\mathcal{D}}_n^\circ}^n(V, \widehat{\mathcal{D}}_n^\circ) = \text{Ext}_{\widehat{\mathcal{D}}_n}^n(N, \widehat{\mathcal{D}}_n) = N^*. \quad (4.2)$$

Indeed, it is well known that for *commutative* noetherian rings localization commutes with Ext for finitely generated modules, and the standard proof of this fact [15, Prop. 3.3.10] works also in our case, because ϖ is central in $\widehat{\mathcal{D}}_n^\circ$. By construction $K \otimes_{\mathfrak{o}_K} T = 0$, so tensoring (4.1) with K we get from (4.2) that $K \otimes_{\mathfrak{o}_K} L = N^*$. The natural map $L \rightarrow M$ is injective by construction because its kernel consists precisely of ϖ -torsion of L . Finally, we show (c). Since tensoring is right exact, we have an exact sequence of left $\bar{\mathcal{D}}_n$ -modules

$$\bar{\mathcal{D}}_n \otimes_{\widehat{\mathcal{D}}_n^\circ} \text{Ext}_{\widehat{\mathcal{D}}_n^\circ}^n(V, \widehat{\mathcal{D}}_n^\circ) \rightarrow \bar{\mathcal{D}}_n \otimes_{\widehat{\mathcal{D}}_n^\circ} L \rightarrow 0.$$

Now, (c) follows from part (a) of Lemma 2.2 because the right hand side is a quotient of a $\bar{\mathcal{D}}_n$ -module which we have already observed to be of minimal dimension. This finishes the proof of (1) $\Rightarrow$ (2).

(2) $\Rightarrow$ (3). This is a consequence of Lemma 3.2. Indeed, it is known [2, Ch. 1, Prop. 5.3] that finitely generated $\bar{\mathcal{D}}_n$ -modules of minimal dimension are of finite length. It follows by induction from part (a) of Lemma 2.2 that a semisimplification of a $\bar{\mathcal{D}}_n$ -module of finite length has minimal dimension if and only if the module itself has minimal dimension. Therefore we may use Lemma 3.2 to get the desired implication. We also get (A) as a byproduct.

(3) $\Rightarrow$ (1). Let $L \subset M$ be a lattice such that $\bar{L}$ has minimal dimension. By the very definition we have $\text{Ext}_{\bar{\mathcal{D}}_n}^i(\bar{L}, \bar{\mathcal{D}}_n) = 0$ for $0 \leq i \leq n-1$. Then short exact sequences of Lemma 3.1 for $B_0 = \hat{\mathcal{D}}_n^\circ$ give

$$0 \rightarrow \text{Ext}_{\hat{\mathcal{D}}_n^\circ}^i(L, \hat{\mathcal{D}}_n^\circ) \otimes_{\hat{\mathcal{D}}_n^\circ} \bar{\mathcal{D}}_n \rightarrow \text{Ext}_{\bar{\mathcal{D}}_n}^i(\bar{L}, \bar{\mathcal{D}}_n) = 0,$$

i.e.,

$$\text{Ext}_{\hat{\mathcal{D}}_n^\circ}^i(L, \hat{\mathcal{D}}_n^\circ) \otimes_{\hat{\mathcal{D}}_n^\circ} \bar{\mathcal{D}}_n = 0 \quad (\text{for } i < n).$$

By noetherianity of $\hat{\mathcal{D}}_n^\circ$ (Lemma 2.3) we know that the right $\hat{\mathcal{D}}_n^\circ$ -modules $\text{Ext}_{\hat{\mathcal{D}}_n^\circ}^i(L, \hat{\mathcal{D}}_n^\circ)$ are finitely generated and therefore must be zero by Nakayama's lemma part of Lemma 2.5. As we have already explained while proving that (1) $\Rightarrow$ (2), we always have isomorphisms $\text{Ext}_{\hat{\mathcal{D}}_n}^i(M, \hat{\mathcal{D}}_n) = \text{Ext}_{\hat{\mathcal{D}}_n^\circ}^i(L, \hat{\mathcal{D}}_n^\circ) \otimes_{\mathfrak{o}_K} K$. We conclude that $\text{Ext}_{\hat{\mathcal{D}}_n}^i(M, \hat{\mathcal{D}}_n)$ must vanish for $i < n$, i.e., M has minimal dimension. This closes the circle of implications.

To prove (B) we use Lemma 3.3. Assume that equivalent conditions of Theorem 1.2 hold for M and let $L \subset M$ be a lattice which has a reduction of minimal dimension. Consider the complex

$$\text{DR}^\bullet(L) = L \rightarrow \bigoplus_{i=1}^d L \cdot dx_i \rightarrow \bigoplus_{i < j} L \cdot dx_i \wedge dx_j \rightarrow \cdots$$

with differentials as in (1.4). This is a bounded complex of complete (by Lemma 2.5) and torsion-free (since lattices are ϖ -torsion-free) $\mathfrak{o}_K$ -modules. Note that by construction we have

$$\text{DR}^\bullet(L) \otimes_{\mathfrak{o}_K} K = \text{DR}_{\hat{\mathcal{D}}_n}^\bullet(M)$$

and

$$\text{DR}^\bullet(L) \otimes_{\mathfrak{o}_K} k = \text{DR}_{\bar{\mathcal{D}}_n}^\bullet(\bar{L}).$$

The latter has finitely-dimensional cohomology over k by part (d) of Lemma 2.2. We may now apply Lemma 3.3 and conclude that $\dim_K H_{dR}^i(M) < \infty$ for all i and moreover $\chi_{dR}(M) = \chi_{dR}(\bar{L})$. ■

Acknowledgments. This work was part of my Ph.D. thesis. I thank my doctoral advisors: Piotr Achinger and Adrian Langer for numerous discussions and helpful suggestions. I thank the referees of my thesis: Wojciech Gajda, Grzegorz Kapustka, and Jérôme

Poineau, as well as the anonymous referees of this manuscript for valuable comments. I thank Stefano Aloé, Javier Fresán, and Gabriel Ribeiro for their hospitality during my stay in Palaiseau.

Funding. This work was supported by the project KAPIBARA funded by the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme (grant agreement No 802787). Part of this work was created during author’s visit at the École polytechnique in Palaiseau. This visit was supported by the “Inicjatywa Doskonałości-Uczelnia Badawcza” programme funded by the University of Warsaw.

References

- [1] M. F. Atiyah and I. G. Macdonald, *Introduction to commutative algebra*. Addison-Wesley Ser. Math., Westview Press, Boulder, CO, 2016 Zbl [1351.13002](#) MR [3525784](#)
- [2] J.-E. Björk, *Rings of differential operators*. North-Holland Mathematical Library 21, North-Holland Publishing, Amsterdam, 1979 Zbl [0499.13009](#) MR [0549189](#)
- [3] R. Hotta, K. Takeuchi, and T. Tanisaki, *D-modules, perverse sheaves, and representation theory*. Progr. Math. 236, Birkhäuser, Boston, MA, 2008 Zbl [1136.14009](#) MR [2357361](#)
- [4] N. M. Katz, *Exponential sums and differential equations*. Ann. of Math. Stud. 124, Princeton University Press, Princeton, NJ, 1990 Zbl [0731.14008](#) MR [1081536](#)
- [5] A. Langer, Moduli spaces of semistable modules over Lie algebroids. [v1] 2021, [v2] 2022, arXiv:[2107.03128v2](#)
- [6] J. C. McConnell, *I*-adic completions of noncommutative rings. *Israel J. Math.* **32** (1979), no. 4, 305–310 Zbl [0412.16009](#) MR [0571085](#)
- [7] Z. Mebkhout, *Le formalisme des six opérations de Grothendieck pour les $\mathcal{D}_X$ -modules cohérents*. Travaux en Cours 35, Hermann, Paris, 1989 Zbl [0686.14020](#) MR [1008245](#)
- [8] Z. Mebkhout and L. Narváez-Macarro, *La théorie du polynôme de Bernstein-Sato pour les algèbres de Tate et de Dwork-Monsky-Washnitzer*. *Ann. Sci. École Norm. Sup. (4)* **24** (1991), no. 2, 227–256 Zbl [0765.14009](#) MR [1097693](#)
- [9] L. Narváez Macarro, Division theorem over the Dwork-Monsky-Washnitzer completion of polynomial rings and Weyl algebras. In *Rings, Hopf algebras, and Brauer groups (Antwerp/Brussels, 1996)*, pp. 175–191, Lecture Notes in Pure and Appl. Math. 197, Dekker, New York, 1998 Zbl [0944.13018](#) MR [1615789](#)
- [10] A. Pangalos, *p*-adic Weyl algebras. Ph.D. thesis, Universität Münster, 2007, <https://d-nb.info/987679023/34> visited on 11 June 2026
- [11] F. Rączka, Holonomic $\mathcal{D}$ -modules on rigid analytic varieties. 2024, arXiv:[2405.03028v1](#)
- [12] J. J. Rotman, *An introduction to homological algebra*. 2nd edn., Universitext, Springer, New York, 2009 Zbl [1157.18001](#) MR [2455920](#)
- [13] L. H. Rowen, *Ring theory. Vol. I*. Pure Appl. Math. 127, Academic Press, Boston, MA, 1988 Zbl [0651.16001](#) MR [0940245](#)
- [14] P. Schneider, *Modular representation theory of finite groups*. Springer, Dordrecht, 2013 Zbl [1268.20012](#) MR [3012316](#)
- [15] C. A. Weibel, *An introduction to homological algebra*. Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge, 1994 Zbl [0797.18001](#) MR [1269324](#)

Communicated by Henning Krause

Received 10 June 2024; revised 23 February 2026.

Feliks Rączka

Department of Mathematics, Institute of Mathematics, Polish Academy of Sciences,
ul. Śniadeckich 8, 00-656 Warsaw; Institute of Mathematics, University of Warsaw, ul. Banacha 2,
02-097 Warsaw, Poland; fraczka@impan.pl, f.raczka@uw.edu.pl

Author IDs: zbMATH [raczka.feliks](#) MR [1452594](#) ORCID [0000-0002-1126-8382](#)