



Maciej Borodzik · Beibei Liu · Ian Zemke

Lattice homology, formality, and plumbed L-space links

Received November 23, 2022; revised September 27, 2023

Abstract. We define a link lattice complex for plumbed links, generalizing constructions of Ozsváth, Stipsicz and Szabó, and of Gorsky and Némethi. We prove that for all plumbed links in rational homology 3-spheres, the link lattice complex is homotopy equivalent to the link Floer complex as an A_∞ -module. Additionally, we prove that the link Floer complex of a plumbed L-space link is a free resolution of its homology. As a consequence, we give an algorithm to compute the link Floer complexes of plumbed L-space links, in particular of algebraic links, from their multivariable Alexander polynomial.

Keywords: link lattice homology, link Floer homology, plumbed L-space links.

Contents

1. Introduction	5274
1.1. Overview	5274
1.2. The link lattice complex	5275
1.3. Algebraic and plumbed L-space links	5276
1.4. Gorsky and Némethi's link lattice homology	5278
1.5. Structure of the paper	5279
2. Algebraic background	5280
2.1. A_∞ -modules	5280
2.2. The homological perturbation lemma	5281
2.3. Free resolutions and A_∞ -actions	5282
2.4. Example of non-formal chain complexes	5284
3. Plumbed manifolds and plumbed links	5285
3.1. Review of plumbed manifolds	5285
3.2. Plumbed manifolds and resolutions of analytic singularities	5287
3.3. Embedded resolutions	5288
3.4. Rationality	5290
3.5. Algebraic links and L-space links	5291

Maciej Borodzik: Institute of Mathematics, Polish Academy of Sciences, 00-656 Warszawa, Poland; mcboro@mimuw.edu.pl

Beibei Liu: Department of Mathematics, The Ohio State University, Columbus, OH 43210, USA; bbliumath@gmail.com

Ian Zemke: Department of Mathematics, University of Oregon, Eugene, OR 97403, USA; izemke@uoregon.edu

Mathematics Subject Classification 2020: 57K18 (primary); 14J17, 32S05, 57R58 (secondary).

4. Link lattice homology	5293
4.1. Background on link Floer homology	5293
4.2. Lattice homology	5295
4.3. The link lattice complex	5296
4.4. The module structure	5296
4.5. Maslov gradings	5299
4.6. Alexander gradings	5300
4.7. Conjugation symmetry	5301
4.8. Spin^c structures	5303
4.9. Independence from the framing on arrow components	5303
4.10. Freeness of the lattice complex	5305
4.11. Type- D modules over $\mathcal{K}$	5307
4.12. An example	5309
5. The equivalence with link Floer homology	5311
5.1. The link surgery complex and sublinks	5312
5.2. Link lattice homology and the link surgery formula	5313
5.3. Proof of Theorem 5.1	5317
5.4. Absolute gradings	5320
6. Plumbed L-space links	5323
6.1. Plumbed L-space links and free resolutions	5323
6.2. Computing the Floer chain complex from Alexander polynomials	5324
6.3. Comparison with Gorsky and Némethi's link lattice homology	5328
7. Computations	5330
7.1. Generators and relations for the homology of an L-space link	5331
7.2. The type- A module of $T(n, n)$	5333
7.3. The free complex of $T(3, 3)$	5336
7.4. The free complex of $T(4, 4)$	5337
References	5339

1. Introduction

1.1. Overview

Lattice homology is a combinatorial invariant of plumbed 3-manifolds defined by Némethi [33, 34]; see also [36, Chapter 11]. Némethi's construction is a formalization of earlier work of Ozsváth and Szabó [43], which computes the Heegaard Floer homology groups of many plumbed 3-manifolds. If Y is a plumbed 3-manifold, we write $\mathbb{H}\mathbb{F}(Y)$ for its lattice homology, which is a module over the power series ring $\mathbb{F}[[U]]$.

Work of Némethi, Ozsváth, Stipsicz and Szabó [33, 34, 41, 43] proves that

$$\mathbb{H}\mathbb{F}(Y) \cong \mathbf{HF}^-(Y)$$

for many important families of plumbed 3-manifolds. Here, $\mathbf{HF}^-(Y)$ denotes the Heegaard Floer homology over $\mathbb{F}[[U]]$. More recently, the third author proved the isomorphism in general [59].

Given a knot K in S^3 , Ozsváth–Szabó [44] and Rasmussen [53] defined a refinement of Heegaard Floer homology, called knot Floer homology. There are several equivalent formulations of this invariant. For our purposes, it is most convenient to consider knot

Floer homology as taking the form of a free, finitely generated chain complex $\mathcal{CFK}(K)$ over the 2-variable polynomial ring $\mathbb{F}[\mathcal{U}, \mathcal{V}]$.

Ozsváth and Szabó also defined a version of Heegaard Floer theory for links in 3-manifolds [48]. For a link $L \subseteq S^3$, we will focus on the description of the link Floer complex as a finitely generated free chain complex $\mathcal{CFL}(L)$ over the polynomial ring $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$, where $\ell = |L|$. For our purposes, it is also helpful to consider the link Floer complexes over the completed ring $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]]$, denoted

$$\mathcal{CFL}(L) := \mathcal{CFL}(L) \otimes_{\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]} \mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]].$$

Their construction also applies more generally when L is a link in a rational homology 3-sphere Y . In this case, we denote the link Floer complex by $\mathcal{CFL}(Y, L)$.

A relative version of lattice homology for plumbed knots was defined by Ozsváth, Stipsicz and Szabó [40]. Modulo notational differences, their version of knot lattice homology is analogous to the complex $\mathcal{CFK}(K)$. They proved that for a plumbed L-space knot in S^3 , knot lattice homology coincides with the knot Floer complex $\mathcal{CFK}(K)$.

Gorsky and Némethi [11] defined a relative version of lattice homology for L-space links. (Note that their construction does not require the link to be plumbed.) They constructed a spectral sequence from their version of link lattice homology to a version of link Floer homology, which is the homology of the quotient complex $\mathcal{CFL}(L)/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$, and proved that it degenerates for all algebraic links. In particular, their version of link lattice homology is isomorphic (as a graded group) to the version of link Floer homology for algebraic links in S^3 .

In this paper, we construct a new version of link lattice homology. Our link lattice complex is more closely related to the construction of Ozsváth, Stipsicz and Szabó [40], and is modeled on the full link Floer complex $\mathcal{CFL}(L)$. We use this link lattice complex to study plumbed L-space links, a family which includes all algebraic links in S^3 .

1.2. The link lattice complex

Suppose that L is a plumbed link in a plumbed 3-manifold Y . Such a pair (Y, L) is represented by a weighted graph Γ , whose vertices are partitioned into two sets,

$$V_\Gamma = V_G \cup V_\uparrow.$$

The vertices V_G are equipped with integral weights. The vertices in $V_\uparrow$ have no weights, and we refer to them as *arrow* vertices. Unless explicitly specified otherwise, we will always assume that Γ is a tree.

From the tree Γ , we obtain a partitioned link $L_\Gamma = L_G \cup L_\uparrow$ in S^3 . This link may be described as a connected sum of Hopf links, with one unknotted component for each vertex of Γ , and one clasp for each edge of Γ . The manifold Y is the result of surgery on L_G , with framing Λ obtained from the weights. Inside of $Y \cong S^3_\Lambda(L_G)$, the link L is identified with $L_\uparrow$.

Given a plumbing tree Γ representing a plumbed link (Y, L) , we will construct a chain complex

$$\mathbb{C}\mathrm{FL}(\Gamma, V_{\uparrow}).$$

Given an orientation of $L_{\uparrow}$, we will equip the chain complex $\mathbb{C}\mathrm{FL}(\Gamma, V_{\uparrow})$ with the structure of a module over the ring $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_{\ell}, \mathcal{V}_{\ell}]]$, where $\ell = |V_{\uparrow}|$, as well as with a Maslov grading and a $\mathbb{Q}^{\ell}$ -valued Alexander grading.

It is helpful to view $\mathbb{C}\mathrm{FL}(\Gamma, V_{\uparrow})$ as an A_{∞} -module over $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_{\ell}, \mathcal{V}_{\ell}]]$ with only m_1 and m_2 non-vanishing. We note that the complex $\mathbb{C}\mathrm{FL}(\Gamma, V_{\uparrow})$ is not free over this ring unless $|V_{\uparrow}| = 1$.

A central result of the paper is the following:

Theorem 1.1. *Suppose that Γ is a plumbing link diagram which is a tree, and write (Y, L) for the associated 3-manifold and link. If Y is a rational homology sphere, then $\mathcal{C}\mathcal{F}\mathcal{L}(Y, L)$ is homotopy equivalent to $\mathbb{C}\mathrm{FL}(\Gamma, V_{\uparrow})$ as an absolutely graded A_{∞} -module over $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_{\ell}, \mathcal{V}_{\ell}]]$.*

See Theorem 5.1 for further details. In the above, we write $\mathcal{C}\mathcal{F}\mathcal{L}(Y, L)$ for the full link Floer complex $\mathcal{C}\mathcal{F}\mathcal{L}(Y, L)$ completed over the power series ring $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_{\ell}, \mathcal{V}_{\ell}]]$.

1.3. Algebraic and plumbed L -space links

We recall that a rational homology 3-sphere Y is called an L -space if

$$\widehat{HF}(Y, \mathfrak{s}) \cong \mathbb{F}$$

for each $\mathfrak{s} \in \mathrm{Spin}^c(Y)$. A link $L \subseteq S^3$ is called an L -space link if all sufficiently positive surgeries are L -spaces. We note that since Dehn surgery does not depend on the orientation of the link, the property of being an L -space link is independent of orientations.

An important family of plumbed links in S^3 is formed by the algebraic links, which are the links of complex plane curve singularities. According to Gorsky and Némethi [12], algebraic links in S^3 are L -space links.

There is a useful characterization of L -space links in terms of the link Floer complex. If $L \subseteq S^3$, then L is an L -space link if and only if the homology group $\mathcal{H}\mathcal{F}\mathcal{L}(L)$ is torsion-free as an $\mathbb{F}[U]$ -module, where U acts by $\mathcal{U}_i \mathcal{V}_i$ for any i . (Since $\mathcal{U}_i \mathcal{V}_i$ and $\mathcal{U}_j \mathcal{V}_j$ have chain homotopic actions for all i and j , the definition is independent of the choice of i .)

Ozsváth and Szabó [47] proved the knot Floer complex of an L -space knot is a staircase complex. A very natural question is whether an analog of that result holds for L -space links. L -space links have been extensively studied in the literature: see e.g. [2, 4, 9–11, 26–28]. Despite the interest in L -space links and many interesting results concerning them, there is to date no result which characterizes the structure of the link Floer complex of an L -space link in parallel with Ozsváth and Szabó's result for L -space knots.

We prove the following:

Theorem 1.2. *Suppose that $L \subseteq S^3$ is a plumbed L -space link. Then the link Floer complex $\mathcal{C}\mathcal{F}\mathcal{L}(L)$ is homotopy equivalent to a free resolution of its homology over the ring*

$\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$. Equivalently, the complex $\mathcal{CFL}(L)$ is homotopy equivalent to its homology $\mathcal{HFL}(L)$ as an A_∞ -module over $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$, where we equip $\mathcal{HFL}(L)$ with the A_∞ -module structure which has $m_j = 0$ unless $j = 2$.




Fig. 1.1. The knot and link Floer complexes of $T(3, 4)$, $T(2, 2)$ and $T(2, 4)$. Each dot denotes a generator in a free basis. The horizontal direction indicates the grading of the free resolution.

Remark 1.3. (1) Theorem 1.2 is a natural generalization of the result of Ozsváth and Szabó for L-space knots because the staircase complex of an L-space knot is easily seen to be a free resolution of its homology over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$.

(2) Theorem 1.2 also holds more generally for plumbed L-space links L in plumbed 3-manifolds Y which are themselves L-spaces. For details, see Section 6.

(3) A dg -module M over a ring A such that $(H_*(M), m_2)$ is homotopy equivalent to M as an A_∞ -module over A is called *formal*. Hence, we may restate Theorem 1.2 by saying that plumbed L-space links have formal link Floer complexes.

We do not know whether Theorem 1.2 holds for non-plumbed L-space links. We state the following open question:

Question 1.4. Are there non-plumbed L-space links $L \subseteq S^3$ for which $\mathcal{CFL}(L)$ is not homotopy equivalent to a free resolution of $\mathcal{HFL}(L)$?

From Theorem 1.2, we obtain an algorithm to compute the link Floer complex of a plumbed L-space link. Namely, we observe that for an L-space link L , the $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$ -module $\mathcal{HFL}(L)$ is completely determined by the H -function

(or equivalently the d -invariants of large surgeries). According to [11], the H -function of an L-space link in S^3 is determined by the multivariable Alexander polynomials of L and its sublinks. After determining $\mathcal{HFL}(L)$, one may compute $\mathcal{CFL}(L)$ by computing its free resolution. Finding such a resolution is algorithmic (see e.g. [51]) and may be done using computer algebra software such as Macaulay2 [14]. We carry this out for $T(3, 3)$ and $T(4, 4)$ in Section 7.

Our algorithm to compute the full link Floer complexes enables us to compute the minus and the hat version of link Floer homology of plumbed L-space links by setting respectively $\mathcal{V}_i = 0$ and $\mathcal{U}_i = \mathcal{V}_i = 0$ for all i . Moreover, the hat version of link Floer homology detects the Thurston norm of the link complement in the 3-sphere [49]. Hence, the algorithm also gives a way to compute the Thurston norm of plumbed L-space links in S^3 .

Corollary 1.5. *The link Floer complex of a plumbed L-space link L in S^3 is computable from the multivariable Alexander polynomials of the link L and its sublinks.*

Remark 1.6. Computing free resolutions is in general a challenging task. For many purposes (e.g. taking tensor products), it is more practical to understand the homology group $\mathcal{HFL}(L)$ and use the fact that $\mathcal{CFL}(L)$ is homotopy equivalent as an A_∞ -module to $\mathcal{HFL}(L)$ with only m_2 non-trivial. Often the homology group $\mathcal{HFL}(L)$ has a much more concise description than its free resolution. For example in Section 7, we give a simple description of the module $\mathcal{HFL}(T(n, n))$. We describe free resolutions for $T(3, 3)$ and $T(4, 4)$, which are considerably more complex.

We note that for algebraic links in S^3 (which are particular examples of plumbed L-space links) the fact that the link Floer complex is determined by the Alexander polynomial may be seen indirectly using the fact that the Alexander polynomial of an algebraic link determines the link (see Yamamoto [55]). This is not the case for algebraic links in other 3-manifolds, or other plumbed L-space links in S^3 ; see [3] and also Proposition 3.20. Nonetheless, our techniques give a concrete algorithm for computing the link Floer complex based on its Alexander polynomial. Although foundational, Yamamoto's work does not give a practical algorithm for computing link Floer complexes of algebraic links in S^3 .

1.4. Gorsky and Némethi's link lattice homology

We also consider Gorsky and Némethi's link lattice homology, which is defined for all L-space links. If L is an L-space link, then they described a chain complex $\mathcal{K}(L)$, which they called the link lattice complex. They proved that if L is an algebraic link, then

$$H_*(\mathcal{K}(L)) \cong H_*(\mathcal{CFL}(L)/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)), \quad (1.1)$$

as graded groups, where the right-hand side is identified with the minus version of link Floer homology.

In our paper, we give an alternate perspective on the complex $\mathcal{K}(L)$. Namely, we show that $\mathcal{K}(L)$ is homotopy equivalent as an A_∞ -module to the derived tensor product

$$\mathcal{HFL}(L) \tilde{\otimes}_{\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]} \mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]/(\mathcal{V}_1, \dots, \mathcal{V}_\ell),$$

where we equip $\mathcal{HFL}(L)$ with the A_∞ -module structure which has only m_2 non-trivial. As a corollary of Theorem 1.2, we obtain the following result:

Corollary 1.7. *If L is a plumbed L-space link, then Gorsky and Némethi's link lattice complex $\mathcal{K}(L)$ is homotopy equivalent to $\mathcal{CFL}(L)/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$ as a graded chain complex.*

We note Gorsky and Némethi only prove the isomorphism in (1.1) at the level of graded groups. Our proof of Corollary 1.7 improves on their result because it equips $\mathcal{K}(L)$ with a dg -module structure over $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$ (i.e. an A_∞ -module structure with only m_1 and m_2 non-trivial) and proves the isomorphism at the level of A_∞ -modules.

Gorsky and Némethi also constructed a spectral sequence from the tensor product $\mathcal{HFL}(L) \otimes \Lambda_{\mathcal{R}_\ell}^*(\xi_1, \dots, \xi_\ell)$ to the homology of $\mathcal{CFL}(L)/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$. Here, $\Lambda_{\mathcal{R}_\ell}^*(\xi_1, \dots, \xi_\ell)$ is the exterior algebra on ℓ generators and $\mathcal{R}_\ell = \mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$. Our Corollary 1.7 naturally recovers this spectral sequence, as follows. We first recall that the module $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$ has a free resolution over $\mathcal{R}_\ell$ equal to the Koszul complex $\Lambda_{\mathcal{R}_\ell}^*(\xi_1, \dots, \xi_\ell)$, where we equip each ξ_i with the differential $d(\xi_i) = \mathcal{V}_i$ (extended via the Leibniz rule). The chain complex $(\Lambda_{\mathcal{R}_\ell}^*(\xi_1, \dots, \xi_\ell), d)$ has a cube filtration, and the spectral sequence associated to the induced cube filtration on $\mathcal{HFL}(L) \otimes \Lambda_{\mathcal{R}_\ell}^*(\xi_1, \dots, \xi_\ell)$ coincides with Gorsky and Némethi's spectral sequence.

1.5. Structure of the paper

Section 2 describes background on A_∞ -modules and the homological perturbation lemma. We give a basic example from knot Floer theory of two complexes with isomorphic homology groups, such that one of them is chain homotopy equivalent to its homology (regarded as a chain complex with trivial differentials), while the other one is not.

Next, we recall some topological background in Section 3. We provide the definitions of plumbed manifolds and plumbed links. We state the result of Gorsky and Némethi that a plumbed link in a rational analytic singularity is an L-space link [12]. In particular, we show that plumbed L-space links are a natural generalization of algebraic links in S^3 . In Section 4, we define our link lattice complex. We describe the gradings and the $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$ -module structure.

Section 5 proves the equivalence of the link lattice complex and the link Floer complex, stated above in Theorem 1.1. This is the main technical result of the paper.

In Section 6 we focus on plumbed L-space links. Using Theorem 5.1 and the homological perturbation lemma, we prove Theorem 1.2, which states that the link Floer complex of a plumbed L-space link is a free resolution of its homology. As a consequence, we

show that the link Floer complex of a plumbed L-space link is computable from its multivariable Alexander polynomial; see Theorem 6.6. Additionally, we describe how our link lattice complex recovers the theory described by Gorsky and Némethi in the case of plumbed L-space links; see Theorem 6.10.

Section 7 describes some algorithms and examples. We provide a concrete way of presenting the $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_n, \mathcal{V}_n]$ -module $\mathcal{HFL}(L)$ for an L-space link L , from its H -function. The algorithm of Lemma 7.1 provides a representation of $\mathcal{HFL}(T(n, n))$ compatible with the description of Gorsky and Hom [9, Section 5]. Next, we compute $\mathcal{CFL}(T(3, 3))$ and $\mathcal{CFL}(T(4, 4))$ by constructing explicit free resolutions.

2. Algebraic background

In this section we recall the notion of an A_∞ -module (Section 2.1). Then, in Section 2.2, we state the homological perturbation lemma in A_∞ -category. In Section 2.3, we consider the homological perturbation lemma in the context of free resolutions of modules. In Section 2.4, we present two complexes over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ with isomorphic homology, but which are not chain homotopy equivalent. These two complexes appear naturally in knot Floer homology.

2.1. A_∞ -modules

Throughout the paper, we make use of the category of A_∞ -modules. The motivation is that A_∞ -module structures may be transferred along homotopy equivalences of groups. Suppose that $\mathcal{A}$ is a ring that is an algebra over a ring $\mathbf{k}$ of characteristic 2. Given a finitely generated chain complex C over $\mathcal{A}$, in very general circumstances, one may pick a homotopy equivalence over $\mathbf{k}$ between C and $H_*(C)$. Given such a homotopy equivalence, the homotopy transfer lemma (see [19]) equips $H_*(C)$ with the structure of an A_∞ -module over $\mathcal{A}$, such that C and $H_*(C)$ are homotopy equivalent as A_∞ -modules. Note that unless $\mathcal{A}$ is a field, it is rarely the case that C and $H_*(C)$ are homotopy equivalent as dg -modules over $\mathcal{A}$. When the homotopy equivalence between C and $H_*(C)$ is suitably simple, the A_∞ -module maps on $H_*(C)$ are usually computed using a version of the homological perturbation lemma, stated in Lemma 2.3.

We now recall the basics of A_∞ -modules. We mostly follow the notation of Lipshitz, Ozsváth and Thurston [24, 25].

Let $\mathcal{A}$ be an associative algebra with unit over a ring $\mathbf{k}$ of characteristic 2. We will frequently consider $\mathbf{k} = \mathbb{F}_2$. We write μ_2 for the multiplication on $\mathcal{A}$.

Definition 2.1. A left A_∞ -module ${}_{\mathcal{A}}M$ over $\mathcal{A}$ is a left $\mathbf{k}$ -module M equipped with $\mathbf{k}$ -module maps

$$m_{j+1}: \mathcal{A}^{\otimes j} \otimes_{\mathbf{k}} M \rightarrow M, \quad j \geq 0,$$

such that $m_1 \circ m_1 = 0$, and for each n and any $a_1, \dots, a_n \in \mathcal{A}$ and $\mathbf{x} \in M$, the following holds:

$$\sum_{i=0}^n m_{n-i+1}(a_n, a_{n-1}, \dots, a_{i+1}, m_{i+1}(a_i, \dots, a_1, \mathbf{x})) + \sum_{k=1}^{n-1} m_n(a_n, a_{n-1}, \dots, \mu_2(a_{k+1}, a_k), \dots, a_1, \mathbf{x}) = 0.$$

Lipshitz, Ozsváth and Thurston refer to A_∞ -modules as *type-A* modules, in contrast to *type-D* modules, which we now introduce.

Definition 2.2. A right *type-D* module $N^{\mathcal{A}}$ over $\mathcal{A}$ is a right $\mathbf{k}$ -module N , together with a $\mathbf{k}$ -linear structure map

$$\delta^1: N \rightarrow N \otimes_{\mathbf{k}} \mathcal{A}$$

such that

$$(\mathrm{id}_N \otimes \mu_2) \circ (\delta^1 \otimes \mathrm{id}_{\mathcal{A}}) \circ \delta^1 = 0.$$

2.2. The homological perturbation lemma

It is a general fact that A_∞ -algebra structures may be transferred along homotopy equivalences of the chain complex underlying an A_∞ -algebra. This was proved by Kadeishvili [19]. Homological perturbation theory gives concrete formulas for the resulting A_∞ -module structure under certain restrictions on the chain homotopy equivalence (see [21, Theorem 3]). An exposition of the technique may be found in the Ph.D. thesis of Lefèvre-Hasegawa [23].

Lemma 2.3. Suppose that $\mathcal{A}$ is an associative algebra over a ground ring $\mathbf{k}$, ${}_{\mathcal{A}}M = (M, m_j)$ is an A_∞ -module over $\mathcal{A}$, (Z, ∂) is a chain complex over $\mathbf{k}$, and that we have three maps of left $\mathbf{k}$ -modules

$$i: Z \rightarrow M, \quad \pi: M \rightarrow Z \quad \text{and} \quad h: M \rightarrow M$$

satisfying the following:

- (1) i and π are chain maps.
- (2) $\pi \circ i = \mathrm{id}_Z$.
- (3) $i \circ \pi = \mathrm{id}_M + \partial(h)$, where $\partial(h) := m_1 \circ h + h \circ m_1$.
- (4) $h \circ i = 0$.
- (5) $\pi \circ h = 0$.
- (6) $h \circ h = 0$.

Then there are A_∞ -module structure maps m_j^Z on Z , as well as A_∞ -module morphisms

$$I_*: {}_{\mathcal{A}}Z \rightarrow {}_{\mathcal{A}}M, \quad \Pi_*: {}_{\mathcal{A}}M \rightarrow {}_{\mathcal{A}}Z \quad \text{and} \quad H_*: {}_{\mathcal{A}}M \rightarrow {}_{\mathcal{A}}M$$

satisfying $m_1^Z = \partial$, $I_1 = i$, $\Pi_1 = \pi$ and $H_1 = h$, and such that the analogs of relations (1)–(6) are also satisfied by the A_∞ -module morphisms I_* , Π_* and H_* .

Remark 2.4. It is important to note that the maps i, π, h in the assumption of the homological perturbation lemma are required to be only $\mathbf{k}$ -module maps, and not necessarily $\mathcal{A}$ -module maps. We refer the interested reader to [23, Section 1.4] for a detailed proof.

The extended A_∞ -module maps in the homological perturbation lemma have a concrete description below. The structure maps on Z are given by the diagrams shown in Figure 2.1. Therein $m_{>1}$ denote the A_∞ structure maps of M , with m_1 excluded, and Δ is the comultiplication on the tensor algebra $T^*\mathcal{A}$.




Fig. 2.1. The maps appearing in the homological perturbation lemma for A_∞ -modules. The notation is introduced in [25, Section 2]. Briefly, a single arrow represents an element of $\mathcal{A}M$, while a double arrow represents an element of $\bigoplus_i \mathcal{A}^{\otimes i}$.

2.3. Free resolutions and A_∞ -actions

In this section, we describe a useful relation between free resolutions and A_∞ -module structures. We assume that $\mathcal{A}$ is an algebra over $\mathbb{F} = \mathbb{Z}/2$. Suppose that $(\mathcal{A}M, m_j)$ is a type- A module which has $m_j = 0$ for $j \neq 2$. That is, M is a left $\mathcal{A}$ -module in the ordinary sense. A *free resolution* of M is a collection $(F_i, f_i)_{i \in \mathbb{N}}$ of free $\mathcal{A}$ -modules and $\mathcal{A}$ -linear maps which form an exact sequence of the following form:

$$\cdots \rightarrow F_i \xrightarrow{f_i} F_{i-1} \xrightarrow{f_{i-1}} \cdots \xrightarrow{f_2} F_1 \xrightarrow{f_1} F_0 \xrightarrow{f_0} M \rightarrow 0.$$

Write $\mathcal{F}$ for the chain complex whose underlying space is the direct sum of the F_i , $i \geq 0$, and whose differential consists of the maps f_i for $i \geq 1$.

By definition, a free resolution ${}_{\mathcal{A}}\mathcal{F}$ is quasi-isomorphic to ${}_{\mathcal{A}}M$, since the canonical projection map from $\mathcal{F}$ to M is a chain map which induces an isomorphism on

homology. In the category of A_∞ -modules, quasi-isomorphisms are always invertible as A_∞ -morphisms. An exposition of this principle may be found in [20, Section 4]. In the present case, it is also helpful to construct the homotopy equivalence explicitly, since we will use it and similar homotopy equivalences later.

Proposition 2.5. *Suppose that ${}_{\mathcal{A}}\mathcal{F}$ is a free resolution of a left $\mathcal{A}$ -module ${}_{\mathcal{A}}M$. Then ${}_{\mathcal{A}}\mathcal{F}$ and ${}_{\mathcal{A}}M$ are homotopy equivalent as A_∞ -modules over $\mathcal{A}$.*

Proof. There is a canonical projection map $\pi: \mathcal{F} \rightarrow M$, which is just f_0 on F_0 , and 0 on every other summand. We pick any map of $\mathbb{F}$ -vector spaces $i: M \rightarrow F_0$ such that $f_0 \circ i = \text{id}_M$. The map i induces a direct sum splitting (of $\mathbb{F}$ -vector spaces) $F_0 = \text{im } f_1 \oplus \text{im } i$. The map $f_1: F_1 \rightarrow \text{im } f_1$ is surjective, so we pick its $\mathbb{F}$ -linear section, which we denote by h_0 . The map h_0 induces a splitting of F_1 into $\text{im } h_0 \oplus \text{im } f_2$. We define a map $h_1: \text{im } f_2 \rightarrow F_2$ to be a section of the map f_2 . We proceed in this manner to split the entire free resolution to obtain a diagram

$$\begin{array}{ccccccc} \cdots & \rightarrow & \text{im } f_{i+1} \oplus \text{im } h_{i-1} & \xrightarrow{f_i} & \text{im } f_i \oplus \text{im } h_{i-2} & \xrightarrow{f_{i-1}} & \cdots \\ & & \xleftarrow{h_{i-1}} & & \xleftarrow{h_{i-2}} & & \\ & & & & \xleftarrow{h_1} & & \\ & & & & & & \xleftarrow{h_0} \end{array} \quad \begin{array}{c} \xrightarrow{f_2} \\ \xrightarrow{f_1} \\ \xrightarrow{f_0} \end{array} \quad \begin{array}{c} \text{im } f_2 \oplus \text{im } h_0 \\ \text{im } f_1 \oplus \text{im } i \\ M \end{array} \rightarrow 0.$$

Clearly the maps i and π are chain maps. Furthermore, $\pi \circ i = \text{id}_M$, and

$$i \circ \pi = \text{id}_{\mathcal{F}} + [\partial, h],$$

where h is the direct sum of the h_i . Additionally,

$$h \circ i = 0, \quad h \circ h = 0 \quad \text{and} \quad \pi \circ h = 0.$$

In particular, the homological perturbation lemma induces an A_∞ -module structure on M , which we denote by ${}_{\mathcal{A}}\tilde{M}$, such that ${}_{\mathcal{A}}\tilde{M}$ and ${}_{\mathcal{A}}\mathcal{F}$ are homotopy equivalent as A_∞ -modules. We claim that the higher actions on ${}_{\mathcal{A}}\tilde{M}$ vanish. Indeed, the map h always maps F_i to F_{i+1} , while $m_2^{\mathcal{F}}$ preserves F_i , and π is only non-vanishing on F_0 . A quick inspection of the leftmost map in Figure 2.1 shows that $m_j = 0$ for $j > 2$.

From the claim it follows that ${}_{\mathcal{A}}\tilde{M} = {}_{\mathcal{A}}M$, completing the proof. $\blacksquare$

Recall that we say an A_∞ -module ${}_{\mathcal{A}}M$ is *formal* if ${}_{\mathcal{A}}M$ is homotopy equivalent to ${}_{\mathcal{A}}H_*(M)$ (viewed as an A_∞ -module with $m_j = 0$ for $j \neq 2$). The following is a helpful restatement of Proposition 2.5:

Corollary 2.6. *Suppose $\mathcal{A}$ is an algebra over $\mathbb{F} = \mathbb{Z}/2$. Let ${}_{\mathcal{A}}M$ be an A_∞ -module over $\mathcal{A}$, and let $\mathcal{F}$ be the total complex of a free resolution of $H_*(M)$. Then ${}_{\mathcal{A}}M$ is homotopy equivalent to ${}_{\mathcal{A}}\mathcal{F}$ as an A_∞ -module if and only if ${}_{\mathcal{A}}M$ is formal.*

We now describe a very helpful criterion for formality, which generalizes Corollary 2.6.

Lemma 2.7. *Suppose that $\mathcal{A}$ is an algebra over $\mathbb{F} = \mathbb{Z}/2$ and*

$${}_{\mathcal{A}}\mathcal{C} = (\cdots \xrightarrow{\partial_{i+1}} C_i \xrightarrow{\partial_i} C_{i-1} \xrightarrow{\partial_{i-1}} \cdots)$$

is a (possibly unbounded) chain complex of $\mathcal{A}$ -modules. That is, ${}_{\mathcal{A}}\mathcal{C}$ is an A_∞ -module with only m_1 and m_2 non-trivial, such that furthermore the action m_2 preserves C_i , and the differential m_1 on ${}_{\mathcal{A}}\mathcal{C}$ is given by the maps $\partial_i: C_i \rightarrow C_{i-1}$. We refer to the subscript i as the algebraic grading. If the homology of ${}_{\mathcal{A}}\mathcal{C}$ is supported in a single algebraic grading, then ${}_{\mathcal{A}}\mathcal{C}$ is formal.

Proof. We consider first the case that there is some $n \in \mathbb{Z}$ such that $C_i = 0$ for $i < n$ and such that the homology is supported in grading n , i.e. $H_*(\mathcal{C}) = C_n/\text{im}(\partial_{n+1})$. We can then define a chain map π of $\mathcal{A}$ -modules from ${}_{\mathcal{A}}\mathcal{C}$ to $H_*(\mathcal{C})$ by sending C_n to $C_n/\text{im}(\partial_{n+1})$ and sending C_i to 0 for $i \neq n$. This gives a quasi-isomorphism of ${}_{\mathcal{A}}\mathcal{C}$ onto ${}_{\mathcal{A}}H_*(\mathcal{C})$ (viewing this homology group as an A_∞ -module with $m_j = 0$ if $j \neq 2$). Since quasi-isomorphisms of A_∞ -modules are homotopy equivalences, Corollary 2.6 implies that ${}_{\mathcal{A}}\mathcal{C}$ is formal in this restricted case.

We now consider the general case that $H_*(\mathcal{C})$ is supported in a single grading n , i.e. $H_*(\mathcal{C}) = \ker(\partial_n)/\text{im}(\partial_{n+1})$. Write $Z_n \subseteq C_n$ for the kernel of ∂_n . Note that Z_n is an $\mathcal{A}$ -submodule. We consider the chain complex

$${}_{\mathcal{A}}\mathcal{C}' = (\cdots \xrightarrow{\partial_{n+2}} C_{n+1} \xrightarrow{\partial_{n+1}} Z_n).$$

Note that there is a natural inclusion map of ${}_{\mathcal{A}}\mathcal{C}'$ into ${}_{\mathcal{A}}\mathcal{C}$, which is a quasi-isomorphism. The complex ${}_{\mathcal{A}}\mathcal{C}'$ is formal by the previous argument. Therefore ${}_{\mathcal{A}}\mathcal{C}$ is as well. $\blacksquare$

2.4. Example of non-formal chain complexes

As mentioned in Section 2.1, if $\mathcal{A}$ is a field, then any finitely generated chain complex is homotopy equivalent to its homology. If $\mathcal{A}$ is a PID, then it is not hard to show that any finitely generated free chain complex is quasi-isomorphic to its homology. For general rings, this is not always the case. In this section, we illustrate the case $\mathcal{A} = \mathbb{F}[\mathcal{U}, \mathcal{V}]$ with examples from the theory of knot Floer homology.

We consider two free $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ -complexes C and D with generators $\mathbf{u}, \mathbf{v}, \mathbf{x}, \mathbf{y}, \mathbf{z}$, respectively $\mathbf{a}, \mathbf{b}$, and $\mathbf{c}$, and with differential represented by arrows as shown below:

$$C = \mathbf{u} \oplus \begin{array}{ccc} \mathbf{w} & \xrightarrow{\mathcal{U}} & \mathbf{x} \\ \downarrow \mathcal{V} & & \downarrow \mathcal{V} \\ \mathbf{y} & \xrightarrow{\mathcal{U}} & \mathbf{z} \end{array} \quad \text{and} \quad D = \begin{array}{ccc} & & \mathbf{a} \\ & & \downarrow \mathcal{V} \\ \mathbf{c} & \xrightarrow{\mathcal{U}} & \mathbf{b} \end{array}$$

For appropriate choices of gradings, there is an isomorphism $H_*(C) \cong H_*(D)$, since both are isomorphic to $\mathbb{F}[\mathcal{U}, \mathcal{V}] \oplus \mathbb{F}$, where $\mathbb{F}$ has vanishing action of $\mathcal{U}$ and $\mathcal{V}$. On the other hand, it is easy to see that C and D are not homotopy equivalent over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$. For example, if we tensor both with the module $\mathbb{F}[\mathcal{U}, \mathcal{V}]/(\mathcal{U}, \mathcal{V})$ and take homology, we obtain vector spaces of different rank over $\mathbb{F}$.

We now equip both $H_*(C)$ and $H_*(D)$ with A_∞ -actions by applying the homological perturbation lemma. Since C is a free resolution of its homology, the induced A_∞ -action has only m_2 non-trivial.

For D , the homology is the $\mathbb{F}$ -span of $\mathcal{U}^i \mathcal{V}^j (\mathcal{U}\mathbf{a} + \mathcal{V}\mathbf{c})$ and $\mathbf{b}$, for all $i, j \in \mathbb{N}$. We may define a homotopy equivalence of chain complexes over $\mathbb{F}$ between D and the complex $\mathbb{F}[\mathcal{U}, \mathcal{V}] \oplus \mathbb{F}$ (with vanishing differential). Write $\mathbf{e}$ for the generator of $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ and $\mathbf{f}$ for the generator of $\mathbb{F}$. The inclusion map $i: H_*(D) \rightarrow D$ is given by

$$i(\mathcal{U}^n \mathcal{V}^m \mathbf{e}) = \mathcal{U}^n \mathcal{V}^m (\mathcal{U}\mathbf{a} + \mathcal{V}\mathbf{c}), \quad i(\mathbf{f}) = \mathbf{b},$$

for all $n, m \geq 0$. We define a projection map $\pi: D \rightarrow H_*(D)$ by setting

$$\pi(\mathcal{U}^n \mathcal{V}^m \mathbf{a}) = \begin{cases} \mathcal{U}^{n-1} \mathcal{V}^m \mathbf{e} & \text{if } n > 0, \\ 0 & \text{otherwise,} \end{cases} \quad \pi(\mathcal{U}^n \mathcal{V}^m \mathbf{b}) = \begin{cases} \mathbf{f} & \text{if } n = m = 0, \\ 0 & \text{otherwise.} \end{cases}$$

The map π vanishes on multiples of $\mathbf{c}$. The maps π and i are clearly chain maps, and $\pi \circ i = \text{id}$. We define a homotopy $h: D \rightarrow D$ by setting $h(\mathbf{a}) = h(\mathbf{c}) = 0$, and

$$h(\mathcal{U}^n \mathcal{V}^m \mathbf{b}) = \begin{cases} \mathcal{U}^{n-1} \mathcal{V}^m \mathbf{c} & \text{if } n > 0, \\ \mathcal{V}^{m-1} \mathbf{a} & \text{if } n = 0, m > 0, \\ 0 & \text{otherwise.} \end{cases}$$

It is straightforward to see that $i \circ \pi = \text{id} + [\partial, h]$. Furthermore, $h \circ i$, $h \circ h$ and $\pi \circ h$ vanish. In particular, the maps i , π and h induce an A_∞ -module structure maps $m_j^{H(D)}$ on $H_*(D)$.

We claim that $m_3^{H(D)}(\mathcal{U}, \mathcal{V}, \mathbf{f}) = \mathbf{e}$. To see this, we use the formula on the left of Figure 2.1. In the present situation, $\mathbf{a} = \mathcal{U} \otimes \mathcal{V}$, $\mathbf{z} = \mathbf{f}$. Note that $\Delta(\mathcal{U} \otimes \mathcal{V}) = (\mathcal{U}) \otimes (\mathcal{V})$, so we compute

$$m_3^{H(D)}(\mathcal{U}, \mathcal{V}, \mathbf{f}) = \pi(m_2^D(\mathcal{U}, h(m_2^D(\mathcal{V}, i(\mathbf{f})))) = \pi(m_2^D(\mathcal{U}, \mathbf{a})) = \mathbf{e}.$$

3. Plumbed manifolds and plumbed links

The goal of this section is to recall the notions of plumbed manifolds, resolution graphs, and plumbed links. We pay particular attention to algebraic links, which are L-space links by work of Gorsky and Némethi [12]. Additionally, in Proposition 3.18 we describe a slightly wider class of plumbed links which are also L-space links.

3.1. Review of plumbed manifolds

To set up the notation, we recall the constructions of 3-manifolds via plumbing. We refer the reader to [38] for a detailed exposition.

Suppose G is a finite graph. We let V_G be the set of its vertices. We assume that each $v \in V_G$ has an associated weight $\lambda_v \in \mathbb{Z}$. From V_G we construct a real 4-manifold X_G as follows. For each $v \in V_G$, we consider T_v , the oriented disk bundle over S^2 with Euler

number λ_v . The manifold X_G is obtained by taking a disjoint union of all the T_v and gluing them using the following recipe. Whenever two vertices $v, v' \in V_G$ are connected by an edge e , we trivialize the bundles T_v and $T_{v'}$ over chosen disks in the base. Then we glue these bundles together by an orientation-preserving diffeomorphism that swaps the base and the fiber. Refer to [8, Example 4.6.2] or [38] for more details.

By convention, if G is not connected, we take a boundary connected sum of the manifolds X_{G_i} corresponding to the connected components G_i of G .

Definition 3.1. The manifold X_G is called the *plumbed 4-manifold* associated with G . Its boundary $Y_G = \partial X_G$ is the *plumbed 3-manifold* associated with G .

The construction of a plumbed manifold can be done in a relative setting, providing a pair consisting of a 3-manifold and a link contained in it. The starting data is a graph Γ with vertices partitioned into two sets, $V_G \sqcup V_\uparrow$. We call $V_\uparrow$ the *arrow* vertices, and V_G the *non-arrow* vertices. We do not add weights to $V_\uparrow$. We call such a graph an *arrow-decorated plumbing graph*.

Remark 3.2. From a topological perspective, it is most natural to require each vertex of $V_\uparrow$ to have valence 1. However, in the combinatorial construction of link lattice homology, we do not need to make this assumption.

We write $G \subseteq \Gamma$ for the full subgraph spanned by the non-arrow vertices. The vertices $V_\uparrow$ determine a link $L_\uparrow$ in Y_G as follows. Suppose $v \in V_\uparrow$ is adjacent to a non-arrow vertex $w \in V_G$. We let L_v be a circle fiber of the S^1 -bundle $\partial T_w \rightarrow S^2$ such that the projection of L_v onto S^2 is disjoint from all the disks used to plumb the disk bundles of other non-arrow vertices. If more than one arrow vertex is adjacent to the same non-arrow vertex w , we require each of the corresponding components of $L_\uparrow$ to be fibers of $\partial T_w \rightarrow S^2$ over distinct points.

We define $L_\uparrow$ as the union of the circle fibers L_v ranging over $v \in V_\uparrow$.

Definition 3.3. The link $L_\uparrow \subseteq Y_G$ is called the *plumbed link* associated with Γ . We say that a link $L \subseteq Y$ is a *plumbed link* if there exists an arrow-decorated plumbing graph Γ and a diffeomorphism $(Y, L) \cong (Y_G, L_\uparrow)$.

Remark 3.4. One can also consider more general plumblings of disk bundles over higher genus surfaces. To do so, one considers a plumbing graph where each vertex $v \in V_G$ is assigned an additional weight, corresponding to the genus of the base space of the disk bundle. However, the resulting manifold Y_G is not a rational homology sphere if at least one surface has positive genus. In the present paper, we are mostly concerned with rational homology spheres, so we restrict the discussion to the case where all surfaces are spheres. We refer to [38] for more details.

Definition 3.5. Suppose G is a plumbing tree with no arrow vertices. We define the *incidence matrix* Q_G as follows. The diagonal entries are the weights associated to vertices, while the off-diagonal terms are 1 or zero, depending on whether or not the two vertices are connected by an edge.

By construction, Q_G represents the intersection form on X_G . As the intersection form on X_G determines the homology of Y_G , we have the following result.

Lemma 3.6. *There is an isomorphism $H_1(Y_G; \mathbb{Z}) \cong \text{coker } Q_G$. In particular, Y_G is a rational homology sphere if and only if $\det Q_G \neq 0$.*

There is a well-known description of $(Y_G, L_\uparrow)$ in terms of Dehn surgery (see [8, Example 4.6.2]). We form a partitioned link $L_\Gamma = L_G \sqcup L_\uparrow$ in S^3 as follows. For each vertex of Γ , we add an unknotted component to L_Γ , and for each edge, we add a clasp between the corresponding components. The link L_Γ may alternatively be described as an iterated connected sum of Hopf links. The weights on the vertices in V_G determine an integral framing Λ on L_G , as in Definition 3.5. Then

$$(Y_G, L_\uparrow) \cong (S^3_\Lambda(L_G), L_\uparrow).$$

We remark a slight abuse of notation: $L_\uparrow$ denotes both the link in S^3 (as a part of $L_\Gamma \subseteq S^3$) and its image in the plumbed manifold Y_G .

3.2. Plumbed manifolds and resolutions of analytic singularities

One of the main motivations for introducing plumbed manifolds comes from resolutions of singularities. We give now a short account of plumbed manifolds obtained from surface singularities. We refer the reader to the introductory lectures of Némethi [32], or to [36, Section 3.3], [29], [37, Chapter 4] for more details and references.

First we focus on the absolute case corresponding to graphs with no arrow vertices. Later on we discuss the relative case of embedded resolutions, leading to plumbed links.

Suppose (X, x_0) is a (germ of a) normal complex analytic surface. The word ‘normal’ refers to the property of the local ring $\mathcal{O}_{x_0}(X)$ being integrally closed (see [15, Exercise I.3.7]). This implies, among other things, that x_0 is an isolated singular point. See [22, 32] for more details. The surface (X, x_0) can be analytically embedded into $(\mathbb{C}^N, 0)$ for N sufficiently large. Let B_ε be a ball in $\mathbb{C}^N$ with center at 0 and radius $\varepsilon > 0$. It is known (see [31]) that the diffeomorphism type of the intersection $L_X := X \cap \partial B_\varepsilon$ is independent of ε and of the embedding of X into $\mathbb{C}^N$, provided $\varepsilon > 0$ is small enough. Moreover, the pair $(B_\varepsilon, X \cap B_\varepsilon)$ is topologically a cone over $(\partial B_\varepsilon, L_X)$. The space L_X is a smooth real 3-dimensional manifold. We call it the *link of the surface singularity* (X, x_0) .

We stress that we study the local behavior of X near x_0 . From the perspective of algebraic geometry, this is emphasized by saying that X is a germ of a surface. The reader unfamiliar with this notion might assume that we replace X by $X \cap B_\varepsilon$, where B_ε is as above.

The manifold L_X admits another description. We let $(\tilde{X}, E)$ be a resolution of (X, x_0) , that is, a smooth complex analytic surface together with a map $\pi: (\tilde{X}, E) \rightarrow (X, x_0)$ which is one-to-one except on $\pi^{-1}(x_0) = E$. Now $E = \sum E_i$ is a union of smooth complex curves (Riemann surfaces) intersecting transversally. Each E_i is assigned a number λ_i which is its self-intersection. The curves E_i are referred to as the *exceptional components* of the map π .

With the resolution we can assign two objects. One is the *dual graph* G_X of the resolution. Its vertices correspond to the divisors E_i . Each vertex is assigned a weight λ_i . There is an additional weight of a vertex by the genus of E_i (in the present paper we will consider only the case where each E_i is a sphere; see Remark 3.4). We add $|E_i \cap E_j|$ edges between vertices v_i and v_j when $i \neq j$. We add no self-edges. There is a matrix Q_G associated with G_X as in Definition 3.5. We have the following result.

Proposition 3.7. (a) *The link L_X of the singularity (X, x_0) is diffeomorphic to Y_{G_X} .*
 (b) *L_X is a rational homology sphere if each E_i is a sphere and G_X is a tree.*

The manifold L_X determines the graph G_X up to a precisely described equivalence relation; see [38]. The way the resolution is constructed implies that Q_G is negative definite. A deep theorem of Grauert [13] shows that this characterizes links of analytic singularities among all plumbed links.

Theorem 3.8. *If Q_G is negative definite, then Y_G is a link of an analytic singularity.*

We stress that the statement is far from true if the word ‘analytic’ is replaced by ‘algebraic’. Moreover, in general, there is no uniqueness. While G_X determines the diffeomorphism type of L_X , there might be analytically different singularities with the same link L_X . There is a vast research area concerning which invariants of X depend on the link L_X , and which depend on the analytic structure. We refer the reader to the book [36]. Examples of invariants depending on the analytic structure of X include the geometric genus p_g (see [36, Section 6.8]), the embedding dimension (minimal N for which X embeds into $\mathbb{C}^N$, see [36, Example 6.7.17]), and the Hilbert–Samuel function (see [36, Section 5.1.40]).

3.3. Embedded resolutions

We now consider embedded singularities, which are pairs of analytic spaces $Z \subseteq X$, together with a point $x_0 \in Z$, where X and Z are possibly singular at x_0 . We restrict our attention to the case where $\dim_{\mathbb{C}} X = 2$ and $\dim_{\mathbb{C}} Z = 1$. For an introduction to embedded singularities, we refer to [36], especially Section 2.2. An overview of singularity theory is given in [37, Section 4.3]. Graph links, and their connection with singularity theory, are described in [6].

Definition 3.9. An *embedded singularity* is a triple (X, Z, x_0) , where (X, x_0) is a (germ of a) normal complex analytic surface and $Z \subseteq X$ is a complex analytic curve passing through x_0 .

Example 3.10. If $X = \mathbb{C}^2$, an embedded singularity is precisely a plane curve singularity.

Embed X analytically in $\mathbb{C}^N$ with x_0 mapped to 0. Take a small ball B_ε around x_0 in $\mathbb{C}^N$ as above. For sufficiently small $\varepsilon > 0$, the triple $(B_\varepsilon, X \cap B_\varepsilon, Z \cap B_\varepsilon)$ is topologically a cone over $(\partial B_\varepsilon, L_X, L_Z)$, where L_X and L_Z are, respectively, the intersections

of X and Z with ∂B_ε . The diffeomorphism type of the pair (L_X, L_Z) depends on neither the choice of ε nor the choice of embedding.

Definition 3.11. The pair (L_X, L_Z) is called the *link of the embedded singularity* (X, Z, x_0) .

Example 3.12. Suppose $(X, x_0) = (\mathbb{C}^2, 0)$, and Z is a plane algebraic curve passing through 0. Then $L_X = S^3$, and the link L_Z is precisely the algebraic link in the ordinary sense.

Since the study of singularities is local, we consider only the germ of the singularity. We note that, by definition, (X, Z) and $(X \cap B_\varepsilon, Z \cap B_\varepsilon)$ have the same germ. In particular, we may and will assume that (X, Z) is topologically a cone over (L_X, L_Z) .

We can recover (L_X, L_Z) from an embedded resolution. By an *embedded resolution* of (X, Z, x_0) we mean the triple $(\tilde{X}, \tilde{Z}, E)$ together with a proper analytic map $\pi: (\tilde{X}, \tilde{Z}, E) \rightarrow (X, Z, x_0)$ satisfying the following conditions:

- $\pi: \tilde{X} \rightarrow X$ is one-to-one away from E . In particular, the restriction $\pi|_{\tilde{Z}}$ is one-to-one away from $\tilde{Z} \cap E$.
- $\tilde{X}$ is a smooth surface and $\tilde{Z}$ is a smooth complex curve.
- Each algebraic component of E is a projective (that is, closed) smooth complex curve.
- The union $E \cup \tilde{Z}$ has only transverse double points as singularities.

As in the non-embedded case, the smooth complex curves whose union is E are referred to as the *exceptional components*.

Given the embedded resolution, we can create a dual graph of the resolution. The construction is in two steps. First, out of E , we construct the graph G_X as above. Next, if $v_i \in V_{G_X}$ and E_i is the corresponding component of E , we adjoin $|E_i \cap \tilde{Z}|$ arrow vertices to v_i . We denote the resulting graph by $\Gamma_{X,Z}$. Recall that we work locally (topologically, we have replaced X by $X \cap B_\varepsilon$). Therefore, $\tilde{Z}$ is a union of disks, each intersecting the graph E in precisely one point. That is, every arrow vertex of $\Gamma_{X,Z}$ corresponds to a connected component of $\tilde{Z}$. The following result is classical.

Proposition 3.13 (see e.g. [36, Proposition 3.3.8]). *The pair $(Y_{G_X}, L_\uparrow)$ is diffeomorphic to (L_X, L_Z) .*

Our next aim is to explain the relative analog of Grauert's Theorem 3.8. As the statement is slightly technical, we give some extra explanation. Let $g: X \rightarrow \mathbb{C}$ be a reduced analytic map such that $g^{-1}(0) = Z$. Here, *reduced* means that g is not divisible by the square of a non-invertible analytic function on X . Then g induces an analytic map $\tilde{g}: \tilde{X} \rightarrow \mathbb{C}$ via $\tilde{g} = g \circ \pi$. Let v be a vertex of $\Gamma := \Gamma_{X,Z}$. The vertex v corresponds either to an exceptional component E_v (if v is a non-arrow vertex), or to a component $\tilde{Z}_v$ of $\tilde{Z}$, if v is an arrow vertex. In both cases, $\tilde{g}$ vanishes on that component. We let $m_v > 0$ denote the order of vanishing, and call it the *multiplicity* of the vertex v . Note that since g is reduced, $m_v = 1$ for all arrow vertices (see [37, Section 4.3.2]). The multiplicities and

the weights satisfy the following compatibility relation (see [37, (4.1.5)]):

$$\lambda_v m_v + \sum_{w \in V_v} m_w = 0 \quad (3.1)$$

for each non-arrow vertex $v \in V_G$, where V_v denotes the set of all vertices in V_Γ adjacent to v . Note that (3.1), together with the condition $m_v = 1$ for all arrow vertices, uniquely determines all other multiplicities. However, unless Q_G is unimodular, the multiplicities need not be integral. If they are not integral, such a plumbed link cannot be realized as a link of an embedded analytic singularity.

We now state a relative analog of Grauert's Theorem 3.8 (see [52, Corollaire 5.5]).

Proposition 3.14. *Let Γ be a graph with vertices $V = V_G \cup V_\uparrow$ such that Q_G is negative definite. If assigning multiplicity 1 to each arrow vertex of Γ leads to integral positive multiplicities on all vertices of V_G via the compatibility relation (3.1), then $(Y_G, L_\uparrow)$ is a link of an embedded analytic singularity.*

3.4. Rationality

Suppose (X, x_0) is an analytic singularity. We define its *geometric genus* to be $p_g = h^1(\mathcal{O}_{\tilde{X}})$, where $\tilde{X}$ is a resolution of X (see [32, Section 2]). Many properties of geometric genus are given in various chapters of [36]. The definition of p_g does not depend on the choice of resolution. Geometric genus is an invariant of the analytic structure of X ; there are known examples of singularities with the same link, but different geometric genus [32, Section 4.8]. Put differently, in general p_g cannot be read off from the combinatorics of the resolution graph G .

An exception is the case of *rational* singularities, which are characterized by the property that $p_g = 0$ (see [36, Section 7.1]). Given a graph G , we can determine whether it represents a rational singularity; the answer is due to Artin [1] (see also [32, Theorem 3.8] and [36, Theorem 7.1.2]). For instance, if Y_G is a link of a rational singularity, then $b_1(Y_G) = 0$.

By studying the relation between p_g and combinatorial invariants of X_G encoded by G , Némethi proved the following groundbreaking result:

Theorem 3.15 ([35]). *The singularity (X, x_0) is rational if and only if L_X is an L-space.*

The proof of Theorem 3.15 can also be done using the recently proved equivalence of lattice and Heegaard Floer homology [59]. As the first step, one uses Némethi's theorem stating that if G is negative definite, then X is a rational singularity if and only if the reduced lattice homology of Y_G is zero (see [33, 34]). Next, one refers to [59] to show that the reduced lattice homology of Y_G vanishes if and only if Y_G is an L-space.

Remark 3.16. Theorem 3.15 only characterizes graphs representing an L-space among graphs with negative definite incidence matrix Q_G . There are indefinite graphs representing L-spaces. For example, if G is a linear plumbing such that Q_G is non-degenerate,

then Y_G is a lens space, regardless of whether Q_G is definite or not. To the best of our knowledge, there is not a generalization of Theorem 3.15 to indefinite graphs.

3.5. Algebraic links and L-space links

Let us recall the following definition.

Definition 3.17. Let L be a link in a rational homology 3-sphere. We say L is an *L-space link* if all sufficiently large positive surgeries on L are L-spaces.

Though usually one considers L-space links in S^3 or an integer homology sphere (see e.g. [12, 28]), we note that the same definition can be applied to links in rational homology 3-spheres. Suppose K is a rationally null-homologous knot in Y . Then Morse framings on K can be identified with an affine $\mathbb{Z}$ -subspace of $\mathbb{Q}$ by taking the intersection number of the framing (viewed as a parallel longitude of K) with a rational Seifert surface. Hence, large surgeries on rationally null-homologous links are surgeries with Morse framings which are sufficiently large in $\mathbb{Q}^\ell$ with respect to this identification.

In [12] Gorsky and Némethi studied which plumbed links in plumbed manifolds are L-space links. Their main result is that every algebraic link in S^3 is an L-space link [12, Theorem 2]. Their proof works in a more general setting, leading to the following statement, which is given in [12, Theorem 12] and the ensuing discussion.

Proposition 3.18. *Suppose Γ is a graph such that Q_G is negative definite and Y_G is a link of a rational singularity. Then $L_\uparrow$ is an L-space link.*

We stress that the assumption on multiplicities as in Proposition 3.14 is never used in the proof of Proposition 3.18. That is, Proposition 3.18 does not require that the link $L_\uparrow$ be a link of an analytic singularity; it only makes a restriction on the graph G being the graph representing a rational singularity.

To see this, we quickly sketch the argument of [12, proof of Theorem 2] proving Proposition 3.18. One first extends the graph G to another graph, $\tilde{G}_0$, which replaces all arrow vertices of Γ by chains of -2 -weighted vertices ending with a -1 -weighted vertex. By construction, $\tilde{G}_0$ can be contracted to G by successive blow-downs, that is, $\tilde{G}_0$ represents the manifold Y_G . Hence, it is a rational graph. Next, a sufficiently large positive surgery on $L_\uparrow$ can be represented as a subgraph of $\tilde{G}_0$ as long as the chains of -2 -weighted vertices are long enough. Since any subgraph of a rational graph is rational by Laufer's criterion, large positive surgeries on $L_\uparrow$ are represented by rational graphs. Rational graphs represent L-spaces by [33, Theorems 6.3, 8.3]. That is, a sufficiently large positive surgery on $L_\uparrow$ is an L-space. This means that $L_\uparrow$ is an L-space link.

It is well-known that an algebraic knot is determined by its Alexander polynomial. A natural question is whether this result generalizes to plumbed L-space links. The following well-known fact is due to Yamamoto.

Proposition 3.19 (see [55]). *Two algebraic links in S^3 with the same Alexander polynomial are equal.*

The result of Yamamoto relies on the characterization of algebraic links in S^3 , due to Zariski [56]; we refer to [6] for this characterization. In particular, this result does not admit direct generalizations to links in other 3-manifolds. We now give a few counterexamples for some naive attempts to generalize the result. Proposition 3.20 is not used in the present paper. Rather it indicates that algebraic links cannot be distinguished by Alexander polynomials, hence, they cannot be distinguished by Heegaard Floer homology.

Proposition 3.20. (a) *There exist non-isotopic plumbed L-space links in S^3 with the same Alexander polynomial.*
 (b) *There exist non-isotopic knots that are links of embedded analytic surface singularities with the same Alexander polynomial.*

Proof. Item (a) is classical. We know that the $(2, 3)$ -cable on the positive trefoil is a plumbed knot, and it is an L-space knot by [16, Theorem 1.10]. However, its Alexander polynomial is the same as that of the $T(3, 4)$ torus knot.

Item (b) expands on results of Campillo, Delgado and Gussein-Zade [3]. In fact, in [3, Section 3, Example 2] there are two knots in the Poincaré sphere with the same Alexander polynomial, represented by two plumbing diagrams. We quickly recall their construction. The starting point is the resolution of the E_8 singularity given by the plumbing graph in Figure 3.1. The first knot is obtained by taking the plumbing diagram of the E_8 singularity and adding to it an arrow vertex at the component marked A in Figure 3.1.

The second knot is obtained by drawing an A_4 singularity (i.e. with local equation $x^5 - y^2 = 0$) transversally to a point at the B component. The resolution of that singularity yields the plumbing graph drawn in Figure 3.2 (compare [3, Figure 2]).




Fig. 3.1. Resolution graph of the E_8 singularity. All vertices correspond to spheres with self-intersection -2 . The meaning of components A and B is explained in the text.




Fig. 3.2. Plumbing graph of the second knot in Proposition 3.20 (b). All weights that are not explicitly marked have value -2 .

An explicit algorithm described in [6, Chapter 20] transforms these two plumbing graphs into graph links, which are presented in Figure 3.3. Using the algorithm of [39], we can compute the signature functions of these links, and we present them in Figure 3.4, omitting straightforward calculations. The signatures are different, so the knots are different. ■




Fig. 3.3. The two links of [3] represented as graph links of [6].




Fig. 3.4. The signature functions $x \mapsto \sigma(e^{2\pi ix})$, $x \in [0, 1/2]$, for links of Figure 3.3.

Remark 3.21. The signatures might be different, but the discontinuities of the signature functions appear at the same places. This is consistent with the fact that the Alexander polynomials of the two knots are equal. (It is well-known that the jumps of the signature functions occur only at roots of the Alexander polynomial.) The fact that the signatures of the two knots in Proposition 3.20(b) being different means not only that the knots are not isotopic, but also that they are not concordant.

4. Link lattice homology

In this section, we recall some basics about Heegaard Floer homology, and subsequently define our link lattice complex.

4.1. Background on link Floer homology

To fix the notation and terminology, we give some necessary background on link Floer homology. We assume some familiarity with basics of Heegaard Floer homology and its refinements for knots [44, 46, 48, 53].

Let L be an ℓ -component link in a 3-manifold Y . Recall [48, Section 3.5] that the pair (Y, L) can be encoded in a multi-pointed Heegaard link diagram $(\Sigma, \alpha, \beta, \mathbf{w}, \mathbf{z})$, as follows:

- (1) Σ is a closed oriented genus g surface.
- (2) $\alpha = \{\alpha_1, \dots, \alpha_{g+\ell-1}\}$ and $\beta = \{\beta_1, \dots, \beta_{g+\ell-1}\}$ are collections of simple closed curves on Σ . The curves α_i are pairwise non-intersecting. Also, the curves β_i are pairwise non-intersecting. Moreover, α and β each span a g -dimensional subspace of $H_1(\Sigma; \mathbb{Z})$.

- (3) $\mathbf{w} = \{w_1, \dots, w_\ell\}$, $\mathbf{z} = \{z_1, \dots, z_\ell\}$. Each component of $\Sigma \setminus \alpha$ (respectively of $\Sigma \setminus \beta$) contains a single point of $\mathbf{w}$ and a single point of $\mathbf{z}$.

It is not hard to see that there exists a Heegaard link diagram for any pair (Y, L) . Furthermore, any two diagrams can be connected by a sequence of Heegaard moves for link diagrams (see [48, Theorem 4.7]).

Given a Heegaard link diagram, we consider the Lagrangian tori

$$\mathbb{T}_\alpha = \alpha_1 \times \dots \times \alpha_{g+\ell-1}, \quad \mathbb{T}_\beta = \beta_1 \times \dots \times \beta_{g+\ell-1}$$

in the symmetric product $\text{Sym}^{g+\ell-1}(\Sigma)$. The *link Floer chain complex*, $\mathcal{CFL}(Y, L)$, is a free chain complex over

$$\mathcal{R}_\ell = \mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$$

generated by the intersection points $\mathbf{x} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta$ with the differential counting pseudo-holomorphic curves in $\text{Sym}^{g+\ell-1}(\Sigma)$ via

$$\partial \mathbf{x} = \sum_{\mathbf{y} \in \mathbb{T}_\alpha \cap \mathbb{T}_\beta} \sum_{\substack{\phi \in \pi_2(\mathbf{x}, \mathbf{y}) \\ \mu(\phi)=1}} (\#\mathcal{M}(\phi)/\mathbb{R}) \mathcal{U}_1^{n_{w_1}(\phi)} \dots \mathcal{U}_\ell^{n_{w_\ell}(\phi)} \mathcal{V}_1^{n_{z_1}(\phi)} \dots \mathcal{V}_\ell^{n_{z_\ell}(\phi)} \mathbf{y}.$$

Here the sum is taken over all homotopy classes $\pi_2(\mathbf{x}, \mathbf{y})$ of maps ϕ of the unit disk $\mathbb{D} \subseteq \mathbb{C}$ to $\text{Sym}^{g+\ell-1}(\Sigma)$, where $\phi(-1) = \mathbf{x}$, $\phi(1) = \mathbf{y}$, $\phi(\partial\mathbb{D} \cap \{\text{Im}(z) \leq 0\}) \subseteq \mathbb{T}_\alpha$, $\phi(\partial\mathbb{D} \cap \{\text{Im}(z) \geq 0\}) \subseteq \mathbb{T}_\beta$. Here, $\mu(\phi)$ denotes the Maslov index of the class ϕ . The space $\mathcal{M}(\phi)$ consists of all pseudo-holomorphic curves representing the class ϕ , for a generic 1-parameter family of almost complex structures on $\text{Sym}^{g+\ell-1}(\Sigma)$. For $x \in \Sigma \setminus (\alpha \cup \beta)$, we denote by $n_x(\phi)$ the intersection number of $\{x\} \times \text{Sym}^{g+\ell-2}(\Sigma) \subseteq \text{Sym}^{g+\ell-1}(\Sigma)$ with $\phi(\mathbb{D})$. We refer to [46] for more details.

There is a map $\mathfrak{s}_w$ from $\mathbb{T}_\alpha \cap \mathbb{T}_\beta$ to the set of Spin^c structures on Y . The component of the map ∂ from $\mathbf{x}$ to $\mathbf{y}$ can be non-trivial only if $\mathfrak{s}_w(\mathbf{x}) = \mathfrak{s}_w(\mathbf{y})$. That is, the chain complex $\mathcal{CFL}(Y, L)$ splits as a direct sum of complexes $\mathcal{CFL}(Y, L, \mathfrak{s})$ for $\mathfrak{s} \in \text{Spin}^c(Y)$.

There is a completed version of $\mathcal{CFL}(Y, L)$ regarded as a module over the ring of power series

$$\mathcal{R}_\ell := \mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]],$$

namely we set

$$\mathcal{CFL}(Y, L) := \mathcal{CFL}(Y, L) \otimes_{\mathcal{R}_\ell} \mathcal{R}_\ell.$$

In other words, $\mathcal{CFL}(Y, L)$ has the same generators as $\mathcal{CFL}(Y, L)$ and the same differential, except that we work over a larger ring. The completed version appears in the surgery formula.

When Y is a rational homology 3-sphere, the link Floer homology groups have several gradings. Firstly, there is a $\mathbb{Q} \times \mathbb{Q}$ -valued Maslov bigrading, denoted $(\text{gr}_w, \text{gr}_z)$, as well as a $\mathbb{Q}^\ell$ -valued Alexander grading A . Furthermore,

$$(\text{gr}_w, \text{gr}_z)(\mathcal{U}_i) = (-2, 0), \quad (\text{gr}_w, \text{gr}_z)(\mathcal{V}_i) = (0, -2) \quad \text{and} \quad A(\mathcal{V}_i) = -A(\mathcal{U}_i) = e_i,$$

where e_i is the standard i -th coordinate vector in $\mathbb{Q}^\ell$. Note also that

$$\mathrm{gr}_w - \mathrm{gr}_z = 2 \sum_{i=1}^{\ell} A_i.$$

Remark 4.1. In this paper, we normalize gr_w so that the isomorphism

$$H_*(\mathcal{CFL}(Y, L)/(\mathcal{V}_1 - 1, \dots, \mathcal{V}_\ell - 1)) \cong HF^-(Y)$$

is grading preserving. In the above, we are writing $HF^-(Y)$ for the Heegaard Floer homology computed with a singly pointed Heegaard diagram for Y . We make a similar normalization for gr_z . Equivalently, our grading convention is that HF^- of a 3-manifold is invariant under adding extra basepoints as a graded module (compare [48, Section 6.1]). We note that some authors normalize the Maslov gradings so that $H_*(\mathcal{CFL}(Y, L)/(\mathcal{V}_1 - 1, \dots, \mathcal{V}_\ell - 1))$ is isomorphic instead to $HF^-(Y)[(\ell - 1)/2]$.

4.2. Lattice homology

We recall the definition of lattice homology [34]. We use the notation of Ozsváth, Stipsicz and Szabó [41], since our construction of link lattice homology is slightly easier to describe using their notation. Let G be a plumbing tree, and write V_G for the vertices of G . Write $\mathbb{P}(V_G)$ for the power set of V_G (i.e. the set of all subsets of V_G). The *lattice complex* is the $\mathbb{F}[[U]]$ -module

$$\mathbb{CF}(G) := \prod_{[K, E] \in \mathrm{Char}(G) \times \mathbb{P}(V_G)} \mathbb{F}[[U]] \otimes \langle [K, E] \rangle,$$

where $\mathrm{Char}(G) \subseteq H^2(X_G; \mathbb{Z})$ denotes the set of characteristic elements of $H^2(X_G; \mathbb{Z})$ on the 4-manifold X_G . Recall that $K \in H^2(X_G; \mathbb{Z})$ is *characteristic* if $K(x) \equiv x \cdot x \pmod{2}$ for all $x \in H_2(X_G; \mathbb{Z})$.

We now define the differential on $\mathbb{CF}(G)$. Note that each vertex $v \in V_G$ determines an element of $H_2(X_G)$, which is the class in $H_2(X_G)$ of the base space of the disk bundle T_v used in the plumbing construction. For $I \subseteq E$, one defines

$$2f(K, I) = \left(\sum_{v \in I} K(v) \right) + \left(\sum_{v \in I} v \right) \cdot \left(\sum_{v \in I} v \right).$$

Note that the right-hand side is always an even integer, because K is characteristic. In particular, $f(K, I)$ is an integer. We set $g(K, E) = \min \{f(K, I) : I \subseteq E\}$. Next, we define

$$A_v(K, E) = g(K, E - v), \quad B_v(K, E) = \min \{f(K, I) : v \in I \subseteq E\}.$$

Set

$$a_v(K, E) = A_v(K, E) - g(K, E), \quad b_v(K, E) = B_v(K, E) - g(K, E).$$

By the definition of $g(K, E)$, one can see that $g(K, E - v) \geq g(K, E)$. Similarly, $B_v(K, E) \geq g(K, E)$. Hence, $a_v(K, E)$ and $b_v(K, E)$ are both non-negative integers.

The differential on $\mathbb{C}\mathbb{F}(G)$ is defined by the formula

$$\partial[K, E] = \sum_{v \in E} U^{a_v(K, E)} \otimes [K, E - v] + \sum_{v \in E} U^{b_v(K, E)} \otimes [K + 2v^*, E - v], \quad (4.1)$$

where v^* is the Poincaré dual of v . Note that because of the factor 2, $K + 2v^*$ is characteristic if and only if K is. Equation (4.1) is extended linearly over $\mathbb{F}[[U]]$. We will refer to the first summand in (4.1) as the *A-terms* in the differential, and to the second summand as the *B-terms*.

4.3. The link lattice complex

We now suppose that Γ is a plumbing tree whose vertex set is partitioned into two sets:

$$V_\Gamma = V_G \cup V_\uparrow.$$

Recall that the components of V_G are equipped with a framing, while those of $V_\uparrow$ are not.

The vertices $V_\uparrow$ determine a link $L_\uparrow$ in the 3-manifold Y_G . We assume that each component of $L_\uparrow$ is rationally null-homologous in Y_G . This occurs, for example, when the incidence matrix Q_G is non-singular.

To define the link lattice complex, we first pick a framing on the components of $V_\uparrow$ arbitrarily. In Proposition 4.13, we will show that the choice of framing on $V_\uparrow$ does not affect the link lattice complex.

We define the *link lattice complex* $\mathbb{C}\mathbb{F}\mathbb{L}(\Gamma, V_\uparrow)$ as the quotient of $\mathbb{C}\mathbb{F}(\Gamma)$ by the subspace generated over $\mathbb{F}[[U]]$ by the tuples $[K, E]$ where $V_\uparrow \not\subseteq E$. Equivalently, we may view $\mathbb{C}\mathbb{F}\mathbb{L}(\Gamma, V_\uparrow)$ as being generated by $[K, E]$ where $V_\uparrow \subseteq E$, equipped with the quotient complex differential. We think of the differential on $\mathbb{C}\mathbb{F}\mathbb{L}(\Gamma, V_\uparrow)$ as being given by the same formula as in (4.1), but with the sums taken only over $v \in E \cap V_G$.

4.4. The module structure

Recall that

$$\mathcal{R}_\ell = \mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell].$$

We now describe the action of $\mathcal{R}_\ell$ on link lattice homology, where $\ell = |V_\uparrow|$. Write $V_\uparrow = \{v_1, \dots, v_\ell\}$. For each $i \in \{1, \dots, \ell\}$, there is an induced element $\mu_i^* \in H_2(X_\Gamma, Y_\Gamma) \cong H^2(X_\Gamma)$. This element is dual to the class $v_i \in H_2(X_\Gamma)$ in the sense that $\mu_i^*(v_i) = 1$ and $\mu_i^*(w) = 0$ for all $w \in V_\Gamma \setminus \{v_i\}$. The class μ_i^* is represented by the co-core disk of the 2-handle corresponding to v_i .

Define

$$\delta_i^+(K, E) = g(K + 2\mu_i^*, E) - g(K, E), \quad \delta_i^-(K, E) = g(K - 2\mu_i^*, E) - g(K, E).$$

An easy computation shows that

$$f(K \pm 2\mu_i^*, I) = \begin{cases} f(K, I) \pm 1 & \text{if } v_i \in I, \\ f(K, I) & \text{if } v_i \notin I. \end{cases}$$

In particular,

$$\delta_i^+(K, E) \in \{0, 1\} \quad \text{and} \quad \delta_i^-(K, E) \in \{0, -1\}$$

for all i .

For $i \in \{1, \dots, \ell\}$, we define

$$\mathcal{U}_i \cdot [K, E] = \begin{cases} U[K - 2\mu_i^*, E] & \text{if } \delta_i^-(K, E) = 0, \\ [K - 2\mu_i^*, E] & \text{if } \delta_i^-(K, E) = -1, \end{cases} \quad (4.2)$$

and

$$\mathcal{V}_i \cdot [K, E] = \begin{cases} U[K + 2\mu_i^*, E] & \text{if } \delta_i^+(K, E) = 1, \\ [K + 2\mu_i^*, E] & \text{if } \delta_i^+(K, E) = 0. \end{cases} \quad (4.3)$$

We extend $\mathcal{U}_i$ and $\mathcal{V}_i$ to the entire link lattice complex by declaring them to be $\mathbb{F}[U]$ -equivariant. Equivalently, we set

$$\begin{aligned} \mathcal{U}_i \cdot [K, E] &= U^{g(K-2\mu_i^*, E) - g(K, E) + 1} [K - 2\mu_i^*, E], \\ \mathcal{V}_i \cdot [K, E] &= U^{g(K+2\mu_i^*, E) - g(K, E)} [K + 2\mu_i^*, E]. \end{aligned} \quad (4.4)$$

Lemma 4.2. *If $v_i \in V_\uparrow$, then the endomorphisms $\mathcal{U}_i$ and $\mathcal{V}_i$ are chain maps.*

Proof. The differential of $\mathbb{CFL}(\Gamma, V_\uparrow)$ is given by modifying (4.1) to sum only over $v \in V_G$. Clearly the summands of $\mathcal{V}_i \partial[K, E]$ are in bijection with the summands of $\partial \mathcal{V}_i[K, E]$. It remains to show that the powers of U coincide. We consider the A -terms of the differential first. The power of U from $\mathcal{V}_i \partial[K, E]$ of $[K + 2\mu_i^*, E - v]$ is

$$g(K + 2\mu_i^*, E - v) - g(K, E - v) + g(K, E - v) - g(K, E),$$

whereas the power of U from $\partial(\mathcal{V}_i[K, E])$ is

$$g(K + 2\mu_i^*, E - v) - g(K + 2\mu_i^*, E) + g(K + 2\mu_i^*, E) - g(K, E).$$

These are obviously equal. Similarly, for the B -terms of the differential we use the equality

$$B_v(K, E) = (K(v) + v \cdot v)/2 + g(K + 2v^*, E - v).$$

From now on, the argument is similar to the case of type- A terms. We note that

$$(K + 2\mu_i^*)(v) + v \cdot v = K(v) + v \cdot v \quad (4.5)$$

if $v \neq v_i$. For the B -terms of the differential, the U power of the term from $\partial \mathcal{V}_i[K, E]$ is

$$B_v(K + 2\mu_i^*, E) - g(K + 2\mu_i^*, E) + g(K + 2\mu_i^*, E) - g(K, E).$$

The U power from $\mathcal{V}_i \partial[K, E]$ is

$$g(K + 2v^* + 2\mu_i^*, E - v) - g(K + 2v^*, E - v) + B_v(K, E) - g(K, E).$$

The difference between these terms is $((K + 2\mu_i^*)(v) + v \cdot v)/2 - (K(v) + v \cdot v)/2$, which vanishes by (4.5).

The proof of the claim about the map $\mathcal{U}_i$ uses essentially the same logic. $\blacksquare$

Lemma 4.3. (1) For each i , we have $\mathcal{U}_i \mathcal{V}_i = \mathcal{V}_i \mathcal{U}_i = U$.

(2) For all i, j , the commutators $[\mathcal{U}_i, \mathcal{V}_j]$, $[\mathcal{U}_i, \mathcal{U}_j]$ and $[\mathcal{V}_i, \mathcal{V}_j]$ vanish.

Proof. All of the stated relations are easily derived from (4.4). $\blacksquare$

Lemma 4.4. The action of $\mathcal{R}_\ell$ on $\mathbb{C}\mathbb{F}\mathbb{L}(\Gamma, V_\uparrow)$ extends to an action of the ring of power series

$$\mathcal{R}_\ell = \mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]].$$

Proof. We use the following fact about direct products. Suppose that $(A_i)_{i \in I}$ and $(B_j)_{j \in J}$ are families of vector spaces. Suppose that we have a function $\gamma: I \times J \rightarrow J$ as well as a family of maps

$$f_{i,j}: A_i \otimes B_j \rightarrow B_{\gamma(i,j)}.$$

If for each $j' \in J$, there are only finitely many i and j such that $\gamma(i, j) = j'$ and $f_{i,j} \neq 0$, then there is a well-defined map

$$\sum_{i,j} f_{i,j}: \prod_{i \in I} A_i \rightarrow \prod_{j \in J} B_j,$$

whose component functions are $f_{i,j}$.

Note that both the lattice complex and the power series ring $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]]$ may be viewed as infinite direct products of copies of $\mathbb{F}$. Hence, it is sufficient to show that for each $y = U^n \otimes [K, E]$, there are at most finitely many monomials $a \in \mathcal{R}_\ell$ and generators $x = U^m \otimes [K', E]$ such that $a \cdot x = y$. If $a = \mathcal{U}_1^{i_1} \dots \mathcal{U}_\ell^{i_\ell} \mathcal{V}_1^{j_1} \dots \mathcal{V}_\ell^{j_\ell} \in \mathcal{R}_\ell$ is a monomial, we define $v(a) \in H^2(X_\Gamma)$ by the formula

$$v(a) = (j_1 - i_1)\mu_1^* + \dots + (j_\ell - i_\ell)\mu_\ell^*.$$

Monomials in $\mathcal{R}_\ell$ are equipped with a U -weight

$$w_U(a) = \min(i_1, j_1) + \dots + \min(i_\ell, j_\ell).$$

A generator $U^n \otimes [K, E]$ also has a U -weight $w_U(U^n \otimes [K, E]) = n$. It is straightforward to verify that $w_U(a \cdot x) \geq w_U(a) + w_U(x)$ for any a and x . In particular, if $y = U^n \otimes [K, E]$ is fixed, then there are only finitely many possible U -weights of a and x such that $y = a \cdot x$.

Monomials in $\mathcal{R}_\ell$ also have an Alexander grading $A(a) = (j_1 - i_1, \dots, j_\ell - i_\ell)$. Since $H^2(X_\Gamma)$ is torsion-free and of rank $|L_\Gamma|$, we see that $v(a) = v(a')$ if and only if $A(a) = A(a')$. Also, if $a \cdot [K', E] = U^n \otimes [K, E]$, then necessarily $K' = K - 2v(a)$.

Let $s \in \{1, \dots, \ell\}$. If $A_s(a) = j_s - i_s$ is sufficiently negative, then $\mathcal{V}_s[K - 2v(a), E] = U[K - 2v(a) + 2\mu_s^*, E]$ by the observation that $(K - 2v(a))(v_s) < 0$ and the minima involving $g(K - 2v(a), E)$ and $g(K - 2v(a) + 2\mu_s^*, E)$ will both be attained at some $I \subseteq E$ containing v_s . A similar argument shows that if $A_s(a)$ is sufficiently positive, then $\mathcal{U}_s[K - 2v(a), E] = U[K - 2v(a) - 2\mu_s^*, E]$.

In particular, it follows from the above reasoning that the set of monomials a such that there is an i satisfying $a \cdot U^i \otimes [K - 2v(a), E] = U^n \otimes [K, E]$ is bounded in U -weight and Alexander grading. However, it is easy to verify that the set of elements in $\mathcal{R}_\ell$ with bounded Alexander grading and U -weight is finite, completing the proof. ■

4.5. Maslov gradings

If the intersection form of X_Γ is non-singular, then the lattice complex $\mathbb{CFL}(\Gamma, V_\uparrow)$ inherits a Maslov grading from $\mathbb{CF}(\Gamma)$. We recall the formula

$$\text{gr}(U^i \otimes [K, E]) = -2i + 2g(K, E) + |E| + \frac{1}{4}(K^2 - 3\sigma(X_\Gamma) - 2\chi(X_\Gamma)) \quad (4.6)$$

(see [40, Section 2.3]). Here, $\sigma(X_\Gamma)$ and $\chi(X_\Gamma)$ are the signature and Euler characteristic, respectively. The number K^2 is obtained by pulling back K from $H^2(X_\Gamma)$ to $H^2(X_\Gamma, \partial X_\Gamma)$, squaring using the cup product, and then evaluating on the fundamental class $[X_\Gamma, \partial X_\Gamma] \in H_4(X_\Gamma, \partial X_\Gamma)$.

In the setting of link lattice homology, it is more natural to define the Maslov grading via the formula

$$\text{gr}_w(U^i \otimes [K, E]) = -2i + 2g(K, E) + |E| - |V_\uparrow| + \frac{1}{4}(K|_{X_G}^2 - 3\sigma(X_G) - 2\chi(X_G)). \quad (4.7)$$

This grading is defined when the intersection form Q_G of X_G is non-singular. More generally, this grading may also be defined when Q_G is singular, as long as we restrict to torsion Spin^c structures on Y_G .

Lemma 4.5. *The differential ∂ on $\mathbb{CFL}(\Gamma, V_\uparrow)$ decreases gr_w by 1. Furthermore, if $v_i \in V_\uparrow$, then $\mathcal{V}_i$ preserves gr_w , and $\mathcal{U}_i$ decreases gr_w by 2.*

Proof. The proof that $\text{gr}_w(\partial) = -1$ is essentially identical to the proof of [41, Lemma 3.1] (see also [34]) so we will not repeat it.

We now consider the actions of $\mathcal{V}_i$ and $\mathcal{U}_i$. Note that $(K \pm 2\mu_i^*)|_{X_G}^2 = K|_{X_G}^2$, since μ_i^* has trivial restriction to X_G . Hence the grading changes are entirely due to the powers of U and the $g(K, E)$ terms. With this in mind, the stated grading changes follow immediately from the formulas defining the action of $\mathcal{V}_i$ and $\mathcal{U}_i$ in (4.3) and (4.2). ■

Remark 4.6. Note that gr_w and gr do not in general differ by a constant. Also ∂ is only homogeneously graded on $\mathbb{CFL}(\Gamma, V_\uparrow)$, and not on the entire complex $\mathbb{CF}(\Gamma)$. The reader should compare the lattice complex to the Heegaard Floer mapping cone complex of Ozsváth and Szabó [50]. For $K \subseteq S^3$ with integer framing n , this takes the form of a mapping cone $\mathbb{X}_n(K) = \text{Cone}(v + h_n: \mathbb{A}(K) \rightarrow \mathbb{B}(K))$, which is homotopy equivalent to

$CF^-(S_n^3(K))$. The complex $\mathbb{A}(K)$ is isomorphic to the full knot Floer complex $\mathcal{CFK}(K)$ (a finitely generated chain complex over $\mathbb{F}[\mathcal{U}, \mathcal{V}]$), and hence admits a grading gr_w on $\mathcal{CFK}(K)$. This does not coincide with the Maslov grading on $\mathbb{X}_n(K)$ (see [50, Section 4]) induced by the isomorphism with $CF^-(S_n^3(K))$.

4.6. Alexander gradings

We now define the Alexander multi-grading. If $|V_\uparrow| = \ell$, our Alexander grading will take values in $\mathbb{Q}^\ell$. We assume that Y_G is a rational homology 3-sphere.

If $v \in V_\uparrow$, we may only view v as a class in $H_2(X_G, \partial X_G)$, and not necessarily $H_2(X_G)$. Concretely, $[v] \in H_2(X_G, \partial X_G)$ is obtained by capping one end of the link cobordism $[0, 1] \times L_v \subseteq X_G$ with a disk in S^3 . Since Y_G is a rational homology 3-sphere, the knot $L_v \subseteq Y_G$ is rationally null-homologous, so we may lift the class $[v] \in H_2(X_G, \partial X_G)$ to a class $[\hat{v}] \in H_2(X_G; \mathbb{Q})$. Such a lift is given by capping with a rational Seifert surface for $L_v \subseteq Y_G$.

Definition 4.7. Let Γ be an arrow-decorated plumbing tree, as above, and suppose that $U^i \cdot [K, E]$ is a generator of $\mathbb{CFL}(\Gamma, V_\uparrow)$. Write $V_\uparrow = \{v_1, \dots, v_\ell\}$. We define the *Alexander multi-grading* of $U^i \cdot [K, E]$ to be

$$A(U^i \cdot [K, E]) = \left(\frac{K(v_1 - \hat{v}_1) + \sum_{v \in V_\uparrow} v \cdot (v_1 - \hat{v}_1)}{2}, \dots, \frac{K(v_\ell - \hat{v}_\ell) + \sum_{v \in V_\uparrow} v \cdot (v_\ell - \hat{v}_\ell)}{2} \right),$$

which lies in $\mathbb{Q}^\ell$.

Note that the Alexander multi-grading of $U^i[K, E]$ depends on K , but not on E or i .

Remark 4.8. More generally, if Y_G is not a rational homology 3-sphere, we can construct the Alexander grading A_i when the corresponding component of $L_\uparrow$ is rationally null-homologous, as long as a rational Seifert surface is chosen. For our purposes, such a choice is equivalent to a choice of lift $\hat{v}_i$ of v_i under the map $H_2(X_G; \mathbb{Q}) \rightarrow H_2(X_G, \partial X_G; \mathbb{Q})$. Our grading may be defined in this context as well.

Lemma 4.9. *The differential on $\mathbb{CFL}(\Gamma, V_\uparrow)$ preserves the Alexander multi-grading. Suppose that $v_i, v_j \in V_\uparrow$. The action of $\mathcal{U}_i$ drops the Alexander grading A_j by $\delta_{i,j}$ (Kronecker delta) and the action of $\mathcal{V}_i$ increases the Alexander grading A_j by $\delta_{i,j}$.*

Proof. The A -terms of the differential obviously preserve the Alexander multi-grading since they do not change K . The B -terms send $[K, E]$ to a sum of U -multiples of $[K + 2w^*, E - w]$, where w ranges over the non-arrow components. Note that since $w \in V_G$, w^* is the Poincaré dual of the element $[w] \in H_2(X_G; \mathbb{Z})$, so

$$(K + 2w^*)(v_j - \hat{v}_j) = K(v_j - \hat{v}_j).$$

Hence the B -terms also preserve the Alexander grading.

Next, we consider the actions of $\mathcal{U}_i$ and $\mathcal{V}_i$. We note that $(K + 2\mu_i^*)(\hat{v}_j) = K(\hat{v}_j)$, since $\hat{v}_j$ and μ_i^* are represented by disjoint rational 2-chains. On the other hand, we have

$$(K + 2\mu_i^*)(v_j) = K(v_j) + 2\delta_{i,j}.$$

The conclusion follows from these observations. $\blacksquare$

4.7. Conjugation symmetry

As with the original construction of lattice homology [34, Remark 3.2.7], the link lattice complex admits a conjugation symmetry (see [42, Section 2.2]). This takes the form of a map

$$J(U^i[K, E]) = U^i\left[-K - \sum_{v \in E} 2v^*, E\right].$$

We now observe that the map J is skew Alexander graded:

Lemma 4.10. *The map J satisfies*

$$A(J([K, E])) = -A([K, E]).$$

Proof. By construction of the link lattice complex, $V_\uparrow \subseteq E$. We observe that if $v \in E \setminus V_\uparrow$ and $v_i \in V_\uparrow$, then $2v^*(v_i - \hat{v}_i) = 0$ since v^* is Poincaré dual to a class in $H_2(X_G; \mathbb{Z})$. Hence, if $v_i \in V_\uparrow$, then

$$\begin{aligned} & \left(-K - \sum_{v \in E} 2v^*\right)(v_i - \hat{v}_i) + \sum_{v \in V_\uparrow} v \cdot (v_i - \hat{v}_i) \\ &= \left(-K - \sum_{v \in V_\uparrow} 2v^*\right)(v_i - \hat{v}_i) + \sum_{v \in V_\uparrow} v \cdot (v_i - \hat{v}_i) \\ &= -K(v_i - \hat{v}_i) - \sum_{v \in V_\uparrow} v \cdot (v_i - \hat{v}_i). \end{aligned}$$

Therefore, $A(J([K, E])) = -A([K, E])$. $\blacksquare$

We recall from [40, Section 2.3] that J preserves the grading gr from (4.6). Its interaction with gr_w is more interesting though. Define

$$\text{gr}_z = \text{gr}_w - 2(A_1 + \cdots + A_\ell),$$

where $(A_1, \dots, A_\ell)$ is the Alexander grading. Note that by Lemmas 4.5 and 4.9, the action of $\mathcal{U}_i$ preserves the gr_z -grading, while the action of $\mathcal{V}_i$ drops the gr_z -grading by 2.

Lemma 4.11. *The map J interchanges the gradings gr_w and gr_z .*

Proof. We recall the useful identity (see [40, (2.2)])

$$g(K, E) - g\left(-K - \sum_{v \in E} 2v^*, E\right) = f(K, E). \quad (4.8)$$

Let us write

$$f([K, E]) = \frac{1}{2}(K(v_E) + v_E^2),$$

where $v_E = \sum_{v \in E} v$. Then

$$\begin{aligned} \text{gr}_w([K, E]) - \text{gr}_w(J[K, E]) &= 2f([K, E]) + \frac{1}{4}(K|_{X_G}^2 - (-K - 2v_E^*)|_{X_G}^2) \\ &= K(v_E) + v_E^2 - (K \cup v_E^* + v_E^* \cup v_E^*)[X_G, Y_G]. \end{aligned}$$

Let us write $\hat{v}_E^*$ for the restriction of v_E^* to X_G , pulled back via $H^2(X_G, Y_G) \rightarrow H^2(X_G)$. Use similar notation for individual vertices. Note that if $v_i \in v_\uparrow$, then $v_i^*|_{X_G} = \hat{v}_i^*$, where $\hat{v}_i^* \in H_2(X_G; \mathbb{Q})$ is the 2-chain appearing in the definition of the Alexander grading. Observe that if $v \in E \setminus V_\uparrow$, then $\hat{v}^*$ is still the Poincaré dual of the 2-sphere represented by v , which is contained in X_G . Let us write $v_\uparrow$ for $\sum_{v \in V_\uparrow} v$, and define $\hat{v}_\uparrow$ similarly. We compute that

$$\begin{aligned} f([K, E]) + \frac{1}{4}(K|_{X_G}^2 - (-K - 2v_E^*)|_{X_G}^2) \\ &= K(v_E) + v_E^2 - (K \cup \hat{v}_E^* + \hat{v}_E^* \cup \hat{v}_E^*)[X_G, Y_G] \\ &= K(v_\uparrow - \hat{v}_\uparrow) + v_\uparrow^2 - \hat{v}_\uparrow^2 \\ &= K(v_\uparrow - \hat{v}_\uparrow) + v_\uparrow(v_\uparrow - \hat{v}_\uparrow). \end{aligned}$$

Combining the above with Lemma 4.10 we see that

$$\text{gr}_w([K, E]) = \text{gr}_w(J[K, E]) + 2(A_1 + \cdots + A_\ell)([K, E]) = \text{gr}_z(J[K, E]).$$

The same argument shows

$$\text{gr}_z([K, E]) = \text{gr}_w(J[K, E]). \quad \blacksquare$$

Lemma 4.12. *The map J skew-commutes with $\mathcal{U}_i$ and $\mathcal{V}_i$, i.e.*

$$J \circ \mathcal{V}_i = \mathcal{U}_i \circ J \quad \text{and} \quad J \circ \mathcal{U}_i = \mathcal{V}_i \circ J.$$

Proof. We compute, from (4.2) and (4.3),

$$\begin{aligned} \mathcal{U}_i \cdot J([K, E]) &= U^{1+\delta_i^- (J[K, E])} \left[-K - \sum_{v \in E} 2v^* - 2\mu_i^*, E \right], \\ J(\mathcal{V}_i \cdot [K, E]) &= U^{\delta_i^+ ([K, E])} \left[-K - \sum_{v \in E} 2v^* - 2\mu_i^*, E \right]. \end{aligned}$$

Therefore, it suffices to show that

$$1 + \delta_i^- (J[K, E]) = \delta_i^+ ([K, E]).$$

Rewritten in terms of the g function, the above is equivalent to

$$\begin{aligned} 1 &= g(K + 2\mu_i^*, E) - g(K, E) \\ &\quad + g\left(-K - \sum_{v \in E} 2v^*, E\right) - g\left(-K - 2\mu_i^* - \sum_{v \in E} 2v^*, E\right). \end{aligned}$$

By (4.8), we see that the right-hand side of the above equation is

$$f(K + 2\mu_i^*, E) - f(K, E) = \mu_i^* \left(\sum_{v \in E} v \right) = 1,$$

as claimed. (Recall that $v_i \in E$ by definition of the lattice link complex.) An analogous argument shows that $\mathcal{V}_i \circ J = J \circ \mathcal{U}_i$. ■

4.8. Spin^c structures

We describe how the link lattice complex $\mathbb{CFL}(\Gamma, V_\uparrow)$ naturally splits over $\text{Spin}^c(Y_G)$ as a module over $\mathcal{R}_\ell$. Recall the isomorphism

$$\text{Spin}^c(Y_G) \cong \text{Spin}^c(X_G)/H_2(X_G).$$

Also, the Chern class map $c_1: \text{Spin}^c(X_G) \rightarrow \text{Char}(X_G)$ is an isomorphism of affine $H^2(X_G)$ -sets (where $C \in H^2(X_G)$ acts on $\text{Char}(X_G)$ by $K \mapsto K + 2C$).

In the link lattice complex, we associate the generator $[K, E] \otimes U^i$ with the Spin^c structure $[K]_{X_G}$, viewed as an element of $\text{Char}(X_G)/H_2(X_G) \cong \text{Spin}^c(Y_G)$. This gives the decomposition of $\mathbb{F}[[U]]$ -modules

$$\mathbb{CFL}(\Gamma, V_\uparrow) = \bigoplus_{\mathfrak{s} \in \text{Spin}^c(Y_G)} \mathbb{CFL}(\Gamma, V_\uparrow, \mathfrak{s}). \quad (4.9)$$

Since the differential on $\mathbb{CFL}(\Gamma, V_\uparrow)$ is constructed by modifying (4.1) to sum over only $v \in V_G$, the decomposition (4.9) is preserved by ∂ . The actions of $\mathcal{U}_i$ and $\mathcal{V}_i$ also preserve this decomposition, because they change K to $K \pm 2\mu_i^*$, and $\mu_i^* \in H^2(X_\Gamma)$ has trivial restriction to $H^2(X_G)$. That is, (4.9) yields the decomposition of chain complexes of $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]]$ -modules over the Spin^c structures of Y_G .

4.9. Independence from the framing on arrow components

We now show that our chain complex $\mathbb{CFL}(\Gamma, V_\uparrow)$ is independent of the choice of framing on the arrow components, up to canonical isomorphism.

Suppose that $\mathcal{G}$ is a weighted plumbing tree obtained by weighting the arrow vertices of Γ by (any) integral weights, and using the weights from Γ on V_G . In the previous section, we obtained a model of the link lattice complex which we denote by $\mathbb{CFL}_{\mathcal{G}}(\Gamma, V_\uparrow)$. In this section, we describe a canonical isomorphism

$$F_{\mathcal{G}, \mathcal{G}'}: \mathbb{CFL}_{\mathcal{G}}(\Gamma, V_\uparrow) \rightarrow \mathbb{CFL}_{\mathcal{G}'}(\Gamma, V_\uparrow)$$

for any two weight-extensions $\mathcal{G}$ and $\mathcal{G}'$ of Γ .

Let $L_\Gamma \subseteq S^3$ denote the link associated to Γ as in Section 3.1. We write $L_{\mathcal{G}}$ to denote L_Γ , equipped with the framing from $\mathcal{G}$. Write n for $|V_\Gamma|$. Following the notation of Manolescu and Ozsváth [30], we define the *linking lattice* $\mathbb{H}(L_{\mathcal{G}})$ to be the

affine $\mathbb{Z}^n$ -subspace of $\mathbb{Q}^n$ consisting of the vectors $\mathbf{s} = (s_1, \dots, s_n)$ such that $s_i \in \mathbb{Z} + \text{lk}(L_i, L_\Gamma - L_i)/2$. As sets, we clearly have

$$\mathbb{H}(L_{\mathcal{G}}) = \mathbb{H}(L_{\mathcal{G}'}).$$

The lattices $\mathbb{H}(L_{\mathcal{G}})$ and $\mathbb{H}(L_{\mathcal{G}'})$ are distinguished by their natural actions of $H_2(X_{\mathcal{G}}) \cong H_2(X_{\mathcal{G}'}) \cong \mathbb{Z}^n$. The action of $H_2(X_{\mathcal{G}})$ on $\mathbb{H}(L_{\mathcal{G}})$ is as follows. Given $v \in V_\Gamma$, write λ_v for the longitude of K_v determined by the framing of K_v . By writing $H_1(S^3 \setminus L_\Gamma) \cong \mathbb{Z}^n$, we can identify λ_v as an element in $\mathbb{Z}^n$, and the action of v on $\mathbb{H}(L_{\mathcal{G}})$ can be identified as translation by this element in $\mathbb{Z}^n$.

Next, there is a canonical isomorphism

$$\Phi_{\mathcal{G}}: \text{Char}(X_{\mathcal{G}}) \rightarrow \mathbb{H}(L_{\mathcal{G}})$$

given by the formula

$$\Phi_{\mathcal{G}}(K) = \left(\frac{K(v_1) + v_{\mathcal{G}} \cdot v_1}{2}, \dots, \frac{K(v_n) + v_{\mathcal{G}} \cdot v_n}{2} \right). \quad (4.10)$$

In the above, we write $v_{\mathcal{G}} = v_1 + \dots + v_n \in H_2(X_{\mathcal{G}})$.

Given two weight-extensions $\mathcal{G}$ and $\mathcal{G}'$ of Γ , we define the group isomorphism

$$F_{\mathcal{G}, \mathcal{G}'}: \mathbb{CFL}_{\mathcal{G}}(\Gamma, V_\uparrow) \rightarrow \mathbb{CFL}_{\mathcal{G}'}(\Gamma, V_\uparrow)$$

via the formula

$$F_{\mathcal{G}, \mathcal{G}'}([K, E] \otimes U^i) = [(\Phi_{\mathcal{G}'}^{-1} \circ \Phi_{\mathcal{G}})(K), E] \otimes U^i.$$

The map $F_{\mathcal{G}, \mathcal{G}'}$ is clearly an isomorphism of $\mathbb{F}[[U]]$ -modules. In fact, we have the following:

Proposition 4.13. *The map $F_{\mathcal{G}, \mathcal{G}'}$ is a (gr_w, A) -grading preserving chain isomorphism between $\mathbb{CFL}_{\mathcal{G}}(\Gamma, V_\uparrow)$ and $\mathbb{CFL}_{\mathcal{G}'}(\Gamma, V_\uparrow)$.*

Proof. The differentials on $\mathbb{CFL}_{\mathcal{G}}(\Gamma, V_\uparrow)$ and $\mathbb{CFL}_{\mathcal{G}'}(\Gamma, V_\uparrow)$ are similar to that in (4.1), except that we sum only over the vertices $v \in V_G$. Note that there are no summands in the differential for $v \in V_\uparrow$. We consider the restricted action of $H_2(X_G) \subseteq H_2(X_\Gamma)$ on $\text{Char}(X_{\mathcal{G}})$, $\text{Char}(X_{\mathcal{G}'})$, $\mathbb{H}(L_{\mathcal{G}})$ and $\mathbb{H}(L_{\mathcal{G}'})$. Since the framings of the vertices of V_G coincide on $X_{\mathcal{G}}$ and $X_{\mathcal{G}'}$, the identification $\mathbb{H}(L_{\mathcal{G}}) \cong \mathbb{H}(L_{\mathcal{G}'})$ is equivariant with respect to the action of $H_2(X_G)$.

It is straightforward to verify that the map $\Phi_{\mathcal{G}}$ is equivariant with respect to the action of $H_2(X_G)$. The same argument shows that $\Phi_{\mathcal{G}'}^{-1}$ is equivariant as well.

Let $K \in \text{Char}(X_{\mathcal{G}})$ and write $K' = (\Phi_{\mathcal{G}'}^{-1} \circ \Phi_{\mathcal{G}})(K)$. The assertion of the proposition is a consequence of the following two claims:

- (1) $K|_{X_G} = K'|_{X_G}$.
- (2) For all E , we have $f(K, E) = f(K', E)$.

We verify these two claims presently.

The first claim follows from the proof of [41, Lemma 4.6], which we repeat for the benefit of the reader using our present notation. Write $\mathbf{s} = \Phi_{\mathcal{G}}(K) = \Phi_{\mathcal{G}'}(K')$. Since $H^2(X_G)$ is torsion-free, it suffices to show that $K|_{X_G}(v) = K'|_{X_G}(v)$ for each $v \in V_G$. Note that if $v_i \in V_G$, then $K(v_i) = 2s_i - v_{\mathcal{G}} \cdot v_i$. This quantity only depends on $\mathbf{s}$, the framing of K_i (the link component represented by the vertex v_i), and on the linking numbers of K_i and other link components. In particular, $K(v_i) = K'(v_i)$, completing the proof of the first claim.

We now consider the second claim. By definition,

$$2f(K, E) = K(v_E) + v_E \cdot v_E,$$

where v_E is the sum of v for $v \in E$. We may rearrange the above expression to obtain

$$\begin{aligned} 2f(K, E) &= (v_E - v_{\mathcal{G}}) \cdot v_E + \sum_{v \in E} (K(v) + v_{\mathcal{G}} \cdot v) \\ &= (v_E - v_{\mathcal{G}}) \cdot v_E + \sum_{v_i \in E} s_i. \end{aligned}$$

The right-hand side depends only on $\mathbf{s}$, the framing of L_G , and the linking numbers of the components of $L_{\mathcal{G}}$, but not on the framings of $V_{\uparrow}$. This establishes the second claim.

From these considerations, it follows that $F_{\mathcal{G}, \mathcal{G}'}$ preserves the gr_w -grading, and is also a chain map.

We now establish that $F_{\mathcal{G}, \mathcal{G}'}$ preserves the Alexander grading. Suppose that $v_i \in V_{\uparrow}$. The corresponding component of the Alexander grading is half of

$$\begin{aligned} K(v_i - \hat{v}_i) + \sum_{v \in V_{\uparrow}} v \cdot (v_i - \hat{v}_i) \\ &= K(v_i) + \sum_{v \in V_{\uparrow}} v \cdot v_i - K(\hat{v}_i) - \sum_{v \in V_{\uparrow}} v \cdot \hat{v}_i \\ &= K(v_i) + \sum_{v \in V_{\Gamma}} v \cdot v_i - \sum_{v \in V_G} v \cdot v_i - K(\hat{v}_i) - \sum_{v \in V_{\uparrow}} v \cdot \hat{v}_i. \end{aligned}$$

The first two terms above sum to $2s_i$. The last three terms may be rewritten as follows:

$$- \sum_{v \in V_G} PD[v](v_i) - K(\hat{v}_i) - \sum_{v \in V_{\uparrow}} PD[v](\hat{v}_i).$$

In particular, each term is the evaluation of an element of $H^2(X_{\Gamma})$ on an element of $H_2(X_G; \mathbb{Q})$. We observe that $PD[v]|_{X_G}$, for $v \in \Gamma$, is independent of the framing on $V_{\uparrow}$. Furthermore, $K|_{X_G} = K'|_{X_G}$. Hence $A_i([K, E]) = A_i([K', E])$. ■

4.10. Freeness of the lattice complex

Since $\mathcal{U}_i \mathcal{V}_i = U$ for all i , for $\ell > 1$ the link lattice complex is not free over $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_{\ell}, \mathcal{V}_{\ell}]]$. Nonetheless, we prove that the link lattice complex is free over $\mathbb{F}[[\mathcal{U}_i, \mathcal{V}_i]]$ for each index i .

Proposition 4.14. *Suppose that Γ is an arrow-decorated plumbing graph with a chosen vertex $v_i \in V_\uparrow$. Write $\mathcal{U}_i$ and $\mathcal{V}_i$ for the variables associated to v_i . Then the module $\text{CFL}(\Gamma, V_\uparrow)$ is the completion of a free module over $\mathbb{F}[[\mathcal{U}_i, \mathcal{V}_i]]$.*

Proof. We consider the set of generators $[K, E]$ modulo the equivalence relation generated by $[K, E] \sim [K + 2\mu_i^*, E]$ for all K and E . Equivalence classes may be identified with elements of $(\text{Char}(\Gamma) \times \mathbb{P}(V_G))/\mathbb{Z}$, where $1 \in \mathbb{Z}$ acts on $\text{Char}(\Gamma)$ by $2\mu_i^*$.

Fix an equivalence class, and let W denote the $\mathbb{F}[U]$ -span of the generators in this class. We may write $W \cong \bigoplus_{s \in \mathbb{Z}} W_s$ where $W_s \cong \mathbb{F}[U]$ for each s . We order the W_s so that $\mathcal{V}_i \cdot W_s \subseteq W_{s+1}$ and $\mathcal{U}_i \cdot W_s \subseteq W_{s-1}$. We make the following claims, from which the result will follow fairly easily:

- (f-1) For each s , exactly one of $\mathcal{V}_i: W_s \rightarrow W_{s+1}$ and $\mathcal{U}_i: W_{s+1} \rightarrow W_s$ is multiplication by 1, and the other is multiplication by U . This follows from the fact that $\mathcal{U}_i \cdot \mathcal{V}_i$ acts by U .
- (f-2) If $[K, E]$ is fixed, then $[K, E]$ is not in the image of $\mathcal{U}_i^n$ or $\mathcal{V}_i^n$ for arbitrarily large n . This follows immediately from Lemma 4.4.
- (f-3) If $\mathcal{V}_i \cdot [K, E] = [K + 2\mu_i^*, E]$, then $\mathcal{V}_i \cdot [K + 2\mu_i^*, E] = [K + 4\mu_i^*, E]$. Similarly, if $\mathcal{U}_i \cdot [K, E] = [K - 2\mu_i^*, E]$ then $\mathcal{U}_i \cdot [K - 2\mu_i^*, E] = [K - 4\mu_i^*, E]$.

We now prove claim (f-3), focusing on the argument for $\mathcal{V}_i$ since the claim about $\mathcal{U}_i$ is similar. We recall from Section 4.4 that

$$f(K + 2\mu_i^*, E) = \begin{cases} f(K, E) & \text{if } v_i \notin E, \\ f(K, E) + 1 & \text{if } v_i \in E. \end{cases}$$

Hence, $\delta_i^+(K, E) = 0$ if and only if there is a $J \subseteq E$ such that $f(K, J) = g(K, E)$ and $v_i \notin J$. In particular, if $\mathcal{V}_i \cdot [K, E] = [K + 2\mu_i^*, E]$, then there exists such a J . Hence

$$g(K, E) = f(K, J) = f(K + 4\mu_i^*, J) \geq g(K + 4\mu_i^*, E) \geq g(K, E),$$

so we have equality throughout. It follows that $\mathcal{V}_i \cdot [K + 2\mu_i^*, E] = [K + 4\mu_i^*, E]$.

Note that claim (f-3) implies there is no $[K, E]$ which is in the image of both $\mathcal{U}_i$ and $\mathcal{V}_i$, since if $[K, E]$ were in the image of both, then the above claims show that

$$\mathcal{U}_i \mathcal{V}_i [K - 2\mu_i^*, E] = \mathcal{U}_i [K, E] = [K - 2\mu_i^*, E],$$

which contradicts $\mathcal{U}_i \mathcal{V}_i = U$.

Claims (f-1) and (f-2) imply that there exist generators $[K, E]$ in W which are in the image of $\mathcal{U}_i$, and there also exist generators which are in the image of $\mathcal{V}_i$. (This rules out the modules $\mathbb{F}[\mathcal{U}_i, \mathcal{V}_i, \mathcal{V}_i^{-1}]$ and $\mathbb{F}[\mathcal{U}_i, \mathcal{U}_i^{-1}, \mathcal{V}_i]$.)

From the above considerations, we deduce that there is a unique generator $[K, E]$ such that $\mathcal{U}_i [K, E] = [K - 2\mu_i^*, E]$ and $\mathcal{V}_i [K, E] = [K + 2\mu_i^*, E]$. By (f-3) this $[K, E]$ must be a free generator of W over $\mathbb{F}[\mathcal{U}_i, \mathcal{V}_i]$. ■

4.11. Type- D modules over $\mathcal{K}$

If G is a weighted plumbing tree (without arrow vertices) and v is a distinguished vertex, we now describe how to view the lattice complex as a type- D module over the algebra $\mathcal{K}$, described by the third author [60].

We first recall the algebra $\mathcal{K}$ from [60]. It is an algebra over the idempotent ring $\mathbf{I} \cong \mathbf{I}_0 \oplus \mathbf{I}_1$, where $\mathbf{I}_\varepsilon \cong \mathbb{F}$ for $\varepsilon = 0, 1$. We define

$$\mathbf{I}_0 \cdot \mathcal{K} \cdot \mathbf{I}_0 \cong \mathbb{F}[\mathcal{U}, \mathcal{V}], \quad \mathbf{I}_0 \cdot \mathcal{K} \cdot \mathbf{I}_1 = 0 \quad \text{and} \quad \mathbf{I}_1 \cdot \mathcal{K} \cdot \mathbf{I}_1 \cong \mathbb{F}[\mathcal{U}, \mathcal{V}, \mathcal{V}^{-1}].$$

Finally, $\mathbf{I}_1 \otimes \mathcal{K} \otimes \mathbf{I}_0$ is isomorphic to the direct sum of two copies of $\mathbb{F}[\mathcal{U}, \mathcal{V}, \mathcal{V}^{-1}]$, viewed as being generated by two distinguished elements σ and τ . These elements satisfy

$$\begin{aligned} \sigma \cdot \mathcal{U} &= U \mathcal{V}^{-1} \cdot \sigma, & \sigma \cdot \mathcal{V} &= \mathcal{V} \cdot \sigma, \\ \tau \cdot \mathcal{U} &= \mathcal{V}^{-1} \cdot \tau, & \tau \cdot \mathcal{V} &= U \mathcal{V} \cdot \tau, \end{aligned}$$

where $U = \mathcal{U} \mathcal{V}$.

In this section, we describe how to construct a type- D module $\mathcal{X}(G)^\mathcal{K}$ from the data of $\mathbb{C}\mathbb{F}(G)$. The construction of $\mathcal{X}(G)^\mathcal{K}$ may be equivalently described as a tensor product of the Hopf, merge and solid torus modules from [60], though we presently give a direct construction in terms of lattices. We define $\mathcal{X}(G)^\mathcal{K}$ at the end of the section, after proving several properties of the lattice complex.

Write $\mathbb{C}\mathbb{F}_0(G)$ for the codimension 1 subcube of $\mathbb{C}\mathbb{F}(G)$ generated by the tuples $[K, E]$ where $v \in E$. Write $\mathbb{C}\mathbb{F}_1(G)$ for the codimension 1 subcube generated by the tuples $[K, E]$ where $v \notin E$. We view $\mathbb{C}\mathbb{F}(G)$ as a mapping cone

$$\mathbb{C}\mathbb{F}(G) \cong \text{Cone}(F_v^A + F_v^B: \mathbb{C}\mathbb{F}_0(G) \rightarrow \mathbb{C}\mathbb{F}_1(G)),$$

where F_v^A and F_v^B are the summands of the differential which are weighted by $U^{a_v(K, E)}$ and $U^{b_v(K, E)}$, respectively.

We observe that $\mathbb{C}\mathbb{F}_0(G)$ is exactly the link lattice complex if we designate the special vertex v as the sole arrow vertex. In particular, Proposition 4.14 implies that it is a completion of a free $\mathbb{F}[[\mathcal{U}, \mathcal{V}]]$ -module (where $\mathcal{U}$ and $\mathcal{V}$ are the actions for v).

We may define actions of $\mathcal{U}$ and $\mathcal{V}$ also on $\mathbb{C}\mathbb{F}_1(G)$, using the same formulas as in Section 4.4. We first observe that the formulas have a comparatively easier description than on $\mathbb{C}\mathbb{F}_0(G)$:

Lemma 4.15. *On $\mathbb{C}\mathbb{F}_1(G)$, we have*

$$\mathcal{U} \cdot [K, E] = U[K - 2\mu^*, E] \quad \text{and} \quad \mathcal{V} \cdot [K, E] = [K + 2\mu^*, E]$$

for all K and E such that $v \notin E$.

Proof. If $v \notin E$, then $f(K, E) = f(K \pm 2\mu^*, E)$, and hence

$$g(K, E) = g(K \pm 2\mu^*, E). \tag{4.11}$$

The desired equalities follow by applying this fact to the definition of $\mathcal{U}$ and $\mathcal{V}$ from Section 4.4. ■

As a consequence of the above, we may define an action of $\mathcal{V}^{-1}$ on $\mathbb{C}\mathbb{F}_1(G)$ by the formula $\mathcal{V}^{-1} \cdot [K, E] = [K - 2\mu^*, E]$. As an additional consequence of Lemma 4.15, we have the following easy analog of Proposition 4.14:

Corollary 4.16. *The complex $\mathbb{C}\mathbb{F}_1(G)$ is the completion of a free module over $\mathbb{F}[\mathcal{U}, \mathcal{V}, \mathcal{V}^{-1}]$.*

Let F_v^A denote the A -term of the differential which involves v . Write F_v^B for the B -term of the differential which involves v .

Lemma 4.17. *The map F_v^A satisfies*

$$F_v^A(\mathcal{U} \cdot \mathbf{x}) = \mathcal{U} \cdot F_v^A(\mathbf{x}) \quad \text{and} \quad F_v^A(\mathcal{V} \cdot \mathbf{x}) = \mathcal{V} \cdot F_v^A(\mathbf{x})$$

for all $\mathbf{x} \in \mathbb{C}\mathbb{F}_0(G)$. Similarly,

$$F_v^B(\mathcal{U} \cdot \mathbf{x}) = \mathcal{V}^{-1} \cdot F_v^B(\mathbf{x}) \quad \text{and} \quad F_v^B(\mathcal{V} \cdot \mathbf{x}) = \mathcal{V}U \cdot F_v^B(\mathbf{x}).$$

Proof. We begin with the map F_v^A . From direct computation,

$$\begin{aligned} \mathcal{V} \cdot F_v^A([K, E]) &= U^{g(K+2\mu^*, E-v)-g(K, E)} \cdot [K + 2\mu^*, E - v] \\ &= F_v^A(\mathcal{V} \cdot [K, E]). \end{aligned}$$

An entirely analogous argument shows that $[F_v^A, \mathcal{U}] = 0$.

Next, we consider the commutation of F_v^B with $\mathcal{U}$. We compute that

$$\begin{aligned} F_v^B(\mathcal{U} \cdot [K, E]) &= U^{g(K+2v^*-2\mu^*, E-v)-g(K, E)+1+(K-2\mu^*)(v)/2+v^2/2} \cdot [K - 2\mu^* + 2v^*, E - v] \\ &= \mathcal{V}^{-1} \cdot U^{g(K+2v^*, E-v)-g(K, E)+K(v)/2+v^2/2} \cdot [K + 2v^*, E - v] \\ &= \mathcal{V}^{-1} \cdot F_v^B([K, E]). \end{aligned} \tag{4.12}$$

Going from the second line to the third, we are using (4.11).

Next, we consider the commutation of F_v^B with $\mathcal{V}$. We compute

$$\begin{aligned} F_v^B(\mathcal{V} \cdot [K, E]) &= U^{g(K+2\mu^*+2v^*, E-v)-g(K, E)+(K+2\mu^*)(v)/2+v \cdot v/2} \cdot [K + 2\mu^* + 2v^*, E - v] \\ &= U \mathcal{V} \cdot (U^{g(K+2v^*, E-v)-g(K, E)+K(v)/2+v \cdot v/2} \cdot [K + 2v^*, E - v]) \\ &= U \mathcal{V} \cdot F_v^B([K, E]), \end{aligned} \tag{4.13}$$

where we again use (4.11). The proof is complete. $\blacksquare$

We are now able to define the type- D module $\mathcal{X}(G)^{\mathcal{X}}$. As a right $\mathbf{I}$ -module, we write $\mathcal{X}(G) = \mathbb{X}_0 \oplus \mathbb{X}_1$, where each $\mathbb{X}_\varepsilon$ is a vector space over $\mathbb{F}$. We define $\mathbb{X}_0$ to be the $\mathbb{F}$ -vector space generated by a free $\mathbb{F}[\mathcal{U}, \mathcal{V}]$ -basis of $\mathbb{C}\mathbb{F}_0(G)$ from Proposition 4.14. We

define $\mathbb{X}_1$ to be the $\mathbb{F}$ -vector space generated by a free $\mathbb{F}[\mathcal{U}, \mathcal{V}, \mathcal{V}^{-1}]$ -basis of $\mathbb{C}\mathbb{F}_1(G)$ from Corollary 4.16.

We now define the structure map

$$\delta^1: \mathcal{X}(G) \rightarrow \mathcal{X}(G) \otimes_{\mathbf{I}} \mathcal{K}.$$

The construction is entirely analogous to the setting of the link surgery formula; see [60, Section 8.5] for a parallel construction. If $\mathbf{x}$ is a basis element of $\mathbb{C}\mathbb{F}_0(G)$ and $\partial(\mathbf{x})$ has a summand $\mathcal{U}^i \mathcal{V}^j \cdot \mathbf{y}$, where $\mathbf{y}$ is a basis element, then we define $\delta^1(\mathbf{x})$ to have a summand $\mathbf{y} \otimes \mathcal{U}^i \mathcal{V}^j$. We make a similar definition for basis elements of $\mathbb{C}\mathbb{F}_1(G)$. Next, if $F_v^A(\mathbf{x}) = \mathcal{U}^i \mathcal{V}^j \cdot \mathbf{y}$, we declare $\delta^1(\mathbf{x})$ to also have a summand $\mathbf{y} \otimes \mathcal{U}^i \mathcal{V}^j \sigma$. Similarly, if $F_v^B(\mathbf{x}) = \mathcal{U}^i \mathcal{V}^j \cdot \mathbf{y}$, then we declare $\delta^1(\mathbf{x})$ to have a summand $\mathbf{y} \otimes \mathcal{U}^i \mathcal{V}^j \tau$. It is straightforward to verify that $\mathcal{X}(G)^{\mathcal{K}}$ satisfies the type- D structure relations.

We note that the underlying vector space of $\mathcal{X}(G)^{\mathcal{K}}$ is infinite-dimensional, so completions play a subtle yet important role in the theory. We leave it to the reader to verify that the modules satisfy the *Alexander module* condition described in [60, Section 6].

4.12. An example

In this subsection, we compute the link lattice homology of $T(2, 2)$, the positive Hopf link. The Hopf link in S^3 can be represented by the plumbing graph Γ with one solid and two arrow vertices, together with two edges connecting the solid vertex to the two arrow vertices. The solid vertex has weight -1 , and we assign -3 and -2 to the arrow vertices. Denote the solid vertex by v_0 and the two arrow vertices by v_1, v_2 . The link lattice complex $\mathbb{C}\mathbb{F}\mathbb{L}(\Gamma, V_\Gamma)$ is generated by the elements $[K, E_1]$ and $[K, E_2]$ as an $\mathbb{F}[[U]]$ -module where $K \in \text{Char}(\Gamma)$ and

$$E_1 = \{v_1, v_2\} \quad \text{and} \quad E_2 = \{v_0, v_1, v_2\}.$$

It is not hard to see that

$$\text{Char}(\Gamma) = \{K = [2n + 1, 2m_1 + 3, 2m_2 + 2]; n, m_1, m_2 \in \mathbb{Z}\},$$

where writing $K = [2n + 1, 2m_1 + 3, 2m_2 + 2]$ means that $K(v_0) = 2n + 1$, $K(v_1) = 2m_1 + 3$, $K(v_2) = 2m_2 + 2$. We compute the differentials:

$$\partial[K, E_1] = 0,$$

$$\partial[K, E_2] = U^{a_{v_0}[K, E_2]} \otimes [K, E_1] + U^{b_{v_0}[K, E_2]} \otimes [K + 2v_0^*, E_1].$$

By direct computations,

$$g(K, E_2) = \min\{0, n, m_1, m_2, m_1 + n + 1, m_2 + n + 1, m_1 + m_2, m_1 + m_2 + n + 2\},$$

$$A_{v_0}(K, E_2) = \min\{0, m_1, m_2, m_1 + m_2\},$$

$$B_{v_0}(K, E_2) = \min\{n, m_1 + n + 1, m_2 + n + 1, m_1 + m_2 + n + 2\}.$$

Hence

$$\begin{aligned} \partial[K, E_2] &= \begin{cases} [K, E_1] + U^n[K + 2v_0^*, E_1] & \text{if } n \geq 0, m_1 \geq 0, m_2 \geq 0, \\ [K, E_1] + U^{n+1}[K + 2v_0^*, E_1] & \text{if } n \geq 0, m_1 < 0, m_2 \geq 0 \text{ or } m_1 \geq 0, m_2 < 0, \\ [K, E_1] + U^{n+2}[K + 2v_0^*, E_1] & \text{if } n \geq 0, m_1 < 0, m_2 < 0, \\ U^{-n}[K, E_1] + [K + 2v_0^*, E_1] & \text{if } n < 0, m_1 \geq 0, m_2 \geq 0, \\ U^{-n-1}[K, E_1] + [K + 2v_0^*, E_1] & \text{if } n < 0, m_1 < 0, m_2 \geq 0 \text{ or } m_1 \geq 0, m_2 < 0, \\ U^{-n-2}[K, E_1] + [K + 2v_0^*, E_1] & \text{if } n < -1, m_1 < 0, m_2 < 0. \\ [K, E_1] + U[K + 2v_0^*, E_1] & \text{if } n = -1, m_1 < 0, m_2 < 0. \end{cases} \end{aligned} \quad (4.14)$$

Therefore, the link lattice homology $\mathbb{HFL}(\Gamma, V_\uparrow)$ is generated as a group by the elements $[K, E_1]$ where $K = [-3, 2m_1 + 3, 2m_2 + 2]$ for $m_1 < 0, m_2 < 0$ and $K = [-1, 2m_1 + 3, 2m_2 + 2]$ otherwise. We now consider the module structure, that is, the actions of $\mathcal{U}_1, \mathcal{U}_2, \mathcal{V}_1, \mathcal{V}_2$ on these elements. By (4.4),

$$\begin{aligned} \mathcal{U}_i \cdot [K, E_1] &= U^{g(K-2\mu_i^*, E_1)-g(K, E_1)+1} [K - 2\mu_i^*, E_1], \\ \mathcal{V}_i \cdot [K, E_1] &= U^{g(K+2\mu_i^*, E_1)-g(K, E_1)} [K + 2\mu_i^*, E_1]. \end{aligned}$$

Suppose $K = [2n + 1, 2m_1 + 3, 2m_2 + 2]$. Then

$$\begin{aligned} g(K, E_1) &= \min \{0, m_1, m_2, m_1 + m_2\}, \\ g(K \pm 2\mu_1^*, E_1) &= \min \{0, m_1 \pm 1, m_2, m_1 \pm 1 + m_2\}. \end{aligned}$$

Similarly,

$$g(K \pm 2\mu_2^*, E_1) = \min \{0, m_1, m_2 \pm 1, m_1 + m_2 \pm 1\}.$$

Hence, for $i = 1$ or 2 , we have

$$\begin{aligned} \mathcal{U}_i[K, E_1] &= \begin{cases} U[K - 2\mu_i^*, E_1] & \text{if } m_i > 0, \\ [K - 2\mu_i^*, E_1] & \text{if } m_i \leq 0, \end{cases} \\ \mathcal{V}_i[K, E_1] &= \begin{cases} U[K + 2\mu_i^*, E_1] & \text{if } m_i < 0, \\ [K + 2\mu_i^*, E_1] & \text{if } m_i \geq 0. \end{cases} \end{aligned}$$

It follows that when $m_1 = m_2 = 0$ and $m_1 = m_2 = -1$ the corresponding generators $[K, E_1]$ are in the image of neither $\mathcal{U}_i$ nor $\mathcal{V}_i$. In particular, the lattice link homology of $T(2, 2)$ is generated over $\mathbb{F}[\mathcal{U}_1, \mathcal{U}_2, \mathcal{V}_1, \mathcal{V}_2]$ by the elements

$$\mathbf{X} := [K_1, E_1] \quad \text{and} \quad \mathbf{Y} := [K_2, E_1],$$

where $K_1 := [-1, 3, 2]$ and $K_2 := [-3, 1, 0]$.

Using Definition 4.7, one easily computes

$$A(\mathbf{X}) = (0, 0) \quad \text{and} \quad A(\mathbf{Y}) = (-1, -1).$$

Additionally, since $\chi(X_G) = 2$ and $\sigma(X_G) = -1$, we compute using (4.7) that

$$\begin{aligned} \text{gr}_w(\mathbf{X}) &= \frac{1}{4}(K|_{X_G}^2 - 3\sigma(X_G) - 2\chi(X_G)) = \frac{1}{4}(K|_{X_G}^2 - 1) = 0, \\ \text{gr}_w(\mathbf{Y}) &= -4 + \frac{1}{4}(K|_{X_G}^2 - 1) = -4 + 2 = -2. \end{aligned}$$

Finally, if $i \neq j$, we have the following equalities as elements of homology:

$$\mathcal{U}_i \cdot \mathbf{X} = U[K_1 - 2\mu_i^*, E_1] = [K_1 + 2v_0^* - 2\mu_i^*, E_1] = \mathcal{V}_j \cdot \mathbf{Y}.$$

Comparing the gradings we see that these two relations (corresponding to $(i, j) = (1, 2), (2, 1)$) generate all relations between the homology classes of $\mathbf{X}$ and $\mathbf{Y}$. In Theorem 7.3, we compute the link Floer homology of $T(n, n)$. The readers can verify that the above description coincides with the link Floer homology of $T(2, 2)$ (see Figure 1.1 for the link Floer complex).

Using these techniques, it is also possible to compute the H -function of $T(2, 2)$. By using (4.7), one can compute $\text{gr}_w([K_{m_1, m_2}, E_1])$ for $m_1, m_2 \in \mathbb{Z}$ and $K_{m_1, m_2} = [-3, 2m_1 + 3, 2m_2 + 2]$ if $m_1, m_2 < 0$, $K_{m_1, m_2} = [-1, 2m_1 + 3, 2m_2 + 2]$ otherwise. The function

$$(a, b) \mapsto -\frac{1}{2} \text{gr}_w([K_{a-1/2, b-1/2}, E_1])$$

agrees with the H -function for the Hopf link; see Figure 6.1.

5. The equivalence with link Floer homology

In this section, we prove that link lattice homology and link Floer homology are isomorphic. The argument applies similar logic to the case of 3-manifolds [41, 59].

Theorem 5.1. *Suppose that Γ is an arrow-decorated plumbing tree with vertex set $V_G \cup V_\uparrow$ such that Y_G is a rational homology 3-sphere. For each $\mathfrak{s} \in \text{Spin}^c(Y_G)$, there is an absolutely (gr_w, A) -graded isomorphism of A_∞ -modules over $\mathcal{R}_\ell$,*

$$\mathbb{CFL}(\Gamma, V_\uparrow, \mathfrak{s}) \simeq \mathcal{CFL}(Y_G, L_\uparrow, \mathfrak{s}).$$

Here, both $\mathbb{CFL}$ and $\mathcal{CFL}$ are equipped with the natural A_∞ -module structures which have only m_1 and m_2 non-trivial.

The proof of Theorem 5.1 is completed in Section 5.3. We now provide a sketch of the proof. We will use a relative version of the Manolescu–Ozsváth link surgery formula [30], which computes link Floer homology as a subcube of the full link surgery hypercube. This is stated in Theorem 5.2. Subsequently, we follow the approach of [59] and view L_Γ as a connected sum of Hopf links. Using a tensor product formula from [60] for the link surgery complex, one obtains a combinatorial model for the link surgery complex of L_Γ .

Following the approach of Ozsváth, Stipsicz and Szabó [41], one may identify the lattice complex with a simplified version of the link surgery hypercube obtained by taking the homology of the link surgery complex at each vertex of the cube $\{0, 1\}^\ell$, and using only the length 1 maps of the link surgery hypercube. In [59], the third author shows directly using the connected sum formula for the link surgery formula that the link surgery complex for L_Γ is chain homotopy equivalent to this simplified model of the link surgery complex. When $b_1(Y_G, \mathbb{Q}) = 0$, we show that this homotopy equivalence induces a homotopy equivalence between the link lattice complex and the corresponding quotient complex of the link surgery complex of L_Γ . We show that the morphisms in this homotopy equivalence are well-behaved with respect to the actions of $\mathcal{U}_i$ and $\mathcal{V}_i$, and give a homotopy equivalence of A_∞ -modules.

5.1. The link surgery complex and sublinks

As a first step, we describe a refinement of the Manolescu and Ozsváth link surgery formula. If $L \subseteq S^3$ is a link equipped with integral framing Λ , then Manolescu and Ozsváth construct a chain complex $\mathcal{C}_\Lambda(L)$ over $\mathbb{F}[[U]]$ (defined in terms of the link Floer complex $\mathcal{CFL}(S^3, L)$, equipped with additional data) and prove that

$$H_*(\mathcal{C}_\Lambda(L)) \cong HF^-(S_\Lambda^3(L)).$$

There is a refinement of this result which can be used to compute link Floer homology, as follows. Suppose that $M = J \cup L \subseteq S^3$ is a partitioned link with $|M| = n$ and $|L| = \ell$. Equip J with a framing Λ . We may extend Λ arbitrarily to a framing Λ' on all of M to obtain a link surgery complex $\mathcal{C}_{\Lambda'}(M)$ whose homology is $HF^-(S_{\Lambda'}^3(M))$.

The link surgery complex $\mathcal{C}_{\Lambda'}(M)$ is a *hypercube* of chain complexes, which means that it admits a natural filtration by the integral points of the cube $\{0, 1\}^n$, where $n = |M|$. While more details are given in [30, Section 5], we give some necessary background and introduce the notation. For any $\varepsilon \in \{0, 1\}^n$, we consider the multiplicatively closed subset $S_\varepsilon \subseteq \mathcal{R}_n$ generated by $\mathcal{V}_i$ such that $\varepsilon(i) = 1$. The complex $\mathcal{C}_\varepsilon$ is defined as the algebraic completion of the localization $S_\varepsilon^{-1} \cdot \mathcal{CFL}(M)$. We remark that the original definition in [30] is seemingly different, though the equivalence with the above description follows from [60, Lemma 5.7].

If $\varepsilon, \varepsilon' \in \{0, 1\}^n$, we write $\varepsilon \leq \varepsilon'$ if $\varepsilon_i \leq \varepsilon'_i$ for all i , and $\varepsilon < \varepsilon'$ if $\varepsilon \leq \varepsilon'$ and $\varepsilon \neq \varepsilon'$. If $\varepsilon, \varepsilon' \in \{0, 1\}^n$ and $\varepsilon < \varepsilon'$, Manolescu and Ozsváth construct a map $D_{\varepsilon, \varepsilon'}: \mathcal{C}_\varepsilon \rightarrow \mathcal{C}_{\varepsilon'}$ (see [30, Section 5]). The chain complex $\mathcal{C}_{\Lambda'}(M)$ is the direct sum of the complexes $\mathcal{C}_\varepsilon$ and the differential is the sum of the internal differentials in $\mathcal{C}_\varepsilon$ and of the maps $D_{\varepsilon, \varepsilon'}$.

We now describe a relative complex $\mathcal{C}_\Lambda(J, L)$. Note that each axis direction in $\{0, 1\}^n$ corresponds to a component of M . Hence, we may consider the quotient complex $\mathcal{C}_\Lambda(J, L)$ obtained by quotienting by the subcomplex of $\mathcal{C}_{\Lambda'}(M)$ consisting of those $\mathcal{C}_\varepsilon$ such that $\varepsilon(i) = 1$ for at least one index i corresponding to L . Examining Manolescu and Ozsváth's construction, it is evident that $\mathcal{C}_\Lambda(J, L)$ is independent of the framings of the L components.

Furthermore, the module $\mathcal{C}_\Lambda(J, L)$ has a natural action of the ring $\mathcal{R}_\ell$, corresponding to the variables for L . The underlying spaces $\mathcal{C}_\varepsilon$ are preserved by this $\mathcal{R}_\ell$ -module structure. It follows from [60, Lemma 5.9] that the hypercube maps of $\mathcal{C}_\Lambda(J, L)$ commute with the action of $\mathcal{R}_\ell$, i.e. the action of the variables from L . Note that in general the differential on $\mathcal{C}_\Lambda(J, L)$ will *not* commute with the actions of the variables from J . The next result is important for our purposes:

Theorem 5.2. *Suppose that $M \subseteq S^3$ is a link which is partitioned into two sublinks, $M = J \cup L$. Let Λ be an integral framing on J and write $\ell = |L|$. Then there is a homotopy equivalence of chain complexes over $\mathcal{R}_\ell$:*

$$\mathcal{CFL}(S_\Lambda^3(J), L) \simeq \mathcal{C}_\Lambda(J, L).$$

Furthermore, this isomorphism respects the Spin^c structures under an isomorphism

$$\text{Spin}^c(S_\Lambda^3(J)) \cong \mathbb{H}(M)/(\text{Span}(\mu_1^*, \dots, \mu_\ell^*) + H_2(W_\Lambda(J))),$$

where $W_\Lambda(J)$ is the cobordism from S^3 to the surgery on the link J .

The above is a folklore result. We believe the techniques of [30] can be adapted in a straightforward manner to prove this theorem. Nonetheless, experts in Heegaard Floer surgery formulas may recognize that although conceptually simple, a rigorous proof requires a substantial amount of bookkeeping because of the role of algebraic truncations in the surgery formula. A conceptually simple proof, avoiding truncations, can be found in [58, Corollary 9.2].

5.2. Link lattice homology and the link surgery formula

We now describe how to recast the link lattice complex in terms of the link surgery complex. This is an adaptation of [41, Proposition 4.4] to our present context of links.

Construct an $|L_G|$ -dimensional hypercube of chain complexes as follows: For $\varepsilon \in \{0, 1\}^{|L_G|}$, define $Z_\varepsilon := H_*(\mathcal{C}_\varepsilon)$, where $\mathcal{C}_\varepsilon$ is the corresponding submodule of $\mathcal{C}_\Lambda(L_G, L_\uparrow)$. If $\varepsilon < \varepsilon'$, define a hypercube map $\delta_{\varepsilon, \varepsilon'}: Z_\varepsilon \rightarrow Z_{\varepsilon'}$ by

$$\delta_{\varepsilon, \varepsilon'} := \begin{cases} (D_{\varepsilon, \varepsilon'})_* & \text{if } |\varepsilon' - \varepsilon|_{L^1} = 1, \\ 0 & \text{otherwise.} \end{cases}$$

Write $\mathcal{Z} = (Z_\varepsilon, \delta_{\varepsilon, \varepsilon'})_{\varepsilon \in \{0, 1\}^{|L_G|}}$. Clearly $\mathcal{Z}$ is a hypercube of chain complexes over $\mathcal{R}_\ell$.

Compare the following with [41, Proposition 4.4]:

Proposition 5.3. *Let Γ be an arrow-decorated plumbing tree. The hypercube $\mathcal{Z} = (Z_\varepsilon, \delta_{\varepsilon, \varepsilon'})_{\varepsilon \in \{0, 1\}^{|L_G|}}$ is isomorphic to the link lattice complex $\mathbb{CFL}(\Gamma, V_\uparrow)$.*

Before proving Proposition 5.3, we prove a technical lemma which is helpful in relating the $\mathcal{R}_\ell$ -actions on $\mathbb{CFL}(\Gamma, V_\uparrow)$ and $\mathcal{Z}$. We note that the following lemma is essentially implicit in the definition of lattice homology and also Ozsváth, Stipsicz, and Szabó's construction of the spectral sequence (see [41, Proposition 4.4]), though we have been unable to find an exact reference which is suitable for our purposes. If $L \subseteq S^3$ and $\mathbf{s} \in \mathbb{H}(L)$,

write

$$d(L, \mathbf{s}) = \max \{ \text{gr}_w(x) : x \in H_* \mathfrak{A}^-(L, \mathbf{s}), x \text{ is } U\text{-non-torsion} \}.$$

Here, gr_w denotes the internal Maslov grading from link Floer homology, and $\mathfrak{A}^-(L, \mathbf{s}) \subseteq \mathcal{CFL}(S^3, L)$ is a subcomplex corresponding to the Alexander grading $\mathbf{s}$.

Remark 5.4. For an oriented link $L \subseteq S^3$ in the 3-sphere, Gorsky and Némethi [11] defined a link invariant, called the H -function, by declaring $-2H_L(\mathbf{s}) = d(L, \mathbf{s})$. The H -function is a generalization of the h_k -invariant considered by Rasmussen [53, Section 7]. For algebraic links, it can be related to the semigroup counting function [11, Section 3.5]. We later generalize the H -function to links in rational homology spheres; see Section 6.2.

Lemma 5.5. *Let G be a forest of plumbing trees. If $E \subseteq V_G$, write $L_E \subseteq S^3$ for the sublink of L_G containing exactly the components corresponding to vertices of E . Let $\phi_E: \text{Char}(X_G) \rightarrow \mathbb{H}(L_E)$ be the composition of the restriction map from $\text{Char}(X_G)$ to $\text{Char}(X_E)$ and the isomorphism from $\text{Char}(X_E)$ to $\mathbb{H}(L_E)$ in (4.10). Then*

$$2g(K, E) = d(L_E, \phi_E(K)).$$

Proof. Note that $g(K, E)$ (computed in X_G) is the same as $g(K|_{X_E}, E)$ (computed in X_E). Hence, we may assume without loss of generality that $E = V_G$.

Next, we observe that the framings on the components of L_G play no role in the statement. Indeed, if $K \in \text{Char}(X_G)$ and $\Phi_G(K) = (s_1, \dots, s_n)$, then (4.10) implies that

$$2g(K, V_G) = \min_{I \subseteq V_G} \left(\sum_{i \in I} 2s_i - v_{G-I} \cdot v_I \right), \quad (5.1)$$

where v_{G-I} is the sum of v_i for $i \in V_G \setminus I$ and v_I is similar. Our proof will be by induction on the number of vertices. If $\mathbf{s} \in \mathbb{H}(L_G)$, we will write $2g(L_G, \mathbf{s})$ for the quantity on the right-hand side of (5.1). Similarly, if $I \subseteq V_G$ we write

$$2f(\mathbf{s}, I) = \sum_{i \in I} 2s_i - v_{G-I} \cdot v_I. \quad (5.2)$$

We claim that $2g(L_G, \mathbf{s}) = d(L_G, \mathbf{s})$.

We begin with the case that L_G is an n -component unlink $\mathbb{U}_n$. In this case, the homology $\mathcal{HFL}(\mathbb{U}_n) \cong \bigoplus_{\mathbf{s} \in \mathbb{H}(L_G)} \mathfrak{A}^-(\mathbb{U}_n, \mathbf{s})$ is well-known to be

$$\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_n, \mathcal{V}_n] / (\mathcal{U}_i \mathcal{V}_i - \mathcal{U}_j \mathcal{V}_j, i, j \in \{1, \dots, n\}),$$

with the class of 1 having $(\text{gr}_w, \text{gr}_z)$ -bigrading $(0, 0)$. The plumbing diagram of the unlink $\mathbb{U}_n$ consists of n disjoint arrow vertices. After we assign the weight -1 to each arrow vertex, it is straightforward that the link lattice homology is also isomorphic to $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_n, \mathcal{V}_n] / (\mathcal{U}_i \mathcal{V}_i - \mathcal{U}_j \mathcal{V}_j, i, j \in \{1, \dots, n\})$ as an $\mathbb{F}[U]$ -module with the generator $[K, E]$ where $K = [2s_1 + 1, \dots, 2s_n + 1]$ with $s_i \in \mathbb{Z}$ and E consists of all arrow vertices equipped with the same gradings $(\text{gr}_w, \text{gr}_z) = (0, 0)$. That is,

$$d(\mathbb{U}_n, \mathbf{s}) = 2g(\mathbb{U}_n, \mathbf{s}) = \sum_{i=1}^n 2 \min \{0, s_i\}.$$

We now assume that the claim is true for some forest G of plumbing trees. We will prove the claim also holds for $L_G \# H$, where H is the positive Hopf link and the connected sum operation is between one knot component in L_G and one knot component of H . We recall that the complex of the positive Hopf link takes the following form:

$$\mathcal{CFL}(H) \cong \begin{array}{ccc} \mathbf{a} - \mathcal{U}_{n+1} & \rightarrow & \mathbf{b} \\ \downarrow \mathcal{V}_n & & \uparrow \mathcal{U}_n \\ \mathbf{c} & \leftarrow \mathcal{V}_{n+1} - & \mathbf{d} \end{array}. \quad (5.3)$$

The Alexander bigradings of $\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}$ are $(\frac{1}{2}, -\frac{1}{2})$, $(\frac{1}{2}, \frac{1}{2})$, $(-\frac{1}{2}, -\frac{1}{2})$ and $(-\frac{1}{2}, \frac{1}{2})$, respectively. The $(\text{gr}_w, \text{gr}_z)$ -bigradings are $(-1, -1)$, $(0, -2)$, $(-2, 0)$ and $(-1, -1)$, respectively.

We let $L_n \subseteq H$ be the component where the connected sum is taken, and $L_{n+1} \subseteq H$ the remaining component.

Topologically, the link L_G is obtained by taking an unlink, and iteratively taking the connected sum with Hopf links. We recall that the complex for an ℓ -component unlink is obtained by tensoring the $\ell - 1$ Koszul complexes

$$C_i = (\mathbf{x} \xrightarrow{\mathcal{U}_1 \mathcal{V}_1 + \mathcal{U}_i \mathcal{V}_i} \mathbf{y})$$

for $i = 2, \dots, \ell$. The $\ell = 2$ case of this computation is verified using a genus 0 Heegaard diagram for a 2-component unlink, and the general case is proved by iteratively using the connected sum formula [44, Section 7]. In (5.3), we observe that the Hopf link also has a similar 2-step filtration.

In particular, since a general L_G is obtained by tensoring an unlink with a collection of Hopf links, we may write $\mathcal{CFL}(L_G)$ as

$$\mathcal{F}^s \rightarrow \mathcal{F}^{s-1} \rightarrow \dots \rightarrow \mathcal{F}^0.$$

Call the index i for $\mathcal{F}^i$ the *Hopf grading*, where each $\mathcal{F}^i$ is a free $\mathcal{R}_n$ -module. By [41, Lemma 4.2] (cf. [43, Lemma 2.6]), L_G is an L-space link, so $\mathcal{HFL}(L_G)$ is supported in just one Hopf grading. We observe that the homology must be supported in $\mathcal{F}^0$, since the map $\mathcal{F}^1 \rightarrow \mathcal{F}^0$ is not surjective. Indeed, to see that the map is not surjective, define a map from $\mathcal{F}^0$ to $\mathbb{F}$ which sends all $\mathcal{U}_i$ and $\mathcal{V}_i$ to 1, and each basis element of $\mathcal{F}^0$ to 1. Then the composition $\mathcal{F}^1 \rightarrow \mathcal{F}^0 \rightarrow \mathbb{F}$ is zero, whereas the map $\mathcal{F}^0 \rightarrow \mathbb{F}$ is non-zero.

Applying the above argument to $\mathcal{CFL}(L_G \# H)$, we see that any homogeneously graded cycle in $\mathcal{CFL}(H) \otimes_{\mathbb{F}[\mathcal{U}_n, \mathcal{V}_n]} \mathcal{CFL}(L_G)$ which represents an $\mathbb{F}[U]$ -non-torsion element of homology may be written as a sum of an odd number of terms of the form $\alpha \cdot \mathbf{b} \otimes \mathbf{z}$ or $\beta \cdot \mathbf{c} \otimes \mathbf{z}$, where $\mathbf{z} \in \mathcal{CFL}(L_G)$ is an $\mathbb{F}[U]$ -non-torsion cycle, and $\alpha, \beta \in \mathbb{F}[\mathcal{U}_{n+1}, \mathcal{V}_{n+1}]$.

Consider $\mathbf{s} = (s_1, \dots, s_{n+1}) \in \mathbb{H}(L_G \# H)$, where $s_{n+1} \in \frac{1}{2} + \mathbb{Z}$ corresponds to the component $L_{n+1} \subseteq H$. We break the proof into two cases: $s_{n+1} > 0$ and $s_{n+1} < 0$.

We consider first the case where $s_{n+1} > 0$. Recall that $L_G \# H$ is an L-space link [41, Lemma 4.2]. In this case, $H_* \mathcal{A}^-(L_G \# H, \mathbf{s})$ is generated over $\mathbb{F}[U]$ by the elements $\mathcal{V}_{n+1}^{s_{n+1}-1/2} \cdot \mathbf{b} \otimes \mathbf{z}$ and $\mathcal{V}_{n+1}^{s_{n+1}+1/2} \cdot \mathbf{c} \otimes \mathbf{z}$, where $\mathbf{z}$ is $\mathbb{F}[U]$ -non-torsion. Write

$\mathbf{s}' = (s_1, \dots, s_n)$. The above argument yields

$$\begin{aligned} d(L_G \# H, \mathbf{s}) &= \max \{ \text{gr}_w(\mathbf{b}) + d(L_G, \mathbf{s}' - \tfrac{1}{2}e_n), \text{gr}_w(\mathbf{c}) + d(L_G, \mathbf{s}' + \tfrac{1}{2}e_n) \} \\ &= \max \{ d(L_G, \mathbf{s}' - \tfrac{1}{2}e_n), -2 + d(L_G, \mathbf{s}' + \tfrac{1}{2}e_n) \} \\ &= d(L_G, \mathbf{s}' - \tfrac{1}{2}e_n). \end{aligned} \quad (5.4)$$

Here the last equality follows from the facts that $d(L_G, \mathbf{s}' - \tfrac{1}{2}e_n) = -2H_{L_G}(\mathbf{s}' - \tfrac{1}{2}e_n)$ and $H_{L_G}(\mathbf{s}' - \tfrac{1}{2}e_n) \leq H_{L_G}(\mathbf{s}' + \tfrac{1}{2}e_n) + 1$ [2, Proposition 3.4]. Let $I' \subseteq V_G$ be any subset. We may view I' also as a subset of $V_G \cup \{v_{n+1}\}$. We compute easily that

$$\begin{aligned} 2f(\mathbf{s}, I') &= 2f(\mathbf{s}' - \tfrac{1}{2}e_n, I'), \\ 2f(\mathbf{s}, I' \cup \{v_{n+1}\}) &= \begin{cases} 2f(\mathbf{s}' - \tfrac{1}{2}e_n, I') + 2s_{n+1} + 1 & \text{if } v_n \in I', \\ 2f(\mathbf{s}' - \tfrac{1}{2}e_n, I') + 2s_{n+1} - 1 & \text{if } v_n \notin I', \end{cases} \end{aligned}$$

Since $s_{n+1} > 0$, we observe from these equations that

$$f(\mathbf{s}, I') = f(\mathbf{s}' - \tfrac{1}{2}e_n, I') \leq f(\mathbf{s}, I' \cup \{v_{n+1}\})$$

for all $I' \subseteq V_G$ and $\mathbf{s} \in \mathbb{H}(L_G)$. It follows easily that

$$g(L_G \# H, \mathbf{s}) = g(L_G, \mathbf{s}' - \tfrac{1}{2}e_n).$$

By induction $2g(L_G, \mathbf{s}' - \tfrac{1}{2}e_n) = d(L_G, \mathbf{s}' - \tfrac{1}{2}e_n)$, so from (5.4) we deduce that $2g(L_G \# H, \mathbf{s}) = d(L_G \# H, \mathbf{s})$ when $s_{n+1} > 0$.

We now consider the case $s_{n+1} < 0$. In this case, $H_*\mathcal{A}^-(L_G \# H, \mathbf{s})$ is generated by the elements of the form $\mathcal{U}_{n+1}^{-s_{n+1}+1/2} \cdot \mathbf{b} \otimes \mathbf{z}$ and $\mathcal{U}_{n+1}^{-s_{n+1}-1/2} \cdot \mathbf{c} \otimes \mathbf{z}$. A similar argument to the $s_{n+1} > 0$ case yields

$$d(L_G \# H, \mathbf{s}) = d(\mathbf{s}' + \tfrac{1}{2}e_n) + 2s_{n+1} - 1. \quad (5.5)$$

One computes directly that

$$\begin{aligned} 2f(\mathbf{s}, I') &= \begin{cases} 2f(\mathbf{s}' + \tfrac{1}{2}e_n, I') - 2 & \text{if } v_n \in I', \\ 2f(\mathbf{s}' + \tfrac{1}{2}e_n, I') & \text{if } v_n \notin I', \end{cases} \\ 2f(\mathbf{s}, I' \cup \{v_{n+1}\}) &= 2f(\mathbf{s}' + \tfrac{1}{2}e_n, I') + 2s_{n+1} - 1, \end{aligned} \quad (5.6)$$

for all $I' \subseteq V_G$. Since $s_{n+1} < 0$, we have

$$2f(\mathbf{s}, I' \cup \{v_{n+1}\}) = 2f(\mathbf{s}' + \tfrac{1}{2}e_n, I') + 2s_{n+1} - 1 \leq 2f(\mathbf{s}, I'),$$

where the last inequality comes from (5.6), so $2g(L_G \# H, \mathbf{s}) = 2g(L_G, \mathbf{s}' + \tfrac{1}{2}e_n) + 2s_{n+1} - 1$. Combining this with (5.5), we conclude that $2g(L_G \# H, \mathbf{s}) = d(L_G \# H, \mathbf{s})$, completing the proof of Lemma 5.5. $\blacksquare$

Proof of Proposition 5.3. Note that [41, Proposition 4.4] proves the identification on the level of chain complexes of $\mathbb{F}[[U]]$ -modules. It suffices to show that the decomposition

respects the refined actions of the ring $\mathcal{R}_\ell$. We recall the basics of their isomorphism. The lattice $\mathbb{H}(L_\Gamma)$ represents the set of Alexander gradings supported by the link Floer complex $\mathcal{CFL}(L_\Gamma)$. If $L \subseteq L_\Gamma$, and $\varepsilon \in \{0, 1\}^n$ is the indicator function for the components of L , then we may write $\mathfrak{X}_\varepsilon^-(L_\Gamma, \mathbf{s})$ for the subcomplex of $S_\varepsilon^{-1} \cdot \mathcal{CFL}(L_\Gamma)$ in Alexander grading $\mathbf{s}$. According to [41, Lemma 4.2], there is an isomorphism

$$H_*(\mathfrak{X}_\varepsilon^-(L_\Gamma, \mathbf{s})) \cong \mathbb{F}[[U]].$$

Additionally, there is an isomorphism $\mathbb{H}(L_\Gamma) \rightarrow \text{Spin}^c(X_\Gamma)$ (stated in (4.10)). This gives an isomorphism between $\mathbb{Z}$ and $\mathbb{CFL}(\Gamma, V_\uparrow)$ as $\mathbb{F}[[U]]$ -modules. The Maslov grading of the generator of $H_*(\mathfrak{X}_\varepsilon^-(L_\Gamma, \mathbf{s}))$ is $d(L_\Gamma, \mathbf{s})$ and the maps $D_{\varepsilon, \varepsilon'}$ for $|\varepsilon' - \varepsilon| = 1$ are determined by the Maslov grading of the generators of the domain and target. Applying Lemma 5.5 and the argument in [41, Proposition 4.4], the differentials also coincide. It remains to show that the isomorphism respects the $\mathcal{R}_\ell$ -module structure.

To disambiguate the actions, let us write $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]]$ for the action we have described on $\mathbb{CFL}(\Gamma, V_\uparrow)$, and write $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]]$ for the action induced by the identification $\mathbb{CFL}(\Gamma, V_\uparrow) \cong \mathbb{Z}$. As a first step, note that by definition $\mathcal{U}_i$ changes the Alexander grading $\mathbf{s} \in \mathbb{H}(L_\Gamma)$ by $-e_i \in \mathbb{Z}^n$ (where e_i is the unit vector with i -th component 1, and the other components 0). It follows by Lemma 5.5 that $\mathcal{U}_i$ has the same gr_w -grading as $\mathcal{U}_i$, and similarly $\mathcal{V}_i$ has the same gr_w -grading as $\mathcal{V}_i$. Clearly, if $[K, E]$ is a generator, then $\mathcal{U}_i \cdot [K, E] = U^\varepsilon [K - 2\mu_i^*, E]$ and $\mathcal{U}_i \cdot [K, E] = U^{\varepsilon'} [K - 2\mu_i^*, E]$ for some $\varepsilon, \varepsilon' \in \{0, 1\}$. Since $\mathcal{U}_i$ and $\mathcal{U}_i$ have the same gr_w -grading as endomorphisms, we must have $\varepsilon = \varepsilon'$, so that $\mathcal{U}_i$ and $\mathcal{U}_i$ have the same action. The same argument implies $\mathcal{V}_i = \mathcal{V}_i$. ■

5.3. Proof of Theorem 5.1

In this subsection we give the proof of Theorem 5.1, though we delay the discussion of absolute gradings until Section 5.4. The main steps of the proof follow [59, Section 5.2]. We provide a summary and highlight the necessary changes for our present setting.

By Theorem 5.2, we have a homotopy equivalence of chain complexes over $\mathcal{R}_\ell$:

$$\mathcal{CFL}(Y_G, L_\uparrow) \simeq \mathcal{C}_\Lambda(L_G, L_\uparrow).$$

In particular, the above chain homotopy equivalence may be viewed as a homotopy equivalence of A_∞ -modules over $\mathcal{R}_\ell$, where each module has $m_j = 0$ for $j > 2$. Above, we also constructed a hypercube $\mathcal{Z} = (Z_\varepsilon, \delta_{\varepsilon, \varepsilon'})$ of $\mathcal{R}_\ell$ -modules by taking the homology of $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ at each cube point, and using only the length 1 maps from $\mathcal{C}_\Lambda(L_G, L_\uparrow)$. By Proposition 5.3, we have a chain isomorphism

$$\mathcal{Z} \simeq \mathbb{CFL}(\Gamma, V_\uparrow).$$

Hence, it suffices to construct an A_∞ -homotopy equivalence

$$\mathcal{Z} \simeq \mathcal{C}_\Lambda(L_G, L_\uparrow). \quad (5.7)$$

This homotopy equivalence follows by the same logic as in [59, Section 6.2], which we summarize for the benefit of the reader.

We will write $\mathcal{C} = (\mathcal{C}_\varepsilon, \Phi_{\varepsilon, \varepsilon'})$ for $\mathcal{C}_\Lambda(L_G, L_\uparrow)$. The underlying complex $\mathcal{C}_\varepsilon$ at each vertex of $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ is obtained from $\mathcal{CF}\mathcal{L}(L_\Gamma)$ by localizing at the variables $\mathcal{V}_i$ such that $\varepsilon(i) = 1$, and then taking an appropriate completion. The complex $\mathcal{CF}\mathcal{L}(L_\Gamma)$ is a tensor product of Hopf link complexes. The Hopf link has a 2-step filtration (see (5.3)), and hence $\mathcal{CF}\mathcal{L}(L_\Gamma)$ has a description as

$$\mathcal{CF}\mathcal{L}(S^3, L_G \cup L_\uparrow) \cong (\mathcal{F}^m \rightarrow \mathcal{F}^{m-1} \rightarrow \dots \rightarrow \mathcal{F}^0), \quad (5.8)$$

where each $\mathcal{F}^i$ is a free $\mathcal{R}_n$ -module, $n = |V_\Gamma|$ and m is the number of edges in Γ (i.e. Hopf link components). Each $\mathcal{C}_\varepsilon$ has a similar filtration, which we denote by $\mathcal{F}_\varepsilon^i$. We call the superscript i in $\mathcal{F}^i$ the *Hopf grading*.

Since L_Γ is an L-space link, each Z_ε is a direct product of copies of $\mathbb{F}[[U]]$. Following [59, Proposition 6.2], there is a natural way to construct a homotopy equivalence between $\mathcal{C}_\varepsilon$ and Z_ε for each ε . This is because the homology of $\mathcal{C}_\varepsilon$ is supported in $\mathcal{F}_\varepsilon^0$ so the projection map of $\mathcal{F}_\varepsilon^0$ onto Z_ε gives a quasi-isomorphism. Since in each Alexander grading (i.e. each $\mathbf{s} \in \mathbb{H}(L_\Gamma)$) the homology of Z_ε is $\mathbb{F}[[U]]$ (in particular, a projective $\mathbb{F}[[U]]$ -module), it is straightforward to construct a splitting over $\mathbb{F}[[U]]$ of the sequence in (5.8) in each Alexander grading. This gives us maps

$$\pi_\varepsilon: \mathcal{C}_\varepsilon \rightarrow Z_\varepsilon, \quad i_\varepsilon: Z_\varepsilon \rightarrow \mathcal{C}_\varepsilon \quad \text{and} \quad h_\varepsilon: \mathcal{C}_\varepsilon \rightarrow \mathcal{C}_\varepsilon,$$

which satisfy

$$\pi_\varepsilon \circ i_\varepsilon = \text{id}, \quad i_\varepsilon \circ \pi_\varepsilon = \text{id} + [\partial, h_\varepsilon], \quad h_\varepsilon \circ h_\varepsilon = 0, \quad \pi_\varepsilon \circ h_\varepsilon = 0, \quad h_\varepsilon \circ i_\varepsilon = 0.$$

Remark 5.6. Note that these maps are usually only $\mathbb{F}[[U]]$ -equivariant, but not necessarily $\mathcal{R}_\ell$ -equivariant. The only exception is π_ε , because it is the canonical projection of $\mathcal{F}_\varepsilon^0$ to $\mathcal{F}_\varepsilon^0/\text{im } \mathcal{F}^1 \cong H_*(Z_\varepsilon)$.

The homological perturbation lemma for hypercubes (see [17, Lemma 2.10]) induces hypercube structure maps $d_{\varepsilon, \varepsilon'}$ on $\bigoplus_{\varepsilon \in \{0,1\}^n} Z_\varepsilon$, which is homotopy equivalent to the hypercube $\mathcal{C}_\Lambda(L_G, L_\uparrow)$. Let us write $\mathcal{W}$ for the hypercube $(Z_\varepsilon, d_{\varepsilon, \varepsilon'})$. The structure maps $d_{\varepsilon, \varepsilon'}$ are given by the concrete formula

$$d_{\varepsilon, \varepsilon'} := \sum_{\varepsilon = \varepsilon_1 < \dots < \varepsilon_n = \varepsilon'} \pi_{\varepsilon_n} \circ \Phi_{\varepsilon_{n-1}, \varepsilon_n} \circ h_{\varepsilon_{n-1}} \circ \dots \circ h_{\varepsilon_2} \circ \Phi_{\varepsilon_1, \varepsilon_2} \circ i_{\varepsilon_1}. \quad (5.9)$$

There are also homotopy equivalences $\Pi: \mathcal{C} \rightarrow \mathcal{W}$, $I: \mathcal{W} \rightarrow \mathcal{C}$ and $H: \mathcal{C} \rightarrow \mathcal{C}$, given by similar formulas, which satisfy

$$\Pi \circ I = \text{id}, \quad I \circ \Pi = \text{id} + \partial(H), \quad H \circ H = 0, \quad \Pi \circ H = 0 \quad \text{and} \quad H \circ I = 0.$$

We will also need to understand the map Π , which is given by

$$\Pi_{\varepsilon, \varepsilon'} = \sum_{\varepsilon = \varepsilon_1 < \dots < \varepsilon_n = \varepsilon'} \pi_{\varepsilon_n} \circ \Phi_{\varepsilon_{n-1}, \varepsilon_n} \circ h_{\varepsilon_{n-1}} \circ \dots \circ \Phi_{\varepsilon_1, \varepsilon_2} \circ h_{\varepsilon_1}. \quad (5.10)$$

A natural strategy is to show that $\mathcal{W} = \mathcal{Z}$. Note that the underlying groups are identical, so it suffices to understand the structure maps. We observe that $\delta_{\varepsilon, \varepsilon'} = d_{\varepsilon, \varepsilon'}$ whenever $|\varepsilon - \varepsilon'|_{L^1} = 1$. In general, it is not the case that $d_{\varepsilon, \varepsilon'} = \delta_{\varepsilon, \varepsilon'}$ when $|\varepsilon - \varepsilon'|_{L^1} > 1$. Instead the main argument of [59] is to show that the analog of $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ is homotopy equivalent to a hypercube which has the same underlying internal chain complexes as $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ and for which the induced hypercube structure on the analog of $\mathcal{W}$ coincides with that of $\mathcal{Z}$. The argument in our present setting is essentially identical to the one in [59, Section 6.2]. In fact, we may take the modified hypercube structure on $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ to be the restriction of the one constructed in [59], viewing $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ as a subcube of the full link surgery hypercube for L_Γ . We write $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ for the resulting hypercube. We will also write

$$\mathcal{C}' = (\mathcal{C}_\varepsilon, \Phi'_{\varepsilon, \varepsilon'})$$

for $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$, and $\mathcal{W}'$ for the hypercube constructed via homological perturbation. There are maps

$$\Pi': \mathcal{C}' \rightarrow \mathcal{W}', \quad I': \mathcal{W}' \rightarrow \mathcal{C}' \quad \text{and} \quad H': \mathcal{C}' \rightarrow \mathcal{C}'$$

constructed similarly to the maps Π , I and H , using formulas as in (5.10).

Concretely, $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ is constructed by tensoring the link surgery complex of Hopf links using the tensor product formula from [59, (3.2)]. The complex $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ is constructed by tensoring the link surgery complexes of Hopf links together using an algebraically simplified model of the tensor product, where several terms have been deleted (this simplified model appears in [59, Theorem 3.4]). The key property of the hypercube maps appearing in $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ is that the maps $\Phi'_{\varepsilon, \varepsilon'}$ are non-increasing in the Hopf grading from (5.8) only when $|\varepsilon - \varepsilon'|_{L^1} \leq 1$. Furthermore, when $|\varepsilon - \varepsilon'|_{L^1} = 1$, the map $\Phi'_{\varepsilon, \varepsilon'}$ preserves the Hopf grading. Since h_ε strictly increases the Hopf grading and π_ε is non-vanishing only on the lowest Hopf grading, the composition in (5.9) will only be non-trivial when $|\varepsilon - \varepsilon'|_{L^1} = 1$. Hence

$$\mathcal{W}' = \mathcal{Z}.$$

In particular, composing these homotopy equivalences, we obtain a homotopy equivalence of chain complexes

$$\mathcal{CFL}(Y_G, L_\uparrow) \simeq \mathcal{Z}. \quad (5.11)$$

It remains to show that the homotopy equivalence in (5.11) may be extended to a homotopy equivalence of A_∞ -modules over $\mathcal{R}_\ell$. There are two subclaims:

- (R-1) The homotopy equivalence between $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ and $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ may be taken to be $\mathcal{R}_\ell$ -equivariant.
- (R-2) The homotopy equivalence $\mathcal{C}'_\Lambda(L_G, L_\uparrow) \simeq \mathbb{CFL}(\Gamma, V_\uparrow)$ (from the homological perturbation lemma for hypercubes) extends to a homotopy equivalence of A_∞ -modules over $\mathcal{R}_\ell$.

We address (R-1) first. In [59, Corollary 4.8], it is shown that the simplified connected sum formula yielding $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ is valid as long as in forming L_Γ by an iterated

connected sum, we never take the connected sum of two knot components which are both homologically essential after we surger on the other components of L_G . In the case that $b_1(Y_G) = 0$, we always avoid this configuration (cf. [59, Lemma 6.5]). The homotopy equivalence between $\mathcal{C}_\Lambda(L_G, L_\uparrow)$ and $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ is concrete and obtained by relating the connected sum formula in [59, (3.2)] and the simplified connected sum formula in [59, Theorem 3.4]. As described in [59, Corollary 4.8], relating these two models amounts to constructing a null-homotopy of an algebraically defined homology action on the link surgery formula. It is observed in [59, Remark 4.3] that this null-homotopy may be taken to be $\mathcal{R}_\ell$ -equivariant.

We now address (R-2). Proposition 5.3 shows that $\mathcal{Z}$ is chain isomorphic as an $\mathcal{R}_\ell$ -module to $\mathbb{CFL}(\Gamma, V_\uparrow)$. It is sufficient to show that the homotopy equivalence $\mathcal{C}' \simeq \mathcal{Z}$ of chain complexes is in fact a homotopy equivalence of A_∞ -modules over $\mathcal{R}_\ell$. To see this, it is sufficient to show that the map $\Pi': \mathcal{C}' \rightarrow \mathcal{Z}$, defined as in (5.10), commutes with the $\mathcal{R}_\ell$ -action. This is sufficient because in the category of A_∞ -modules, quasi-isomorphisms are always invertible as A_∞ -morphisms. To establish the $\mathcal{R}_\ell$ -equivariance, we observe that the projection maps π_ε are themselves equivariant since they are merely quotient maps; compare Remark 5.6. Next, we examine the expression for $\Pi'_{\varepsilon, \varepsilon'}$ as in (5.10). By considering the Hopf grading similarly to what we did with $d_{\varepsilon, \varepsilon'}$ above, we observe that the maps $\Pi'_{\varepsilon, \varepsilon'}$ are non-trivial only when $\varepsilon = \varepsilon'$. In this case, the only non-vanishing contribution is from π_ε , so $\mathcal{R}_\ell$ -equivariance is established, and the proof is complete. ■

Remark 5.7. By using the homological perturbation lemma, stated in Lemma 2.3, a concrete homotopy equivalence of A_∞ -modules between $\mathcal{C}'_\Lambda(L_G, L_\uparrow)$ and $\mathcal{Z}$ may be constructed. Indeed, the hypercube maps Π' , I' and H' , constructed via the homological perturbation lemma for hypercubes, also satisfy the assumptions of the homological perturbation lemma for A_∞ -modules. These maps then induce an A_∞ -module structure on $\mathcal{Z}$ over the ring $\mathcal{R}_\ell$, which is homotopy equivalent to $\mathcal{CFL}(Y_G, L_\uparrow)$. By considering the Hopf grading, similarly to the above, we obtain only non-trivial m_1 and m_2 on $\mathcal{Z}$. We also observe that the morphisms Π' , H' and I' extend to A_∞ -morphisms Π'_j , H'_j and I'_j . Hopf grading considerations show that $\Pi'_j = 0$ unless $j = 1$, in which case the only contribution is from π_ε . We observe that I'_j may be non-trivial for $1 \leq j \leq |L_\Gamma|$.

5.4. Absolute gradings

In this section, we prove the subclaim of Theorem 5.1 concerning the absolute Maslov and Alexander gradings; compare [41, Proposition 4.8].

We begin by stating formulas for the absolute gradings on the link surgery complex, and its subcube refinement for sublinks. Although likely known to experts, these formulas have not appeared in the literature except in special cases. For example, in the case of knots, the result is due to Ozsváth and Szabó [44, Section 4]. A detailed proof of the absolute grading formula can be found in [58, Section 10].

If $L \subseteq S^3$ is a link with framing Λ , let $W_\Lambda(L)$ denote the standard 2-handle cobordism from S^3 to $S^3_\Lambda(L)$. If $\varepsilon \in \{0, 1\}^n$, where $n = |L|$, and $\mathbf{s} \in \mathbb{H}(L)$, write $\mathcal{C}_\varepsilon(\mathbf{s}) \subseteq \mathcal{C}_\Lambda(L)$ for

the subspace of $\mathcal{C}_\varepsilon$ which lies in internal Alexander grading $\mathbf{s}$. Finally, if $\mathbf{s} \in \mathbb{H}(L)$, write $\mathfrak{z}_\mathbf{s} \in \text{Spin}^c(W_\Lambda(L))$ for the Spin^c structure which satisfies

$$\frac{\langle c_1(\mathfrak{z}_\mathbf{s}), \Sigma_i \rangle - \Sigma \cdot \Sigma_i}{2} = -s_i \quad (5.12)$$

for all $i \in \{1, \dots, n\}$. In the above, Σ_i denotes the core of the 2-handle attached along the component $K_i \subseteq L$, and Σ denotes the sum of all Σ_i .

Lemma 5.8 ([58, Theorem 10.2]). *Suppose that $L \subseteq S^3$ is a link with framing Λ and that $b_1(S_\Lambda^3(L)) = 0$. The homotopy equivalence $\mathcal{CF}^-(S_\Lambda^3(L)) \simeq \mathcal{C}_\Lambda(L)$ is absolutely graded if we equip $\mathcal{C}_\varepsilon(\mathbf{s}) \subseteq \mathcal{C}_\Lambda(L)$ with the Maslov grading*

$$\widetilde{\text{gr}} := \text{gr}_w + \frac{c_1(\mathfrak{z}_\mathbf{s})^2 - 2\chi(W_\Lambda(L)) - 3\sigma(W_\Lambda(L))}{4} + |L| - |\varepsilon|,$$

where gr_w is the Maslov grading from $\mathcal{CFL}(L)$.

In [50], Ozsváth and Szabó prove this formula in the context of the knot surgery formula. Their main tool is computing the grading change of a surgery cobordism map, which they denote by f_3^+ . A similar strategy is likely possible in the context of the link surgery formula. Nonetheless, algebraic truncations make writing a simple proof challenging.

There is also a relative version of the statement. Suppose that we have a partitioned link $J \sqcup L \subseteq S^3$ and that J is equipped with an integral framing Λ . If K_i is a link component of $J \sqcup L$, there is a class $\Sigma_i \in H_2(W_\Lambda(J), \partial W_\Lambda(J))$. If K_i is in J , then Σ_i is the core of the corresponding 2-handle. If K_i is in L , then Σ_i is an annulus, with boundary on the images of K_i in S^3 and $S_\Lambda^3(J)$. We write Σ for the sum of the classes for all components of $J \sqcup L$. If the component $K_i \subseteq L$ becomes rationally null-homologous in $S_\Lambda^3(J)$, we write $\hat{\Sigma}_i$ for the class in $H_2(W_\Lambda(J); \mathbb{Q})$ obtained by capping with a rational Seifert surface. Finally, if $\mathbf{s} \in \mathbb{H}(L)$, we write $\mathfrak{z}_\mathbf{s}^J \in \text{Spin}^c(W_\Lambda(J))$ for the Spin^c structure which satisfies (5.12) for all link components K_i in J .

Lemma 5.9 ([58, Theorem 10.8]). *Suppose that $J \cup L \subseteq S^3$ is a partitioned link and Λ is an integral framing on J such that $b_1(S_\Lambda^3(J)) = 0$. Then the isomorphism $\mathcal{CFL}(S_\Lambda^3(J), L) \simeq \mathcal{C}_\Lambda(J, L)$ from Theorem 5.2 is absolutely gr_w -graded if we equip $\mathcal{C}_\varepsilon(\mathbf{s}) \subseteq \mathcal{C}_\Lambda(J, L)$ with the Maslov grading*

$$\widetilde{\text{gr}}_w = \text{gr}_w + \frac{c_1(\mathfrak{z}_\mathbf{s}^J)^2 - 2\chi(W_\Lambda(J)) - 3\sigma(W_\Lambda(J))}{4} + |J| - |\varepsilon|,$$

where gr_w is the internal gr_w -grading on $\mathcal{CFL}(J \cup L)$. The isomorphism is absolutely graded with respect to the Alexander grading $A = (A_1, \dots, A_{|L|})$ if we define A_i on $\mathcal{C}_\varepsilon(\mathbf{s})$ via the formula

$$A_i = s_i + \frac{\langle c_1(\mathfrak{z}_\mathbf{s}^J), \hat{\Sigma}_i \rangle - \Sigma \cdot \hat{\Sigma}_i}{2}.$$

Proposition 5.10. *If $b_1(Y_G) = 0$, the isomorphism from Theorem 5.1 respects the absolute Maslov and Alexander gradings.*

Proof. As a first step, we consider the absolute case when there are no arrow vertices. We recall that we already constructed an isomorphism $\Phi_G: \text{Char}(X_G) \rightarrow \mathbb{H}(L)$ in (4.10). In the present case, it is straightforward to verify from the definitions that $c_1(\mathfrak{z}_s) = -\Phi_G^{-1}(s)$. If $s \in \mathbb{H}(L_G)$ and $s = \Phi_G(K)$, then this is equivalent to

$$c_1(\mathfrak{z}_s) = -K. \quad (5.13)$$

Next, we recall that in Lemma 5.5, we identified the quantity $g(K, E)$ with the gr_w -grading of the generator of the tower in the s -graded subspace of the homology of $\mathcal{CFL}(L_G)$, localized at the $\mathcal{V}_i$ variables for vertices in E . Noting that $(-K)^2 = K^2$, we obtain (4.6).

We now consider the case where there are arrow vertices. Suppose that Γ is a tree with $V_\Gamma = V_G \cup V_\uparrow$. Lemma 5.9 computes the Maslov grading shift. If $K \in \text{Char}(X_\Gamma)$ and $s = \Phi_\Gamma(K)$, then we have, similarly to (5.13),

$$c_1(\mathfrak{z}_s^{L_G}) = -K|_{X_G}. \quad (5.14)$$

In particular, the statement from Lemma 5.5 implies that the Maslov grading on $\mathcal{CFL}(S_\Lambda^3(L_G), L_\uparrow)$ coincides with the one defined in (4.7).

We now consider the Alexander grading. We note that given $s \in \mathbb{H}(L_\Gamma)$, there are two Spin^c structures of interest: $\mathfrak{z}_s^{L_G} \in \text{Spin}^c(X_G)$ and $\mathfrak{z}_s^{L_\Gamma} \in \text{Spin}^c(X_\Gamma)$. It is straightforward to verify that $\mathfrak{z}_s^{L_G} = \mathfrak{z}_s^{L_\Gamma}|_{X_G}$.

If $s = (s_1, \dots, s_{|L_\Gamma|}) \in \mathbb{H}(L_\Gamma)$ and $K_i \in L_\uparrow$, we view $\mathcal{C}_\Lambda(L_G, L_\uparrow, s)$ as having internal Alexander grading A_i equal to s_i . Lemma 5.9 implies that the isomorphism $\mathcal{CFL}(S_\Lambda^3(L_G), L_\uparrow) \simeq \mathcal{C}_\Lambda(L_G, L_\uparrow)$ is Alexander graded if we shift the internal Alexander grading s_i (for a component $K_i \in L_\uparrow$) of $\mathcal{C}_\Lambda(L_G, L_\uparrow, s)$ by $(\langle c_1(\mathfrak{z}_s^{L_G}), \hat{\Sigma}_i \rangle - \Sigma \cdot \hat{\Sigma}_i)/2$.

By the definition of Φ_Γ , if $\Phi_\Gamma(K) = s$, then

$$s_i = \frac{\langle K, \Sigma_i \rangle + \Sigma \cdot \Sigma_i}{2},$$

where the pairings are taken in X_Γ . In particular, the generator $[K, E]$ in the lattice complex will be given i -th Alexander grading

$$s_i + \frac{\langle c_1(\mathfrak{z}_s^{L_G}), \hat{\Sigma}_i \rangle - \Sigma \cdot \hat{\Sigma}_i}{2}.$$

By (5.14), we may write the above as

$$\frac{K(v_i - \hat{v}_i) + \sum_{v \in V_\Gamma} v \cdot (v_i - \hat{v}_i)}{2}.$$

We note that for $v \in V_G$, the pairing $v \cdot (v_i - \hat{v}_i)$ will vanish, and hence we can replace the sum in the above equation with a sum over only $v \in V_\uparrow$. This recovers the formula for the Alexander grading stated in Section 4.6, so the proof is complete. ■

6. Plumbed L-space links

In this section, we compute the link Floer complexes of plumbed L-space links. In Section 6.1, we prove that their complexes are formal (i.e. $\mathcal{CFL}(Y_G, L_\uparrow)$ is homotopy equivalent as an A_∞ -module to $(\mathcal{HFL}(Y_G, L_\uparrow), m_j)$, where $m_j = 0$ unless $j = 2$). Consequently, the chain complexes are also homotopy equivalent to free resolutions of their homology. In Section 6.2 we recall work of Gorsky and Némethi [11] which allows one to compute the module $\mathcal{HFL}(Y_G, L_\uparrow)$ when $Y_G = S^3$ and $L_\uparrow$ is an L-space link. We extend their description to the case where Y_G is a rational homology L-space. Finally, in Section 6.3, we prove that our model of link lattice homology recovers the version of Gorsky and Némethi [11] in the case of plumbed L-space links.

6.1. Plumbed L-space links and free resolutions

Suppose Γ is an arrow-decorated plumbing tree. Our goal is to show that the chain complex $\mathcal{CFL}(Y_G, L_\uparrow)$ is a free resolution of its homology, in particular, the chain complex is determined by the homology up to chain homotopy equivalence.

In this section, we consider plumbings where $L_\uparrow$ is an L-space link. We observe that this implies that Y_G is a rational homology 3-sphere, and furthermore it is itself an L-space. In particular, by Proposition 3.18, the results in this subsection hold for links of embedded analytic singularities, as long as the underlying surface singularity is rational.

Theorem 6.1. *Suppose that Γ is an arrow-decorated plumbing tree with $V_\Gamma = V_G \cup V_\uparrow$. Let $L_\uparrow \subseteq Y_G$ be the associated link and assume that Y_G is a rational homology 3-sphere. If $L_\uparrow$ is an L-space link, then $\mathcal{CFL}(Y_G, L_\uparrow)$ is a free resolution over $\mathcal{R}_\ell$ of $\mathcal{HFL}(Y_G, L_\uparrow)$.*

Proof. To simplify the notation, we assume that Y_G is an integer homology 3-sphere; for rational homology 3-spheres, one may apply the same argument to each Spin^c structure.

Next, we observe that it is sufficient to show that $\mathcal{CFL}(Y_G, L_\uparrow)$ is homotopy equivalent to a free resolution of $\mathcal{HFL}(Y_G, L_\uparrow)$. This may be seen as follows: The $\mathcal{R}_\ell$ -module $\mathcal{HFL}(Y_G, L_\uparrow)$ is finitely generated and hence admits a finitely generated free resolution over $\mathcal{R}_\ell$ by Hilbert's syzygy theorem [18] (see e.g. [51, Theorem 15.2] for a modern exposition). Furthermore, it is straightforward to see that if C and C' are two free, finitely generated chain complexes over $\mathcal{R}_\ell$ which are both $(\text{gr}_w, \text{gr}_z)$ -graded, then C and C' are homotopy equivalent over $\mathcal{R}_\ell$ if and only if $C \otimes_{\mathcal{R}_\ell} \mathcal{R}_\ell$ and $C' \otimes_{\mathcal{R}_\ell} \mathcal{R}_\ell$ are homotopy equivalent. Moreover, the completion of a free resolution of $\mathcal{HFL}(Y_G, L_\uparrow)$ will be a free resolution of $\mathcal{HFL}(Y_G, L_\uparrow)$, by similar reasoning. Hence, $\mathcal{CFL}(Y_G, L_\uparrow)$ will be homotopy equivalent to a free resolution of $\mathcal{HFL}(Y_G, L_\uparrow)$ if and only if $\mathcal{CFL}(Y_G, L_\uparrow)$ is homotopy equivalent to a free resolution of $\mathcal{HFL}(Y_G, L_\uparrow)$.

Since $L_\uparrow \subseteq Y_G$ is an L-space link, $\mathcal{HFL}(Y_G, L_\uparrow)$ is a torsion-free $\mathbb{F}[U]$ -module. By Theorem 5.1, $\mathcal{CFL}(Y_G, L_\uparrow)$ is homotopy equivalent to $\mathbb{CFL}(\Gamma, V_\uparrow)$ as an A_∞ -module over $\mathcal{R}_\ell$.

The link lattice complex has a cube grading:

$$\mathbb{CFL}(\Gamma, V_\uparrow) = (Z_n \xrightarrow{f_{n,n-1}} Z_{n-1} \rightarrow \cdots \rightarrow Z_1 \xrightarrow{f_{1,0}} Z_0),$$

where Z_q is spanned by $U^p[K, E]$ where $|E| = q$, for all $p \geq 0$. Furthermore, each Z_i is itself an $\mathcal{R}_\ell$ -module (in particular, the action of $\mathcal{R}_\ell$ preserves the cube grading). We make the following subclaim:

Claim. *The homology of $\mathbb{CFL}(\Gamma, V_\uparrow)$ is supported in a single cube grading Z_i .*

Note that the main theorem statement follows from the above claim, and from Lemma 2.7.

To prove the above claim, we observe that if there are two Z_i which support the homology, then $\mathcal{HFL}(Y_G, L_\uparrow)$ will split as a direct sum of $\mathcal{R}_\ell$ -modules; we will show that this is impossible.

We argue by considering the localization at the multiplicatively closed subset $S \subseteq \mathcal{R}_\ell$ spanned by the monomials. Since $L_\uparrow \subseteq Y_G$ is an L-space link, it follows that $s \cdot x \neq 0$ for all $s \in S$ and non-zero $x \in \mathcal{HFL}(Y_G, L_\uparrow)$. Therefore, the localization map is injective, so it suffices to show that $S^{-1}\mathcal{HFL}(Y_G, L_\uparrow)$ does not split as the direct sum of two $\mathcal{R}_\ell$ -modules.

We recall that localization is an exact functor, so it suffices to consider the localization on the chain level. We pick an arbitrary $(s_1, \dots, s_n) \in \mathbb{H}(Y, L)$. We observe that after localizing, we may perform a change of basis and replace each basis element $\mathbf{x}$ with a basis element

$$\mathcal{V}_1^{-A_1(\mathbf{x})+s_1} \dots \mathcal{V}_\ell^{-A_\ell(\mathbf{x})+s_\ell} \mathbf{x}$$

(we take $\mathbf{s}$ such that the powers of $\mathcal{V}_i$ are integral). With this choice of basis, all generators of $\mathcal{CFL}(Y_G, L_\uparrow)$ are now concentrated in Alexander grading $\mathbf{s}$. Since the differential preserves the Alexander grading, each component of the differential is weighted by powers of $U_i = \mathcal{U}_i \mathcal{V}_i$ for $i \in \{1, \dots, \ell\}$. Setting $\mathcal{V}_i$ equal to 1 recovers $\mathcal{CF}^\infty(Y_G, L_\uparrow)$, so we conclude that

$$\mathcal{CFL}^\infty(Y_G, L_\uparrow) \cong \mathcal{CF}^\infty(Y_G, w_1, \dots, w_\ell) \otimes_{\mathbb{F}[U_1, \dots, U_\ell]} \mathcal{R}_\ell,$$

where U_i acts on $\mathcal{R}_\ell$ by $\mathcal{U}_i \mathcal{V}_i$. Here, $\mathcal{CF}^\infty(Y_G, w_1, \dots, w_\ell)$ denotes the ℓ -pointed Floer complex for Y_G . It follows from [45, Theorem 10.1] that the singly pointed Floer complex $\mathcal{CF}^\infty(Y_G, w_1)$ is homotopy equivalent to $\mathbb{F}[[U, U^{-1}]]$, and by [48, Proposition 6.5], adding the basepoint w_i has the effect of tensoring with the Koszul complex $\mathbf{x} \xrightarrow{U_1+U_i} \mathbf{y}$. In particular, it follows that $\mathcal{HFL}^\infty(Y_G, L_\uparrow)$ is isomorphic to

$$S^{-1}\mathcal{R}_\ell / (\mathcal{U}_i \mathcal{V}_i - \mathcal{U}_j \mathcal{V}_j : i, j \in \{1, \dots, \ell\}).$$

This does not decompose as a direct sum of $\mathcal{R}_\ell$ -modules, since it has a single generator (the image of $1 \in \mathcal{R}_\ell$) over the ring $S^{-1}\mathcal{R}_\ell$.

This completes the proof of the Claim, and therefore of the main statement as well. ■

6.2. Computing the Floer chain complex from Alexander polynomials

In this subsection, we consider the H -function for oriented ℓ -component links L in rational homology spheres Y . We begin by defining a lattice $\mathbb{H}(Y, L)$, which is an affine

space over $H_1(Y \setminus L; \mathbb{Z})$. The set $\mathbb{H}(Y, L)$ is a subspace of $\mathbb{Q}^\ell \times \text{Spin}^c(Y)$. The simplest definition of $\mathbb{H}(Y, L)$ is that it is the set of $(\mathbf{s}, \mathbf{t}) \in \mathbb{Q}^\ell \times \text{Spin}^c(Y)$ such that $\widehat{HFK}(Y, L, \mathbf{t})$ is non-trivial in some Alexander grading $\mathbf{s}' \in \mathbb{Q}^\ell$ satisfying $\mathbf{s} - \mathbf{s}' \in \mathbb{Z}^\ell$. Since $\widehat{HFK}(Y, L, \mathbf{t}) \neq 0$ for each $\mathbf{t}$ when Y is a rational homology 3-sphere, this construction gives a well-defined set $\mathbb{H}(Y, L)$. The action of an element $\gamma \in H_1(Y \setminus L)$ is given by

$$\gamma \cdot (s_1, \dots, s_\ell, \mathbf{t}) = (s_1 - \text{lk}(\gamma, K_1), \dots, s_\ell - \text{lk}(\gamma, K_\ell), \mathbf{t} + PD[i_*\gamma]),$$

where $\text{lk}(\gamma, K_i) \in \mathbb{Q}$ is the rational linking number, and $i: Y \setminus L \rightarrow Y$ is inclusion.

A more topological description may be obtained by representing $Y \setminus L$ as surgery on a link L' in the complement of an ℓ -component unlink in S^3 . Such a representation induces a link cobordism (W, Σ) from the complement of an ℓ -component unlink to $Y \setminus L$ such that W is a 2-handlebody, and Σ consists of ℓ annuli, each of which cobounds an unknot component in S^3 and a knot component of L . If $\mathbf{t} \in \text{Spin}^c(Y)$, the fiber over $\mathbf{t}$ of the map $\mathbb{H}(Y, L) \rightarrow \text{Spin}^c(Y)$ consists of $(\mathbf{s}, \mathbf{t})$ where

$$\mathbf{s} \in \left(\frac{\langle c_1(\mathbf{t}'), \hat{\Sigma}_1 \rangle - [\hat{\Sigma}_1] \cdot [\Sigma]}{2}, \dots, \frac{\langle c_1(\mathbf{t}'), \hat{\Sigma}_\ell \rangle - [\hat{\Sigma}_\ell] \cdot [\Sigma]}{2} \right) + \mathbb{Z}^\ell, \quad (6.1)$$

and $\mathbf{t}' \in \text{Spin}^c(W)$ is any lift of $\mathbf{t} \in \text{Spin}^c(Y)$. Here, we view Σ as the union of $\Sigma_1, \dots, \Sigma_\ell$. We write $[\Sigma]$ and $[\Sigma_i]$ for the induced classes in $H_2(W, \partial W; \mathbb{Z})$, and we write $[\hat{\Sigma}_i]$ for the lifts under the map $H_2(W; \mathbb{Q}) \rightarrow H_2(W, \partial W; \mathbb{Q})$.

This may be seen to coincide with the definition in terms of Alexander gradings on $\widehat{HFL}(Y, L)$ using a small modification of the cobordism argument from [57, Section 5.5] (which is stated for integrally null-homologous links). See [17, Section 3.2] for an exposition in the setting of rationally null-homologous knots. It is an easy consequence of the cobordism description of the Alexander grading that $\widehat{CFL}(Y, L)$ is supported on $\mathbb{H}(Y, L)$.

Lemma 6.2. *There is an isomorphism*

$$\mathbb{H}(Y, L) \cong H_1(Y \setminus L; \mathbb{Z})$$

of affine spaces over $H_1(Y \setminus L; \mathbb{Z})$.

Proof. This follows from the short exact sequence of affine spaces

$$0 \rightarrow \mathbb{Z}^\ell \rightarrow \mathbb{H}(Y, L) \rightarrow \text{Spin}^c(Y) \rightarrow 0 \quad (6.2)$$

which is parallel to the short exact sequence of homology groups

$$0 \rightarrow \mathbb{Z}^\ell \rightarrow H_1(Y \setminus L; \mathbb{Z}) \rightarrow H_1(Y; \mathbb{Z}) \rightarrow 0.$$

If we pick a base element $(\mathbf{s}, \mathbf{t}) \in \mathbb{H}(Y, L)$, we obtain a map of affine spaces from $H_1(Y \setminus L; \mathbb{Z})$ to $\mathbb{H}(Y, L)$ which makes the natural diagram commute. By the five-lemma, we conclude that $\mathbb{H}(Y, L)$ and $H_1(Y \setminus L; \mathbb{Z})$ are isomorphic as affine spaces. ■

We are now ready to define the H -function of a link in a rational homology sphere.

Definition 6.3. For an oriented link $L \subseteq Y$ in a rational homology sphere Y and $(s, t) \in \mathbb{H}(Y, L)$, we define $H_L: \mathbb{H}(Y, L) \rightarrow \mathbb{Q}$ by declaring that $-2H_L(s, t)$ is the maximal gr_w -grading of a non-zero element in the free part of $H_*(\mathfrak{A}^-(L, s, t))$, where $\mathfrak{A}^-(L, s, t)$ is the subcomplex of $\mathcal{CFL}(Y, L, t)$ lying in Alexander grading s .

The H -function of L-space links in the 3-sphere can be computed from Alexander polynomials of the link and all sublinks [2, 11]. In order to generalize the result to links in a rational homology sphere, we first recall *generalized Alexander polynomials*. Friedl, Juhász and Rasmussen [7, Theorem 1] proved that $\widehat{HFL}(Y, L)$ categorifies the Turaev torsion $\Delta(Y, L)$ of $Y \setminus L$, which we view as an element in $\mathbb{F}[H_1(Y \setminus L)]$, well-defined up to multiplication by monomials. When Y is a rational homology 3-sphere, we refer to $\Delta(Y, L)$ as the *generalized Alexander polynomial*.

In our present setting, it is helpful to view $\Delta(Y, L)$ as taking values in the $\mathbb{F}[H_1(Y \setminus L)]$ -module $\mathbb{F}[\mathbb{H}(Y, L)]$ instead of $\mathbb{F}[H_1(Y \setminus L)]$ itself, via the following formula:

$$\Delta(Y, L) = \chi(\widehat{HFL}(Y, L)) := \sum_{(s,t) \in \mathbb{H}(Y,L)} t^{(s,t)} \cdot \chi(\widehat{HFL}(Y, L, s, t)) \in \mathbb{F}[\mathbb{H}(Y, L)].$$

We may think of $\chi(\widehat{HFL}(Y, L))$ as a normalized version of the Turaev torsion. Similarly to [48, Proposition 8.1], it is not hard to see that the Euler characteristic of $\widehat{HFL}(Y, L)$ is symmetric with respect to the involution of $\mathbb{H}(Y, L)$ given by $(s, t) \mapsto (-s, \bar{t} + PD[L])$ (see [57, Proposition 8.3]).

To make more transparent connections with integer homology spheres, we refine the definition of $\Delta(Y, L)$. For $t \in \text{Spin}^c(Y)$, we set

$$\begin{aligned} \Delta(Y, L, t) &= \chi(\widehat{HFL}(Y, L, t)) \\ &:= \sum_{s: (s,t) \in \mathbb{H}(Y,L)} t^s \cdot \chi(\widehat{HFL}(Y, L, s, t)) \in \mathbb{F}[\mathbb{H}(Y, L)]. \end{aligned} \quad (6.3)$$

We recall that the graded Euler characteristics of $\widehat{HFL}$ and HFL^- are related by the formula

$$\chi(HFL^-(Y, L)) = \frac{1}{\prod_{i=1}^{\ell} (1 - t_i)} \chi(\widehat{HFL}(Y, L))$$

(see [48, Proposition 9.2]).

For any sublink $L' \subseteq L$, define the natural forgetful map

$$\pi_{L, L-L'}: \mathbb{H}(Y, L) \rightarrow \mathbb{H}(Y, L - L')$$

as follows. If $L' = K_1$, the map $\pi_{L, L-L'}$ is given by the formula

$$\pi_{L, L-L'}(s_1, \dots, s_{\ell}, t) = \left(s_2 - \frac{\text{lk}(K_1, K_2)}{2}, \dots, s_{\ell} - \frac{\text{lk}(K_1, K_{\ell})}{2}, t \right),$$

where $L = K_1 \cup \dots \cup K_{\ell}$. For general L' , the map $\pi_{L, L-L'}$ is a composition of several such maps. We refer the reader to [30, Section 3.7] for explicit formulas for the forgetful map for links in S^3 . Compare [2, Section 3].

Given $(\mathbf{s}, \mathbf{t}), (\mathbf{s}', \mathbf{t}') \in \mathbb{H}(Y, L)$, we write $(\mathbf{s}, \mathbf{t}) \geq (\mathbf{s}', \mathbf{t}')$ if and only if $\mathbf{t} = \mathbf{t}'$ and $\mathbf{s} \geq \mathbf{s}'$, that is, $s_i \geq s'_i$ for all i , where $\mathbf{s} = (s_1, \dots, s_\ell)$ and $\mathbf{s}' = (s'_1, \dots, s'_\ell)$.

Lemma 6.4. *For an oriented L-space link $L \subseteq Y$ in a rational homology sphere with $\mathbf{t} \in \text{Spin}^c(Y)$, the H -function H_L satisfies*

$$H_L(\mathbf{s}, \mathbf{t}) = \sum_{L' \subseteq L} (-1)^{|L'| - 1} \sum_{\substack{(\mathbf{s}', \mathbf{t}) \in \mathbb{H}(Y, L') \\ (\mathbf{s}', \mathbf{t}) \geq \pi_{L, L'}(\mathbf{s} + \mathbf{1}, \mathbf{t})}} \chi(HFL^-(Y, L', \mathbf{s}', \mathbf{t})), \quad (6.4)$$

where $\mathbf{1} = (1, \dots, 1)$. In particular, the H -function is determined by the generalized Alexander polynomials $\Delta(Y, L', \mathbf{t})$ of all sublinks $L' \subseteq L$.

Proof. The arguments of [11, Theorem 2.10], which relate H_L to $\chi(HFL^-(Y, L))$ when $Y = S^3$, can be repeated verbatim to obtain (6.4) for a general L-space link L in a rational homology 3-sphere Y , though here we follow the convention in [2, Theorem 3.15] and assume $HFL^-(Y, \emptyset) = 0$. The right-hand side of (6.4) is determined by $\chi(\widehat{HFL}(Y, L'))$ for all sublinks L' , which can be computed from Alexander polynomials of L' by (6.3). Therefore, the H -function is determined by the Alexander polynomials of all sublinks $L' \subseteq L$. ■

Example 6.5. We normalize the multivariable Alexander polynomial of the unlink $\mathbb{U}$ in the 3-sphere to be 0, and the H -function for an ℓ -component unlink in S^3 is the following:

$$H_{\mathbb{U}_\ell}(s_1, \dots, s_\ell) = \sum_i (|s_i| - s_i)/2.$$

If $L \subseteq S^3$ is the Hopf link in the 3-sphere, its Alexander polynomial $\Delta(t_1, t_2)$ is 1 and $s_i \in \mathbb{Z} + 1/2$. By the formula in [2, 28], the value of the H -function at each lattice point (s_1, s_2) is shown as follows:




Fig. 6.1. The H -function of the Hopf link

Theorem 6.6. *Suppose that Γ is an arrow-decorated plumbing tree with $V_\Gamma = V_G \cup V_\uparrow$. Let $L_\uparrow \subseteq Y_G$ be the associated link and assume that Y_G is a rational homology 3-sphere.*

If $L_\uparrow$ is an L -space link, then the full link Floer complex $\mathcal{CFL}(Y_G, L_\uparrow)$ is determined by the generalized Alexander polynomials of $L_\uparrow$ and its sublinks.

Proof. By Theorem 6.1, the chain complex $\mathcal{CFL}(Y_G, L_\uparrow)$ is a free resolution of $\mathcal{HFL}(Y_G, L_\uparrow)$. In particular, $\mathcal{CFL}(Y_G, L_\uparrow)$ is determined by the homology group $\mathcal{HFL}(Y_G, L_\uparrow)$ as an $\mathcal{R}_\ell$ -module. Therefore, it suffices to prove that the generalized Alexander polynomials determine the homology $\mathcal{HFL}(Y_G, L_\uparrow)$. By Lemma 6.4, the H -function is determined by the generalized Alexander polynomials of $L_\uparrow$ and its sublinks, and it remains to prove that the H -function determines the $\mathcal{R}_\ell$ -module structure of $\mathcal{HFL}(Y_G, L_\uparrow)$.

There is a decomposition of $\mathbb{F}[U]$ -modules

$$\mathcal{HFL}(Y_G, L_\uparrow) = \bigoplus_{(\mathbf{s}, \mathbf{t}) \in \mathbb{H}(Y_G, L_\uparrow)} \mathcal{HFL}(Y_G, L_\uparrow, \mathbf{s}, \mathbf{t}).$$

Since $L_\uparrow$ is an L -space link in Y_G , $\mathcal{HFL}(Y_G, L_\uparrow, \mathbf{s}, \mathbf{t}) \cong \mathbb{F}[U]$ for all $(\mathbf{s}, \mathbf{t}) \in \mathbb{H}(Y_G, L_\uparrow)$. The (gr_w, A) -grading of the generator of $\mathcal{HFL}(Y_G, L_\uparrow, \mathbf{s}, \mathbf{t})$ is determined by the H -function. That is, the Alexander grading of the generator equals $\mathbf{s}$ and the Maslov grading gr_w equals $-2H_{L_\uparrow}(\mathbf{s}, \mathbf{t})$. Hence, it suffices to see that the $\mathcal{U}_i$ and $\mathcal{V}_i$ actions on $\mathcal{HFL}(Y_G, L_\uparrow, \mathbf{s}, \mathbf{t})$ are also determined by the H -function.

To see this, note that we may view $\mathcal{U}_i$ as restricting to a map

$$\mathcal{HFL}(Y_G, L_\uparrow, \mathbf{s}, \mathbf{t}) \cong \mathbb{F}[U] \rightarrow \mathcal{HFL}(Y_G, L_\uparrow, \mathbf{s} - e_i, \mathbf{t}) \cong \mathbb{F}[U]. \quad (6.5)$$

Since the map $\mathcal{V}_i$ goes in the opposite direction and $\mathcal{U}_i \mathcal{V}_i = U$, the map $\mathcal{U}_i$ is given by multiplication by either 1 or U , with respect to identifications in (6.5). We observe that the map $\mathcal{U}_i$ has gr_w -grading -2 , and hence the choice of being U or 1 is determined by the gr_w -gradings of the copies of $\mathbb{F}[U]$ in (6.5), which is encoded by the H -function. Similarly, the action of $\mathcal{V}_i$ is also determined by the H -function since $\mathcal{U}_i \mathcal{V}_i = U$, and the action of $\mathcal{U}_i$ is determined by the H -function. Therefore, for L -space links, the Alexander polynomials determine the full link Floer chain complexes. ■

6.3. Comparison with Gorsky and Némethi's link lattice homology

We now compare our chain complex $\mathbb{CFL}(\Gamma, V_\uparrow)$ with the definition of link lattice homology due to Gorsky and Némethi [11].

As a first step, we recall their definition. Let $L \subseteq S^3$ be a link of ℓ components. Write C_i for the mapping cone complex

$$C_i := \text{Cone}(\mathcal{V}_i: \mathcal{R}_\ell \rightarrow \mathcal{R}_\ell), \quad i = 1, \dots, \ell.$$

Write $\mathcal{K}(L)$ for the (non-free) complex of $\mathcal{R}_\ell$ -modules obtained by tensoring the homology group $\mathcal{HFL}(L)$ with $C_1, \dots, C_\ell$ over the ring $\mathcal{R}_\ell$.

Remark 6.7. Note that $C_1 \otimes_{\mathcal{R}_\ell} \dots \otimes_{\mathcal{R}_\ell} C_\ell$ is the Koszul complex for the regular sequence $(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$ in $\mathcal{R}_\ell$ (see [54, Section 4.5]).

If L is an L-space link, Gorsky and Némethi define the link lattice complex to be the chain complex $\mathcal{K}(L)$. They prove the following:

Theorem 6.8 ([11, Theorem 2.9]). *If L is an algebraic link, then*

$$H_*\mathcal{K}(L) \cong H_*(\mathcal{CFL}(L)/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)).$$

Remark 6.9. Gorsky and Némethi prove Theorem 6.8 at the level of graded $\mathbb{F}$ -vector spaces. There are additional module actions of $\mathbb{F}[[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]]$ on both sides. Gorsky and Hom [9, Proposition 3.7] equip $\mathcal{K}(L)$ with commuting endomorphisms $\mathcal{U}_1, \dots, \mathcal{U}_\ell$. In our notation, their action of $\mathcal{U}_i$ coincides with the standard action of $\mathcal{U}_i$ on the $\mathcal{HFL}(L)$ tensor factor of $\mathcal{K}(L)$. Of course, one can also consider the action of $\mathcal{V}_i$, defined symmetrically. Note that as an endomorphism of $\mathcal{K}(L)$, each $\mathcal{V}_i$ is null-homotopic since $\mathcal{K}(L)$ is defined by tensoring $\mathcal{HFL}(L)$ with the Koszul complex of the sequence $(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$. More explicitly, we can write $\mathcal{K}(L)$ as a mapping cone

$$\mathcal{K}(L) = \text{Cone}(\mathcal{K}_i^0(L) \xrightarrow{\mathcal{V}_i} \mathcal{K}_i^1(L)),$$

where $\mathcal{K}_i^0(L)$ (resp. $\mathcal{K}_i^1(L)$) is the codimension 1 subcube which has i -coordinate 0 (resp. 1). As an endomorphism of $\mathcal{K}(L)$, the module action of $\mathcal{V}_i$ preserves both $\mathcal{K}_i^0(L)$ and $\mathcal{K}_i^1(L)$. We define an endomorphism H of $\mathcal{K}(L)$, which sends $\mathcal{K}_i^1(L)$ to $\mathcal{K}_i^0(L)$ via the identity, and observe that on $\mathcal{K}(L)$,

$$\mathcal{V}_i = [\partial, H].$$

We now explain how our Theorem 6.1 quickly recovers Theorem 6.8, and also prove a refinement which takes into account the $\mathcal{R}_\ell$ -action. To state our result, we equip $\mathcal{K}(L)$ with an A_∞ -module structure which has only m_1 and m_2 non-trivial. The action of m_2 corresponds to the standard action on the $\mathcal{HFL}(L)$ factor of the tensor product.

Theorem 6.10. *If $L \subseteq S^3$ is a plumbed L-space link, then there is a homotopy equivalence of A_∞ -modules over $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]$,*

$$\mathcal{K}(L) \simeq \mathcal{CFL}(L)/(\mathcal{V}_1, \dots, \mathcal{V}_\ell).$$

Proof. We first consider the proof only at the level of chain complexes, and then consider the $\mathcal{R}_\ell$ -module structure.

Note that C_i is a free resolution of $\mathcal{R}_\ell/\mathcal{V}_i$ as an $\mathcal{R}_\ell$ -module. Furthermore, the ℓ -dimensional cube-shaped complex $C_1 \otimes \dots \otimes C_\ell$ is a free resolution of $\mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_\ell, \mathcal{V}_\ell]/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$.

We use the algebraic formalism of type- D , type- A and type- DA modules of Lipshitz, Ozsváth and Thurston [24, 25]. Their algebraic formalism gives a convenient model of the derived tensor product, called the *box tensor product*, which is denoted by $\boxtimes$. See [24, Section 2.3.2] for more details. We may view C_i as a type- D module ${}^{\mathcal{R}_\ell}C_i$ whose underlying $\mathbb{F}$ -vector space has two generators, $\mathbf{x}_i$ and $\mathbf{y}_i$, and whose structure map δ^1 is given by

$$\delta^1(\mathbf{x}_i) = \mathcal{V}_i \otimes \mathbf{y}_i.$$

Similarly, the complex $C_1 \otimes \cdots \otimes C_\ell$ naturally corresponds to a type- D module over $\mathcal{R}_\ell$, whose underlying vector space is generated by the points of an ℓ -dimensional cube. We write ${}^{\mathcal{R}_\ell}C_{1,\dots,\ell}$ for this cube-shaped complex.

By definition,

$$\mathcal{H}(L) := \mathcal{HFL}(L)_{\mathcal{R}_\ell} \boxtimes {}^{\mathcal{R}_\ell}C_{1,\dots,\ell}.$$

By Theorem 6.1, if L is a plumbed L-space link, $\mathcal{HFL}(L)$ and $\mathcal{CFL}(L)$ are homotopy equivalent as A_∞ -modules over $\mathcal{R}_\ell$. On the other hand, $\mathcal{CFL}(L)_{\mathcal{R}_\ell}$ is free over $\mathcal{R}_\ell$, which translates to the fact that there is a type- D module $\mathcal{CFL}(L)^{\mathcal{R}_\ell}$ such that

$$\mathcal{CFL}(L)_{\mathcal{R}_\ell} \cong \mathcal{CFL}(L)^{\mathcal{R}_\ell} \boxtimes_{{\mathcal{R}_\ell}} \mathcal{R}_\ell.$$

We observe that since ${}^{\mathcal{R}_\ell}C_{1,\dots,\ell}$ is a free resolution of $\mathcal{R}_\ell/(\mathcal{V}_1, \dots, \mathcal{V}_\ell)$, it follows that

$${}_{\mathcal{R}_\ell}C_{1,\dots,\ell} \simeq \mathcal{R}_\ell \mathcal{R}_\ell / (\mathcal{V}_1, \dots, \mathcal{V}_\ell).$$

Putting these relations together, we obtain

$$\begin{aligned} \mathcal{H}(L) &= \mathcal{HFL}(L)_{\mathcal{R}_\ell} \boxtimes {}^{\mathcal{R}_\ell}C_{1,\dots,\ell} \\ &\simeq \mathcal{CFL}(L)_{\mathcal{R}_\ell} \boxtimes {}^{\mathcal{R}_\ell}C_{1,\dots,\ell} \\ &= \mathcal{CFL}(L)^{\mathcal{R}_\ell} \boxtimes_{{\mathcal{R}_\ell}} \mathcal{R}_\ell \mathcal{R}_\ell \boxtimes {}^{\mathcal{R}_\ell}C_{1,\dots,\ell} \\ &\simeq \mathcal{CFL}(L)^{\mathcal{R}_\ell} \boxtimes_{{\mathcal{R}_\ell}} \mathcal{R}_\ell \mathcal{R}_\ell / (\mathcal{V}_1, \dots, \mathcal{V}_\ell). \end{aligned} \tag{6.6}$$

We now consider the $\mathcal{R}_\ell$ -module structures. We note that the complex ${}^{\mathcal{R}_\ell}C_{1,\dots,\ell}$ extends to a DA -bimodule ${}^{\mathcal{R}_\ell}(C_{1,\dots,\ell})_{\mathcal{R}_\ell}$. This DA -bimodule has the same generators and the same structure map δ^1 as ${}^{\mathcal{R}_\ell}C_{1,\dots,\ell}$. Additionally, there is a δ_2^1 action given by $\delta_2^1(\mathbf{x}, a) = a \otimes \mathbf{x}$ for any $a \in \mathcal{R}_\ell$ and $\mathbf{x} \in C_{1,\dots,\ell}$. We observe that by definition,

$$\mathcal{H}(L)_{\mathcal{R}_\ell} = \mathcal{HFL}(L)_{\mathcal{R}_\ell} \boxtimes {}^{\mathcal{R}_\ell}(C_{1,\dots,\ell})_{\mathcal{R}_\ell}.$$

Additionally, it is easy to check that

$${}_{\mathcal{R}_\ell} \mathcal{R}_\ell \mathcal{R}_\ell \boxtimes {}^{\mathcal{R}_\ell}(C_{1,\dots,\ell})_{\mathcal{R}_\ell} \simeq \mathcal{R}_\ell (\mathcal{R}_\ell / (\mathcal{V}_1, \dots, \mathcal{V}_\ell))_{\mathcal{R}_\ell},$$

so that the manipulation from (6.6) extends to an equivalence of A_∞ -modules. $\blacksquare$

7. Computations

In this section, we give some computational tools. In Section 7.1, we provide an algorithm to compute the link Floer homology of an L-space link from its H -function. Next, in Section 7.2, we describe the $\mathcal{R}_n$ -module $\mathcal{HFL}(T(n, n))$ based on Gorsky and Hom's computation of the H -function of $T(n, n)$. By our Theorem 6.1, this $\mathcal{R}_n$ -module contains equivalent information to $\mathcal{CFL}(T(n, n))$. In Sections 7.3 and 7.4, we compute explicit free resolutions of $\mathcal{HFL}(T(3, 3))$ and $\mathcal{HFL}(T(4, 4))$.

7.1. Generators and relations for the homology of an L-space link

In this section, we describe generators and relations for the modules $\mathcal{HFL}(L)$ when L is an L-space link. We focus on the case $L \subseteq S^3$ to simplify the notation, but this is not essential.

If $\mathbf{s} \in \mathbb{H}(L)$, we write $X_{\mathbf{s}} \in \mathcal{HFL}(L)$ for the generator of the $\mathbb{F}[U]$ -tower in Alexander grading $\mathbf{s}$. By definition,

$$A(X_{\mathbf{s}}) = \mathbf{s} \quad \text{and} \quad \text{gr}_w(X_{\mathbf{s}}) = -2H_L(\mathbf{s}).$$

We let $\mathcal{G}$ denote the set of $X_{\mathbf{s}} \in \mathcal{HFL}(L)$ which satisfy

$$H_L(\mathbf{s} - e_i) - H_L(\mathbf{s}) = 1 \quad \text{and} \quad H_L(\mathbf{s}) - H_L(\mathbf{s} + e_i) = 0, \quad (7.1)$$

for all $i \in \{1, \dots, \ell\}$. Here, e_i denotes the standard unit vector.

Lemma 7.1. *Let L be an ℓ -component L-space link in S^3 .*

- (1) *The set $\mathcal{G}$ is finite and is the unique $\mathcal{R}_{\ell}$ -module generating set of $\mathcal{HFL}(L)$ of minimal length.*
- (2) *The kernel of the natural map*

$$\mathcal{R}_{\ell}^n \rightarrow \mathcal{HFL}(L)$$

is spanned by the following generating set:

(r-1) *All elements of the form*

$$\alpha X_{\mathbf{s}} + \beta X_{\mathbf{s}'},$$

where $X_{\mathbf{s}}, X_{\mathbf{s}'} \in \mathcal{G}$, and α, β are monomials such that $w_U(\alpha) = w_U(\beta) = 0$ and $\gcd(\alpha, \beta) = 1$ with

$$A(\alpha) + \mathbf{s} = A(\beta) + \mathbf{s}' \quad \text{and} \quad \text{gr}_w(\alpha) - 2H_L(\mathbf{s}) = \text{gr}_w(\beta) - 2H_L(\mathbf{s}').$$

We use the notation $w_U(\alpha) = \min\{i_1, j_1\} + \dots + \min\{i_{\ell}, j_{\ell}\}$ for any monomial $\alpha = \mathcal{U}_1^{i_1} \dots \mathcal{U}_{\ell}^{i_{\ell}} \mathcal{V}_1^{j_1} \dots \mathcal{V}_{\ell}^{j_{\ell}}$.

(r-2) *For each i, j and $X_{\mathbf{s}} \in \mathcal{G}$ the element*

$$(\mathcal{U}_i \mathcal{V}_i + \mathcal{U}_j \mathcal{V}_j) X_{\mathbf{s}}.$$

Proof. Since $\mathcal{HFL}(L)$ is a finitely generated $\mathcal{R}_{\ell}$ -module, and $\mathcal{HFL}(L)$ admits Maslov and Alexander gradings, the $\mathcal{R}_{\ell}$ -module $\mathcal{HFL}(L)$ is spanned by finitely many homogeneously graded vectors $X_{\mathbf{s}}$. Equation (7.1) is equivalent to the statement that $X_{\mathbf{s}}$ is in the image of neither $\mathcal{U}_i$ nor $\mathcal{V}_i$ for any $i \in \{1, \dots, \ell\}$. Therefore any minimal length generating set can only contain elements of $\mathcal{G}$. In particular, $\mathcal{G}$ generates $\mathcal{HFL}(L)$.

We now claim that any minimal length generating set must contain all elements of $\mathcal{G}$. To see this, we write $\mathcal{G} = \{X_{\mathbf{s}_1}, \dots, X_{\mathbf{s}_n}\}$ and suppose

$$X_{\mathbf{s}_j} = \sum_{i \neq j} \alpha_i X_{\mathbf{s}_i}$$

for some homogeneously graded $\alpha_i \in \mathcal{R}_\ell$. We may assume each $\alpha_i X_{s_i}$ has the same gr_w and Alexander gradings as X_{s_j} . However, $\mathcal{HFL}(L)$ is rank 1 over $\mathbb{F}$ in each Maslov and Alexander grading in which it is supported. Therefore if $\alpha_i \neq 0$, then $X_{s_j} = \alpha_i X_{s_i}$. However, (7.1) implies that $\alpha_i = 1$ since none of the X_{s_i} admit non-trivial factorizations. Therefore every minimal length generating set contains all of $\mathcal{G}$.

We now consider the claim about relations. Suppose that

$$\sum_{i=1}^n \alpha_i X_{s_i} = 0.$$

We may assume that each α_i is a monomial (or zero) and all $\alpha_i X_{s_i}$ have the same Alexander and Maslov gradings. When all of the $\alpha_i X_{s_i}$ have the same homogeneous Alexander and Maslov gradings, $\sum_{i=1}^n \alpha_i X_{s_i} = 0$ if and only if $\#\{i : \alpha_i \neq 0\}$ is even, since $\mathcal{HFL}(L)$ has rank 1 over $\mathbb{F}$ in each of the gradings in which it is supported. In particular, pairing elements of $\{i : \alpha_i \neq 0\}$ arbitrarily, we may write $\sum_{i=1}^n \alpha_i X_{s_i}$ as an $\mathcal{R}_\ell$ -linear combination of elements of the form $\alpha X_s + \beta X_{s'}$, where $\alpha, \beta \neq 0$ and αX_s and $\beta X_{s'}$ have the same Maslov and Alexander gradings. By canceling common factors, we may assume that $\gcd(\alpha, \beta) = 1$.

If $\max(w_U(\alpha), w_U(\beta)) > 0$, we claim that we can still reduce this relation further. For concreteness, assume that $w_U(\alpha) > 0$. This means that $\mathcal{U}_j \mathcal{V}_j$ divides α for some j . Note that $\beta \neq 1$, so assume that either $\mathcal{V}_i$ or $\mathcal{U}_i$ divides β , for some i . By using (r-2), we may replace the factor of $\mathcal{U}_j \mathcal{V}_j$ in α with $\mathcal{U}_i \mathcal{V}_i$, and assume $\mathcal{U}_i \mathcal{V}_i$ divides α . The resulting monomials α and β have $\gcd(\alpha, \beta) \neq 1$, so we divide out common factors. This operation is non-increasing in the total degree of α and β , so we may repeat this procedure until $\gcd(\alpha, \beta) = 1$ and $w_U(\alpha) = w_U(\beta) = 0$. This completes the proof. ■

Remark 7.2. We can describe the relations (r-2) more concretely as follows. Let $X_s, X_{s'} \in \mathcal{G}$ and write

$$\mathbf{s} = (s_1, \dots, s_\ell) \quad \text{and} \quad \mathbf{s}' = (s'_1, \dots, s'_\ell).$$

We let

$$\mathcal{P}_i = \begin{cases} \mathcal{V}_i & \text{if } s_i < s'_i, \\ \mathcal{U}_i & \text{if } s_i > s'_i, \end{cases} \quad \mathcal{Q}_i = \begin{cases} \mathcal{U}_i & \text{if } s_i < s'_i, \\ \mathcal{V}_i & \text{if } s_i > s'_i. \end{cases}$$

If $s_i = s'_i$, then we view $\mathcal{P}_i = \mathcal{Q}_i = 1$. The relations labeled (r-2) may be rewritten in the form

$$\mathcal{P}_1^{i_1} \dots \mathcal{P}_\ell^{i_\ell} \cdot X_s = \mathcal{Q}_1^{j_1} \dots \mathcal{Q}_\ell^{j_\ell} \cdot X_{s'}$$

ranging over all $X_s, X_{s'} \in \mathcal{G}$ and all sequences $i_1, \dots, i_\ell, j_1, \dots, j_\ell \geq 0$ such that

$$i_k + j_k = |s_k - s'_k| \quad \text{and} \quad \text{gr}_w(\mathcal{P}_1^{i_1} \dots \mathcal{P}_\ell^{i_\ell} \cdot X_s) = \text{gr}_w(\mathcal{Q}_1^{j_1} \dots \mathcal{Q}_\ell^{j_\ell} \cdot X_{s'}),$$

for all k . Note in particular that there are only finitely many relations of the form (r-2).

7.2. The type- A module of $T(n, n)$

We now compute the Heegaard Floer module $\mathcal{HFL}(T(n, n))$ of the torus link $T(n, n)$ in the 3-sphere. Recall that $\mathcal{R}_n = \mathbb{F}[\mathcal{U}_1, \mathcal{V}_1, \dots, \mathcal{U}_n, \mathcal{V}_n]$.

Theorem 7.3. *As an $\mathcal{R}_n$ -module, the group $\mathcal{HFL}(T(n, n))$ has a unique minimal generating set, consisting of n generators, $X_1, \dots, X_n$. The relations are spanned by the following:*

$$\left(\prod_{i \in I_k} \mathcal{U}_i\right) X_k = \left(\prod_{j \in \{1, \dots, n\} \setminus I_k} \mathcal{V}_j\right) X_{k+1}, \quad (7.2)$$

$$\mathcal{U}_i \mathcal{V}_i X_k = \mathcal{U}_j \mathcal{V}_j X_k. \quad (7.3)$$

Here I_k runs through all subsets of $\{1, \dots, n\}$ of cardinality k (so (7.2) has $\binom{n}{k}$ relations for each k), and in (7.3), i, j , and k range from 1 to n .

Proof. Since $T(n, n)$ is an L-space link, by Lemma 7.1 the minimal generating set consists of the set of all X_s which satisfy

$$H(\mathbf{s} - e_i) - H(\mathbf{s}) = 1 \quad \text{and} \quad H(\mathbf{s}) - H(\mathbf{s} + e_i) = 0, \quad (7.4)$$

for each $i \in \{1, \dots, n\}$. Here, $H(\mathbf{s})$ denotes the H -function of $T(n, n)$.

The latter H -function is computed in [9]. Its Alexander polynomial equals

$$\Delta(t_1, \dots, t_n) = ((t_1 \cdots t_n)^{1/2} - (t_1 \cdots t_n)^{-1/2})^{n-2},$$

which is symmetric in $t_1, \dots, t_n$. Hence the H -function for $T(n, n)$ is also symmetric in $s_1, \dots, s_n$. Therefore we consider the case $s_1 \leq \dots \leq s_n$. By [9, Theorem 4.3],

$$H(s_1, \dots, s_n) = h\left(s_1 - \frac{n-1}{2}\right) + h\left(s_2 - \frac{n-1}{2} + 1\right) + \dots + h\left(s_n - \frac{n-1}{2} + n-1\right), \quad (7.5)$$

where $h(s) = (|s| - s)/2$. We note that $h(s)$ is the H -function of the unknot. It is not hard to compute that

$$H(s_1, \dots, s_n) = 0 \quad (7.6)$$

if $s_i \geq (n-1)/2$ for all i , and

$$H(\mathbf{s}) = -(s_1 + \dots + s_n) \quad (7.7)$$

if $s_i \leq -(n-1)/2$ for all i .

We first consider the diagonal vertices, that is, $s_1 = \dots = s_n = m$, $m \in \mathbb{Z} + (n-1)/2$. If $m > (n-1)/2$, then

$$H(s_1, \dots, s_n) = H(s_1 - 1, s_2, \dots, s_n) = 0$$

by (7.6), which does not satisfy (7.4). Similarly, if $m < -(n-1)/2$, then

$$H(s_1, \dots, s_n) = H(s_1, \dots, s_n + 1) + 1$$

by (7.7), which does not satisfy (7.4) either. Now we consider the n diagonal vertices where $-(n-1)/2 \leq m \leq (n-1)/2$, that is,

$$s_1 = \dots = s_n = \frac{n+1}{2} - k$$

for all integers k between 1 and n . The corresponding generators are $X_1, \dots, X_n$ with Alexander gradings

$$\left(\frac{n+1}{2} - k, \frac{n+1}{2} - k, \dots, \frac{n+1}{2} - k \right).$$

By a straightforward computation, (7.4) is satisfied by values of the H -function at these vertices, so $X_1, \dots, X_n$ are all contained in the minimal generating set from Lemma 7.1. We claim these diagonal vertices are the only ones where the H -function satisfies (7.4). It suffices to prove that the non-diagonal vertices do not satisfy (7.4). Recall that we assume $s_1 \leq \dots \leq s_n$. Suppose that $s_n = s$ is the maximal value among the s_i , and there are exactly λ coordinates equal to s where $\lambda < n$, i.e., $s_{n-\lambda+1} = s_{n-\lambda+2} = \dots = s_n = s$, and $s_{n-\lambda} < s$.

If $s > (\lambda - 1) - (n - 1)/2$, then by (7.5),

$$H(s_1, \dots, s_{n-\lambda}, s_{n-\lambda+1} - 1, \dots, s_n) = H(s_1, \dots, s_n),$$

which does not satisfy (7.4).

If $s \leq (\lambda - 1) - (n - 1)/2$, then by (7.5),

$$H(s_1, \dots, s_{n-\lambda} + 1, s_{n-\lambda+1}, \dots, s_n) + 1 = H(s_1, \dots, s_n),$$

which does not satisfy (7.4) either. By Lemma 7.1, $X_1, \dots, X_n$ form a unique minimal generating set of $\mathcal{HFL}(T(n, n))$ over $\mathcal{R}_n$. Based on the values of the H -function, one can compute that

$$\text{gr}_w(X_k) = -k(k-1).$$

We now consider the relations satisfied by $X_1, \dots, X_n$ over $\mathcal{R}_n$. For convenience, if $I = (i_1, \dots, i_n)$ is a sequence of non-negative numbers, write $\mathcal{U}^I$ for $\mathcal{U}_1^{i_1} \dots \mathcal{U}_n^{i_n}$, and define $\mathcal{V}^J$ similarly. Lemma 7.1 and Remark 7.2 immediately imply that if $1 \leq p < q \leq n$, then the relations involving X_p and X_q are spanned by $(\mathcal{U}_i \mathcal{V}_i + \mathcal{U}_j \mathcal{V}_j)X_p$ and $(\mathcal{U}_i \mathcal{V}_i + \mathcal{U}_j \mathcal{V}_j)X_q$ for $i, j \in \{1, \dots, n\}$, as well as the sums

$$\mathcal{U}^I X_p + \mathcal{V}^J X_q$$

ranging over sequences of non-negative integers I and J such that

$$I + J = (q - p, \dots, q - p) \quad \text{and} \quad |I|_{L^1} = \sum_{s=p}^{q-1} s = \frac{(p+q-1)(q-p)}{2}. \quad (7.8)$$

If $q = p + 1$, these are exactly the relations in the statement.

We will show by induction on q that the relations between X_p and X_q are in the span of the relations between consecutive X_i and X_{i+1} , as well as the relations

$$(\mathcal{U}_i \mathcal{V}_i + \mathcal{U}_j \mathcal{V}_j) X_k = 0$$

(labeled (r-2) above). The case $q = p + 1$ is vacuous, so we suppose that $q > p + 1$.

We let $I = (I_1, \dots, I_n)$ and $J = (J_1, \dots, J_n)$ be tuples of non-negative integers such that $I + J = (q - p, \dots, q - p)$ and $|I|_{L^1} = p + \dots + q - 1$. We claim first that there is a tuple I' such that

$$0 \leq I' \leq I, \quad |I'|_{L^1} = q - 1, \quad \text{and} \quad |I'|_{L^\infty} \leq 2. \quad (7.9)$$

(We remark that it may not be possible to find such an I' which satisfies $|I'|_{L^\infty} \leq 1$.) For $k \in \mathbb{N}$, we write

$$a_k := \#\{i : I_i \geq k\}.$$

We observe that

$$|I|_{L^1} = \sum_{k=1}^{q-p} a_k = \frac{(p+q-1)(q-p)}{2} \quad \text{and} \quad a_1 \geq \dots \geq a_{q-p}. \quad (7.10)$$

Our claim about the existence of an I' as above is equivalent to $a_1 + a_2 \geq q - 1$. If, to the contrary, $a_1 + a_2 < q - 1$, then we observe that $a_2 < (q - 1)/2$ and therefore

$$\sum_{k=1}^{q-p} a_k = a_1 + a_2 + \sum_{n=3}^{q-p} a_k < q - 1 + (q - p - 2) \frac{q - 1}{2} = \frac{(q - p)(q - 1)}{2}.$$

However, this contradicts (7.10), so we conclude that an I' satisfying (7.9) must exist.

By induction, if J' denotes $(q - p - 1, \dots, q - p - 1) - (I - I')$, then the relation

$$\mathcal{U}^{I-I'} X_p = \mathcal{V}^{J'} X_{q-1} \quad (7.11)$$

is in the span of the claimed relations between consecutive X_i and X_{i+1} , as well as the relations (r-2). We observe that (7.11) implies

$$\mathcal{U}^I X_p = \mathcal{U}^{I'} \mathcal{U}^{I-I'} X_p = \mathcal{U}^{I'} \mathcal{V}^{J'} X_{q-1}.$$

If $|I'|_{L^\infty} = 1$, then we have the relation

$$\mathcal{U}^{I'} X_{q-1} = \mathcal{V}^{(1, \dots, 1) - I'} X_q,$$

which implies with (7.11) that $\mathcal{U}^I X_p = \mathcal{V}^J X_q$, which would complete the proof.

If instead $|I'|_{L^\infty} = 2$, then we observe that any coordinate i such that $I'_i = 2$ has the property that $(I - I')_i < q - p - 1$ and hence $J'_i > 0$. In particular, $\mathcal{U}^{I'} \mathcal{V}^{J'}$ has a factor $\mathcal{U}_i \mathcal{V}_i$ for each i such that $I'_i = 2$. Using relation (r-2) we can trade each of these $\mathcal{U}_i \mathcal{V}_i$ for a $\mathcal{U}_j \mathcal{V}_j$ where $I'_j = 0$. Proceeding in this manner, we may relate $\mathcal{U}^{I'} \mathcal{V}^{J'} X_{q-1}$ to some

$\mathcal{U}^{I''} \mathcal{V}^{J''} X_{q-1}$ where $|I''|_{L^1} = q - 1$ and $|I''|_{L^\infty} = 1$. The relations between X_{q-1} and X_q now show that

$$\mathcal{U}^{I'} \mathcal{V}^{J'} X_{q-1} = \mathcal{U}^{I''} \mathcal{V}^{J''} X_{q-1} = \mathcal{V}^J X_q,$$

completing the proof. $\blacksquare$

In general, a free resolution of the homology $\mathcal{HFL}(T(n, n))$ can be computed algorithmically (see [51]), or, for a concrete value of n , using a computer algebra system such as Macaulay2 [14].

7.3. The free complex of $T(3, 3)$

We present a free resolution of the torus link $T(3, 3)$. The homology of $T(3, 3)$ is generated by X_1, X_2, X_3 with the following relations:

$$\begin{aligned} \mathcal{U}_i X_1 &= \prod_{j \in \{1, 2, 3\} \setminus \{i\}} \mathcal{V}_j X_2, & \mathcal{V}_i X_3 &= \prod_{j \in \{1, 2, 3\} \setminus \{i\}} \mathcal{U}_j X_2, \\ \mathcal{U}_i \mathcal{V}_i X_k &= \mathcal{U}_j \mathcal{V}_j X_k. \end{aligned}$$

Then a free resolution of the homology is

$$0 \rightarrow C_3 \xrightarrow{\partial_3} C_2 \xrightarrow{\partial_2} C_1 \xrightarrow{\partial_1} C_0 \rightarrow 0,$$

with the spaces C_0, C_1, C_2, C_3 and the maps $\partial_1, \partial_2, \partial_3$ defined as follows.

The space $C_0 = \mathcal{R}_3^3$ is generated by X_1, X_2, X_3 . Take $C_1 = \mathcal{R}_3^8$ generated by $b_1, b_2, b_3, B_1, B_2, B_3$, and Z_1, Z_2 . For symmetry, it is helpful to consider an extra variable Z_3 which satisfies $Z_3 = Z_1 + Z_2$; Z_3 is not a generator of C_1 . The differential $\partial_1: C_1 \rightarrow C_0$ is given by

$$\begin{aligned} \partial_1 b_i &= \mathcal{U}_i X_1 + \prod_{j \in \{1, 2, 3\} \setminus \{i\}} \mathcal{V}_j X_2, & \partial_1 B_i &= \mathcal{V}_i X_3 + \prod_{j \in \{1, 2, 3\} \setminus \{i\}} \mathcal{U}_j X_2, \\ \partial_1 Z_1 &= \mathcal{U}_2 \mathcal{V}_2 X_2 + \mathcal{U}_3 \mathcal{V}_3 X_2, & \partial_1 Z_2 &= \mathcal{U}_1 \mathcal{V}_1 X_2 + \mathcal{U}_3 \mathcal{V}_3 X_2, \\ \partial_1 Z_3 &= \partial_1(Z_1 + Z_2) = \mathcal{U}_1 \mathcal{V}_1 X_2 + \mathcal{U}_2 \mathcal{V}_2 X_2. \end{aligned}$$

The link Floer homology of $T(3, 3)$ is coker ∂_1 . Indeed, the relations $\mathcal{U}_i \mathcal{V}_i X_k = \mathcal{U}_j \mathcal{V}_j X_k$ for $k = 1$ and $k = 3$ follow from the other relations. For example,

$$0 = \mathcal{V}_1(\mathcal{U}_1 X_1 - \mathcal{V}_2 \mathcal{V}_3 X_2) - \mathcal{V}_2(\mathcal{U}_2 X_1 - \mathcal{V}_1 \mathcal{V}_3 X_2) = (\mathcal{U}_1 \mathcal{V}_1 - \mathcal{U}_2 \mathcal{V}_2) X_1.$$

We define the module $C_2 = \mathcal{R}_3^6$ with generators $c_1, c_2, c_3, d_1, d_2, d_3$ and let ∂_2 be the differential

$$\begin{aligned} \partial_2 c_k &= \mathcal{U}_i b_j + \mathcal{U}_j b_i + \mathcal{V}_k Z_k, \\ \partial_2 d_k &= \mathcal{V}_i B_j + \mathcal{V}_j B_i + \mathcal{U}_k Z_k. \end{aligned}$$

Here $\{i, j, k\}$ ranges through all permutations of $\{1, 2, 3\}$. There is a relation between $c_1, \dots, d_3$. That is, there is a module $C_3 = \mathcal{R}_3$ generated by e with $\partial_3: C_3 \rightarrow C_2$ given by

$$\partial_3 e = \mathcal{U}_1 c_1 + \mathcal{U}_2 c_2 + \mathcal{U}_3 c_3 + \mathcal{V}_1 d_1 + \mathcal{V}_2 d_2 + \mathcal{V}_3 d_3.$$

It can be checked either directly or using a computer algebra system that $\ker \partial_1 = \text{im } \partial_2$, $\ker \partial_2 = \text{im } \partial_3$ and $\ker \partial_3 = 0$. That is, the complex we have constructed is a free resolution of $\text{coker } \partial_1$.

Remark 7.4. By examining the resolution of $\mathcal{HFL}(T(3, 3))$ shown in Figure 7.1, we see that the relations described in Lemma 7.1 have some redundancy. For example the relation $(\mathcal{U}_1 \mathcal{V}_1 + \mathcal{U}_2 \mathcal{V}_2) X_1 = 0$ is a consequence of the relations $\mathcal{U}_1 X_1 = \mathcal{V}_2 \mathcal{V}_3 X_2$ and $\mathcal{U}_2 X_2 = \mathcal{V}_1 \mathcal{V}_3 X_2$.




Fig. 7.1. The complex $\mathcal{CFL}(T(3, 3))$ as a free resolution.

7.4. The free complex of $T(4, 4)$

To stress the usefulness and power of Theorem 6.1, we show how to compute the link Floer chain complex of the torus link $T(4, 4)$. We start with the model of the link Floer homology for $T(4, 4)$ as described in Theorem 7.3 and compute its free resolution. It is pretty straightforward to find candidates for the relations between generators of $T(4, 4)$, and then candidates for the relations among relations (second syzygies), and to iterate this procedure. This amounts to creating a complex $(\mathcal{C}_i, \partial_i)$, whose homology at the zero

grading is $\mathcal{HFL}(T(4, 4))$. Showing that the relations are complete, that is, the complex we have constructed is acyclic in higher gradings, is rather tedious. We have used Macaulay [14] to verify this fact.

We can give a quick description of the free resolution of $T(4, 4)$. The free resolution has length 4:

$$0 \rightarrow \mathcal{C}_4 \xrightarrow{\partial_4} \mathcal{C}_3 \xrightarrow{\partial_3} \mathcal{C}_2 \xrightarrow{\partial_2} \mathcal{C}_1 \xrightarrow{\partial_1} \mathcal{C}_0 \rightarrow 0.$$

Let $\mathcal{C}_0$ be the free $\mathcal{R}_4$ -module generated by $X_1, \dots, X_4$. Consider the module $\mathcal{C}_1 \cong \mathcal{R}_4^{20}$ generated by Z_{ij}^k with $k = 2, 3$, $j = i + 1$, $a_1, \dots, a_4$, B_{ij} with $1 \leq i < j \leq 4$ and $A_1, \dots, A_4$. As in the $T(3, 3)$ case, for symmetry, we add variables Z_{ij}^k for $j > i + 1$, satisfying $Z_{ij}^k = Z_{i,i+1}^k + Z_{i+1,i+2}^k + \dots + Z_{j-1,j}^k$. If $j < i$ we set $B_{ij} := B_{ji}$. There is a map $\partial_1: \mathcal{C}_1 \rightarrow \mathcal{C}_0$ given by

$$\begin{aligned} \partial_1 Z_{i,i+1}^k &= (\mathcal{U}_i \mathcal{V}_i + \mathcal{U}_{i+1} \mathcal{V}_{i+1}) X_k, & \partial_1 a_i &= \mathcal{U}_i X_1 + \prod_{j \neq i} \mathcal{V}_j X_2, \\ \partial_1 B_{ij} &= \mathcal{U}_i \mathcal{U}_j X_2 + \prod_{\substack{i' < j' \\ \{i,j,i',j'\} = \{1,2,3,4\}}} \mathcal{V}_{i'} \mathcal{V}_{j'} X_3, & \partial_1 A_i &= \prod_{j \neq i} \mathcal{U}_j X_3 + \mathcal{V}_i X_4. \end{aligned}$$

By Theorem 7.3, the link Floer homology of $T(4, 4)$ is $\text{coker } \partial_1$. We now define the module $\mathcal{C}_2 \cong \mathcal{R}_4^{28}$. It is generated by a_{ij} , A_{ij} with $1 \leq i < j \leq 4$, and c_k^{ij} , C_{ij}^k .

The indexing of c and C generators is a bit complex. We choose $1 \leq k \leq 4$, and $\{i, j\}$ is a subset of $\{1, 2, 3, 4\} \setminus \{k\}$ with $j > i$. To obtain a smaller resolution, we reduce the number of generators by declaring that the c -generators are c_1^{23} and c_1^{34} , c_2^{13} and c_2^{34} , c_3^{12} and c_3^{24} , as well as c_4^{12} and c_4^{23} . Whenever another configuration of indices appears in the differential, we declare it to be the sum of the other two generators with the same subscript, like c_1^{24} is not a generator, but $c_1^{24} = c_1^{23} + c_1^{34}$ etc.

For C -generators, for each three-element subset $\{a, b, c\} \subseteq \{1, 2, 3, 4\}$ we choose two out of three permutations $\{i, j, k\}$ with $i < j$ and these indices are C -generators. The third object is declared to be the sum of the other two. More specifically, we choose C_{23}^1, C_{13}^2 to be generators and $C_{12}^3 = C_{23}^1 + C_{13}^2$, C_{24}^1, C_{12}^4 as generators, and $C_{14}^2 = C_{24}^1 + C_{12}^4$, C_{34}^1, C_{14}^3 as generators, with $C_{13}^4 = C_{34}^1 + C_{14}^3$. Finally, C_{34}^2, C_{24}^3 are generators, whereas $C_{23}^4 = C_{34}^2 + C_{24}^3$.

Therefore, we have eight c -generators and eight C -generators. The map $\partial_2: \mathcal{C}_2 \rightarrow \mathcal{C}_1$ is given by

$$\begin{aligned} \partial_2 a_{ij} &= \mathcal{U}_j a_i + \mathcal{U}_i a_j + \prod_{k \neq i,j} \mathcal{V}_k Z_{ij}^2, \\ \partial_2 A_{ij} &= \mathcal{V}_j A_i + \mathcal{V}_i A_j + \prod_{k \neq i,j} \mathcal{U}_k Z_{ij}^3, \\ \partial_2 C_{ij}^k &= \mathcal{U}_j B_{ik} + \mathcal{U}_i B_{jk} + \mathcal{V}_\ell Z_{ij}^3, \\ \partial_2 c_k^{ij} &= \mathcal{V}_i B_{ik} + \mathcal{V}_j B_{jk} + \mathcal{U}_k Z_{ij}^2, \end{aligned}$$

where $\ell = \{1, 2, 3, 4\} \setminus \{i, j, k\}$.

The module $\mathcal{C}_3 \cong \mathcal{R}_4^{14}$ is generated by $a_{ijk}, A_{ijk}, B'_{ij}$ where $1 \leq i < j < k \leq 4$ with the map $\partial_3: \mathcal{C}_3 \rightarrow \mathcal{C}_2$ given by

$$\begin{aligned}\partial_3 a_{ijk} &= \mathcal{U}_i a_{jk} + \mathcal{U}_j a_{ik} + \mathcal{U}_k a_{ij} + \mathcal{V}_\ell (\mathcal{V}_i c_i^{jk} + \mathcal{V}_j c_j^{ik} + \mathcal{V}_k c_k^{ij}), \\ \partial_3 A_{ijk} &= \mathcal{V}_i A_{jk} + \mathcal{V}_j A_{ik} + \mathcal{V}_k A_{ij} + \mathcal{U}_\ell (\mathcal{U}_i C_{jk}^\ell + \mathcal{U}_j C_{ik}^\ell + \mathcal{U}_k C_{ij}^\ell), \\ \partial_3 B'_{ij} &= \mathcal{V}_k C_{ij}^k + \mathcal{V}_\ell C_{ij}^\ell + \mathcal{U}_i c_j^{k\ell} + \mathcal{U}_j c_i^{k\ell},\end{aligned}$$

where we let $\ell = \{1, 2, 3, 4\} \setminus \{i, j, k\}$.

The module $\mathcal{C}_4 \cong \mathcal{R}_4^2$ is generated by d_{1234}, D_{1234} with the map ∂_4 given by

$$\begin{aligned}\partial_4(d_{1234}) &= \sum_i \mathcal{U}_i a_{jkl} + \sum_{i < j} \mathcal{V}_i \mathcal{V}_j B'_{ij}, \\ \partial_4(D_{1234}) &= \sum_i \mathcal{V}_i A_{jkl} + \sum_{\substack{i < j, k < \ell \\ \{i, j, k, \ell\} = \{1, 2, 3, 4\}}} \mathcal{U}_i \mathcal{U}_j B'_{kl}.\end{aligned}$$

In each of the two equations, where the sum runs over the indices i , we let $1 \leq j < k < \ell \leq 4$ be such that $\{i, j, k, \ell\}$ is a permutation of $\{1, 2, 3, 4\}$.

It can be verified that the above description of $\mathcal{CF}\mathcal{L}(T(4, 4))$ is acyclic except at resolution grading 0, with coker ∂_1 being the link Floer homology of $T(4, 4)$.

Acknowledgments. The authors would like to thank Eugene Gorsky, Chuck Livingston, András Némethi and Lorenzo Traldi for stimulating discussions. We are grateful to Marco Marengon for the help with the complex of the $T(3, 3)$ torus link. We have benefited a lot from Karol Palka's explanations of the state-of-the-art in computing resolutions of modules.

Funding. MB was supported by OPUS 2019/B/35/ST1/01120 grant of the Polish National Science Center. BL is partially supported by NSF grant DMS-2203237. IZ was partially supported by NSF grants DMS-1703685 and DMS-2204375.

References

- [1] Artin, M.: [On isolated rational singularities of surfaces](#). Amer. J. Math. **88**, 129–136 (1966) Zbl [0142.18602](#) MR [0199191](#)
- [2] Borodzik, M., Gorsky, E.: [Immersed concordances of links and Heegaard Floer homology](#). Indiana Univ. Math. J. **67**, 1039–1083 (2018) Zbl [1407.57005](#) MR [3820241](#)
- [3] Campillo, A., Delgado, F., Gusein-Zade, S. M.: [Are algebraic links in the Poincaré sphere determined by their Alexander polynomials?](#) Math. Z. **294**, 593–613 (2020) Zbl [1469.14008](#) MR [4050077](#)
- [4] Cavallo, A., Liu, B.: [Fibered and strongly quasi-positive L-space links](#). Michigan Math. J. **73**, 209–224 (2023) Zbl [1525.57002](#) MR [4555227](#)
- [5] Dai, I., Hom, J., Stoffregen, M., Truong, L.: [More concordance homomorphisms from knot Floer homology](#). Geom. Topol. **25**, 275–338 (2021) Zbl [1464.57019](#) MR [4226231](#)
- [6] Eisenbud, D., Neumann, W.: [Three-dimensional link theory and invariants of plane curve singularities](#). Ann. of Math. Stud. 110, Princeton University Press, Princeton, NJ (1985) Zbl [0628.57002](#) MR [0817982](#)
- [7] Friedl, S., Juhász, A., Rasmussen, J.: [The decategorification of sutured Floer homology](#). J. Topol. **4**, 431–478 (2011) Zbl [1270.57041](#) MR [2805998](#)

- [8] Gompf, R. E., Stipsicz, A. I.: [4-manifolds and Kirby calculus](#). Grad. Stud. Math. 20, American Mathematical Society, Providence, RI (1999) Zbl [0933.57020](#) MR [1707327](#)
- [9] Gorsky, E., Hom, J.: [Cable links and L-space surgeries](#). Quantum Topol. **8**, 629–666 (2017) Zbl [1422.57021](#) MR [3737274](#)
- [10] Gorsky, E., Liu, B., Moore, A. H.: [Surgery on links of linking number zero and the Heegaard Floer \$d\$ -invariant](#). Quantum Topol. **11**, 323–378 (2020) Zbl [1508.57015](#) MR [4118637](#)
- [11] Gorsky, E., Némethi, A.: [Lattice and Heegaard Floer homologies of algebraic links](#). Int. Math. Res. Notices **2015**, 12737–12780 Zbl [1342.57005](#) MR [3431635](#)
- [12] Gorsky, E., Némethi, A.: [Links of plane curve singularities are L-space links](#). Algebr. Geom. Topol. **16**, 1905–1912 (2016) Zbl [1364.14022](#) MR [3546454](#)
- [13] Grauert, H.: [Über Modifikationen und exzeptionelle analytische Mengen](#). Math. Ann. **146**, 331–368 (1962) Zbl [0173.33004](#) MR [0137127](#)
- [14] Grayson, D. R., Stillman, M. E.: Macaulay2, a software system for research in algebraic geometry. <http://www.math.uiuc.edu/Macaulay2/>, visited on October 28, 2024
- [15] Hartshorne, R.: [Algebraic geometry](#). Grad. Texts in Math. 52, Springer, New York (1977) Zbl [0367.14001](#) MR [0463157](#)
- [16] Hedden, M.: [On knot Floer homology and cabling. II](#). Int. Math. Res. Notices **2009**, 2248–2274 Zbl [1172.57008](#) MR [2511910](#)
- [17] Hendricks, K., Hom, J., Stoffregen, M., Zemke, I.: An involutive dual knot surgery formula. arXiv:[2205.12798](#) (2022)
- [18] Hilbert, D.: [Ueber die Theorie der algebraischen Formen](#). Math. Ann. **36**, 473–534 (1890) Zbl [22.0133.01](#) MR [1510634](#)
- [19] Kadeishvili, T. V.: The algebraic structure in the homology of an $A(\infty)$ -algebra. Soobshch. Akad. Nauk Gruzin. SSR **108**, 249–252 (1982) (in Russian) Zbl [0535.55005](#) MR [0720689](#)
- [20] Keller, B.: [Introduction to \$A\$ -infinity algebras and modules](#). Homology Homotopy Appl. **3**, 1–35 (2001) Zbl [0989.18009](#) MR [1854636](#)
- [21] Kontsevich, M., Soibelman, Y.: [Homological mirror symmetry and torus fibrations](#). In: Symplectic geometry and mirror symmetry (Seoul, 2000), World Scientific Publications, River Edge, NJ, 203–263 (2001) Zbl [1072.14046](#) MR [1882331](#)
- [22] Laufer, H. B.: Normal two-dimensional singularities. Ann. of Math. Stud. 71, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo (1971) Zbl [0245.32005](#) MR [0320365](#)
- [23] Lefèvre-Hasegawa, K.: Sur les A_∞ -catégories. arXiv:[math/0310337](#) (2003)
- [24] Lipshitz, R., Ozsváth, P. S., Thurston, D. P.: [Bimodules in bordered Heegaard Floer homology](#). Geom. Topol. **19**, 525–724 (2015) Zbl [1315.57036](#) MR [3336273](#)
- [25] Lipshitz, R., Ozsváth, P. S., Thurston, D. P.: [Bordered Heegaard Floer homology](#). Mem. Amer. Math. Soc. **254**, no. 1216, viii+279 pp. (2018) Zbl [1422.57080](#) MR [3827056](#)
- [26] Liu, B.: [Heegaard Floer homology of L-space links with two components](#). Pacific J. Math. **298**, 83–112 (2019) Zbl [1411.57029](#) MR [3910050](#)
- [27] Liu, B.: [L-space surgeries on 2-component L-space links](#). Trans. London Math. Soc. **8**, 65–94 (2021) Zbl [1501.57010](#) MR [4440446](#)
- [28] Liu, Y.: [L-space surgeries on links](#). Quantum Topol. **8**, 505–570 (2017) Zbl [1381.57019](#) MR [3692910](#)
- [29] Looijenga, E. J. N.: Isolated singular points on complete intersections. 2nd ed., Surv. Mod. Math. 5, International Press, Somerville, MA; Higher Education Press, Beijing (2013) Zbl [1272.14002](#) MR [3112993](#)
- [30] Manolescu, C., Ozsváth, P. S.: Heegaard Floer homology and integer surgeries on links. arXiv:[1011.1317](#) (2010)
- [31] Milnor, J.: Singular points of complex hypersurfaces. Ann. of Math. Stud. 61, Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo (1968) Zbl [0184.48405](#) MR [0239612](#)

- [32] Némethi, A.: Five lectures on normal surface singularities. In: Low dimensional topology (Eger, 1996/Budapest, 1998), Bolyai Soc. Math. Stud. 8, János Bolyai Mathematical Society, Budapest, 269–351 (1999) Zbl 0958.32026 MR 1747271
- [33] Némethi, A.: [On the Ozsváth–Szabó invariant of negative definite plumbed 3-manifolds](#). Geom. Topol. **9**, 991–1042 (2005) Zbl 1138.57301 MR 2140997
- [34] Némethi, A.: [Lattice cohomology of normal surface singularities](#). Publ. Res. Inst. Math. Sci. **44**, 507–543 (2008) Zbl 1149.14029 MR 2426357
- [35] Némethi, A.: [Links of rational singularities, L-spaces and LO fundamental groups](#). Invent. Math. **210**, 69–83 (2017) Zbl 1380.32027 MR 3698339
- [36] Némethi, A.: [Normal surface singularities](#). Ergeb. Math. Grenzgeb. (3) 74, Springer, Cham (2022) Zbl 1523.14001 MR 4510934
- [37] Némethi, A., Szilárd, Á.: [Milnor fiber boundary of a non-isolated surface singularity](#). Lecture Notes in Math. 2037, Springer, Heidelberg (2012) Zbl 1239.32024 MR 3024944
- [38] Neumann, W. D.: [A calculus for plumbing applied to the topology of complex surface singularities and degenerating complex curves](#). Trans. Amer. Math. Soc. **268**, 299–344 (1981) Zbl 0546.57002 MR 0632532
- [39] Neumann, W. D.: Invariants of plane curve singularities. In: Knots, braids and singularities (Plans-sur-Bex, 1982), Monogr. Enseign. Math. 31, Enseignement Mathématique, Genève, 223–232 (1983) Zbl 0586.14023 MR 0728588
- [40] Ozsváth, P., Stipsicz, A. I., Szabó, Z.: [Knots in lattice homology](#). Comment. Math. Helv. **89**, 783–818 (2014) Zbl 1309.57023 MR 3284295
- [41] Ozsváth, P., Stipsicz, A. I., Szabó, Z.: [A spectral sequence on lattice homology](#). Quantum Topol. **5**, 487–521 (2014) Zbl 1316.57023 MR 3317341
- [42] Ozsváth, P., Stipsicz, A. I., Szabó, Z.: [Knot lattice homology in L-spaces](#). J. Knot Theory Ramifications **25**, article no. 1650003, 24 pp. (2016) Zbl 1345.57016 MR 3449536
- [43] Ozsváth, P., Szabó, Z.: [On the Floer homology of plumbed three-manifolds](#). Geom. Topol. **7**, 185–224 (2003) Zbl 1130.57302 MR 1988284
- [44] Ozsváth, P., Szabó, Z.: [Holomorphic disks and knot invariants](#). Adv. Math. **186**, 58–116 (2004) Zbl 1062.57019 MR 2065507
- [45] Ozsváth, P., Szabó, Z.: [Holomorphic disks and three-manifold invariants: properties and applications](#). Ann. of Math. (2) **159**, 1159–1245 (2004) Zbl 1081.57013 MR 2113020
- [46] Ozsváth, P., Szabó, Z.: [Holomorphic disks and topological invariants for closed three-manifolds](#). Ann. of Math. (2) **159**, 1027–1158 (2004) Zbl 1073.57009 MR 2113019
- [47] Ozsváth, P., Szabó, Z.: [On knot Floer homology and lens space surgeries](#). Topology **44**, 1281–1300 (2005) Zbl 1077.57012 MR 2168576
- [48] Ozsváth, P., Szabó, Z.: [Holomorphic disks, link invariants and the multi-variable Alexander polynomial](#). Algebr. Geom. Topol. **8**, 615–692 (2008) Zbl 1144.57011 MR 2443092
- [49] Ozsváth, P., Szabó, Z.: [Link Floer homology and the Thurston norm](#). J. Amer. Math. Soc. **21**, 671–709 (2008) Zbl 1235.53090 MR 2393424
- [50] Ozsváth, P. S., Szabó, Z.: [Knot Floer homology and integer surgeries](#). Algebr. Geom. Topol. **8**, 101–153 (2008) Zbl 1181.57018 MR 2377279
- [51] Peeva, I.: [Graded syzygies](#). Algebra Appl. 14, Springer, London (2011) Zbl 1213.13002 MR 2560561
- [52] Pichon, A.: [Fibrations sur le cercle et surfaces complexes](#). Ann. Inst. Fourier (Grenoble) **51**, 337–374 (2001) Zbl 0971.32013 MR 1824957
- [53] Rasmussen, J. A.: Floer homology and knot complements. Ph.D. thesis, Harvard University (2003) MR 2704683
- [54] Weibel, C. A.: [An introduction to homological algebra](#). Cambridge Stud. Adv. Math. 38, Cambridge University Press, Cambridge (1994) Zbl 0797.18001 MR 1269324

-
- [55] Yamamoto, M.: [Classification of isolated algebraic singularities by their Alexander polynomials](#). *Topology* **23**, 277–287 (1984) [Zbl 0581.57002](#) [MR 0770564](#)
 - [56] Zariski, O.: [On the topology of algebroid singularities](#). *Amer. J. Math.* **54**, 453–465 (1932) [Zbl 58.0614.02](#) [MR 1507926](#)
 - [57] Zemke, I.: [Link cobordisms and absolute gradings on link Floer homology](#). *Quantum Topol.* **10**, 207–323 (2019) [Zbl 1478.57035](#) [MR 3950650](#)
 - [58] Zemke, I.: A general Heegaard Floer surgery formula. [arXiv:2308.15658](#) (2023)
 - [59] Zemke, I.: The equivalence of lattice and Heegaard Floer homology. *Duke Math. J.*, to appear; [arXiv:2111.14962](#)
 - [60] Zemke, I.: Bordered manifolds with torus boundary and the link surgery formula. [arXiv:2109.11520v5](#) (2023)